Carnegie Mellon University
Research Showcase

Department of Mathematical Sciences | Mellon College of Science

1-1-1969

Totally positive differential systems Binyamin Schwarz.

Binyamin Schwarz
Carnegie Mellon University

Follow this and additional works at: http://repository.cmu.edu/math

Recommended Citation
Schwarz, Binyamin, "Totally positive differential systems Binyamin Schwarz." (1969). *Department of Mathematical Sciences.* Paper 27.
http://repository.cmu.edu/math/27

This Technical Report is brought to you for free and open access by the Mellon College of Science at Research Showcase. It has been accepted for inclusion in Department of Mathematical Sciences by an authorized administrator of Research Showcase. For more information, please contact research-showcase@andrew.cmu.edu.

is TP iri (a,b) .if and only if the following two conditions hold:

(a) $a_{\pm j}(t) = 0$, $|i - j| \geq 2$, $i, j = 1, \ldots, n$, $a < t < b$.

(b) $a_{\pm, i+1}(t) \geq 0$, $a_{\pm+1, \pm}(t) \geq 0$, $i = 1, \ldots, n-1$, $a < t < b$.

Proof. As total positivity of the system (1.2) implies its positivity, it follows from Lemma 1 that all off-diagonal elements $a_{ij}(t)$, $i \neq j$, of $A(t)$ have to be non-negative in (a,b). If an element $a_{ij}(t)$, $|i - j| \geq 2$, were to be positive for some t part (ii) of Corollary 1 would imply that the matrix $B^{(2)}(t)$ of the second compound system has an off-diagonal element which is somewhere negative, and Lemma 1, applied to this second compound system, then shows that (1.2) is not TP. Conditions (a) and (b) are thus necessary. Their sufficiency follows from part (i) of Corollary 1 and the sufficiency part of Lemma 1, applied to all compound systems (1.3). (We remark that we also use that the pth compound of the unit matrix $I = (\delta_{ij})_1^n$ is again $I = (\delta_{ij})_1^N$. Hence if $Y(t) = Y(t,r)$ is the solution of (1.2) which satisfies (1.5), then its compound also satisfies $c_p(Y(r)) = I$.)

Theorem 3. Let the n^2 real functions $a_{ij}(t)$, $i,j = 1, \ldots, n$, be continuous in (a,b), $-\infty \leq a < b \leq \infty$, and set $A(t) = (a_{ij}(t))_1^n$. The differential system

$$(1.2) \qquad Y^*(t) = A(t)Y(t),$$