

7-2008

On the equivalence of conglomerability and disintegrability for unbounded random variables

Mark J. Schervish

Carnegie Mellon University, mark@stat.cmu.edu

Teddy Seidenfeld

Carnegie Mellon University, teddy@stat.cmu.edu

Joseph B. Kadane

Carnegie Mellon University, kadane@stat.cmu.edu

Follow this and additional works at: <http://repository.cmu.edu/statistics>

 Part of the [Statistics and Probability Commons](#)

Published In

Statistical Methods and Applications, 23, 4, 501-518.

This Technical Report is brought to you for free and open access by the Dietrich College of Humanities and Social Sciences at Research Showcase @ CMU. It has been accepted for inclusion in Department of Statistics by an authorized administrator of Research Showcase @ CMU. For more information, please contact research-showcase@andrew.cmu.edu.

ON THE EQUIVALENCE OF CONGLOMERABILITY AND DISINTEGRABILITY FOR UNBOUNDED RANDOM VARIABLES

BY MARK J. SCHERVISH, TEDDY SEIDENFELD, AND JOSEPH B.
KADANE,

Carnegie Mellon University

We extend a result of Dubins [3] from bounded to unbounded random variables. Dubins [3] showed that a finitely additive expectation over the collection of bounded random variables can be written as an integral of conditional expectations (disintegrability) if and only if the marginal expectation is always within the smallest closed interval containing the conditional expectations (conglomerability). We give a sufficient condition to extend this result to the collection \mathcal{Z} of all random variables that have finite expected value and whose conditional expectations are finite and have finite expected value. The sufficient condition also allows the result to extend some, but not all, subcollections of \mathcal{Z} . We give an example where the equivalence of disintegrability and conglomerability fails for a subcollection of \mathcal{Z} that still contains all bounded random variables.

1. Introduction. Dubins [3] establishes an equivalence between conglomerability in a partition and disintegrability in that partition for coherent previsions over bounded random variables. Loosely speaking (see Definition 3 for precise definitions) conglomerability in a partition means that the marginal prevision of each random variable lies between the smallest and largest of all of the conditional previsions given elements of the partition. Disintegrability means that the integral of the conditional prevision equals the marginal prevision.

Specifically, let Ω be a fixed non-empty set, π a partition of Ω , and \mathcal{X} the collection of bounded, real-valued random variables defined on Ω . The concept of coherent prevision on a collection of random variables was introduced by de Finetti [2].

DEFINITION 1. *Let \mathcal{P} be a collection of random variables defined on Ω , and let $P : \mathcal{P} \rightarrow \mathbb{R}$ be a prevision. We say that P is incoherent if there exists a finite subset of \mathcal{P} , $\{X_1, \dots, X_n\}$ and scalars $\alpha_1, \dots, \alpha_n$ and $\epsilon > 0$*

AMS 2000 subject classifications: Primary 60A05; secondary 28C05

Keywords and phrases: finite additivity, law of total probability, Daniell integral, coherence, ultrafilters

such that, for all $\omega \in \Omega$,

$$(1) \quad \sum_{i=1}^n \alpha_i [X_i(\omega) - P(X_i)] < -\epsilon.$$

If P is not incoherent, we say that P is coherent.

Given an arbitrary set Ω and the collection \mathcal{X} of all bounded random variables defined on Ω , it is always possible to find a coherent prevision P defined on \mathcal{X} . However, this existence result, like many others in this domain, relies on the Axiom of Choice. We avail ourselves of the Axiom of Choice whenever it is needed in this paper. Of course, there are many coherent previsions on \mathcal{X} , but each of them will be a finitely additive probability when restricted to the collection of indicator functions of subsets of Ω . That is, using the standard notation of letting the name of an event stand for its indicator function, $P(\Omega) = 1$, $P(A \cup B) = P(A) + P(B)$ when $A \cap B = \emptyset$, and $P(A) \geq 0$ for all $A \subseteq \Omega$. By finite additivity and linearity of prevision, the prevision of a simple function $X = \sum_{i=1}^n \alpha_i A_i$ equals $\sum_{i=1}^n \alpha_i P(A_i)$. This resembles the formula for the integral of a simple function in the usual measure theoretic derivation. To carry the resemblance further, the values of P on \mathcal{X} are uniquely determined from the finitely additive probability by means of the fact that

$$(2) \quad P(X) = \sup_{\text{simple } Y \leq X} P(Y) = \inf_{\text{simple } Y \geq X} P(Y).$$

The first equation in (2) is the same way that the Lebesgue integral of a nonnegative function X is defined in terms of the integrals of simple functions. Indeed P can be expressed as a finitely additive integral. For each $X \in \mathcal{X}$, we denote $P(X) = \int_{\Omega} X(\omega) P(d\omega)$. Generalizing from the definition of Daniell integral [6, Chapter 13], we can call a prevision a finitely additive Daniell integral.

DEFINITION 2. *Let \mathcal{L} be a linear space of functions that contains all constants, and let L be a linear functional defined on \mathcal{L} that satisfies $L(X) \geq 0$ whenever $X(\omega) \geq 0$ for all ω . Then L is called a nonnegative linear functional or a finitely additive Daniell integral over \mathcal{L} . We can write $L(X) = \int_{\Omega} X(\omega) L(d\omega)$.*

Finitely additive Daniell integrals behave in many ways like the countably additive Lebesgue integral. One property that the two integrals share is the following transformation property that we use.

LEMMA 1. *Let \mathcal{L} be a linear space of functions on Ω , and let $H : \Omega \rightarrow \Theta$ be a function. Let L be a finitely additive Daniell integral over \mathcal{L} . Let \mathcal{W} be a linear space of functions on Θ that includes all constants and such that, $Z(H) \in \mathcal{L}$ for every $Z \in \mathcal{W}$. Define $L_H(Z) = L(X)$, where $X = Z(H)$. Then L_H is a finitely additive Daniell integral on \mathcal{W} that we call the integral induced from L by H .*

PROOF. If $Z_1, Z_2 \in \mathcal{W}$ and $\alpha, \beta \in \mathbb{R}$, then

$$\begin{aligned} L_H(\alpha Z_1 + \beta Z_2) &= L[\alpha Z_1(H) + \beta Z_2(H)] = \alpha L[Z_1(H)] + \beta L[Z_2(H)] \\ &= \alpha L_H(Z_1) + \beta L_H(Z_2). \end{aligned}$$

If $Z(\theta) \geq 0$ for all $\theta \in \Theta$, then $X(\omega) = Z(H(\omega)) \geq 0$ for all $\omega \in \Omega$. Hence $L_H(Z) = L(X) \geq 0$. \square

In terms of integrals, Lemma 1 says that

$$(3) \quad \int_{\Theta} Z(\theta) L_H(d\theta) = \int_{\Omega} Z(H(\omega)) L(d\omega).$$

Extending a finite coherent prevision P on a set of random variables \mathcal{P} to a finitely additive Daniell integral is straightforward.

LEMMA 2. *Let P be a coherent prevision on a set \mathcal{P} of random variables such that $P(X)$ is finite for all $X \in \mathcal{P}$. Then there exists a finitely additive Daniell integral L on a linear space \mathcal{L} that contains \mathcal{P} such that $L(X) = P(X)$ for every $X \in \mathcal{P}$.*

PROOF. Let \mathcal{L} be the linear span of \mathcal{P} . For each $Y \in \mathcal{L}$, express Y as $\sum_{i=1}^n \alpha_i X_i$ with $X_1, \dots, X_n \in \mathcal{P}$. Define $L(Y) = \sum_{i=1}^n \alpha_i P(X_i)$. To see that L is well-defined, assume that Y can be expressed two different ways as

$$Y = \sum_{i=1}^n \alpha_i X_i = \sum_{j=1}^m \beta_j Y_j.$$

Let $\ell_1 = \sum_{i=1}^n \alpha_i P(X_i)$ and $\ell_2 = \sum_{j=1}^m \beta_j P(Y_j)$. If $\ell_1 > \ell_2$, then for all ω

$$\sum_{i=1}^n \alpha_i [X_i(\omega) - P(X_i)] - \sum_{j=1}^m \beta_j [Y_j(\omega) - P(Y_j)] = \ell_2 - \ell_1 < 0.$$

This would make P incoherent, a contradiction. A similar contradiction arises if $\ell_1 < \ell_2$. So L is well-defined. It is straightforward to see that L is

linear. To see that $X \geq 0$ implies $L(X) \geq 0$, suppose to the contrary that $L(X) = -\epsilon < 0$. Write $X = \sum_{i=1}^n \alpha_i X_i$. Then, for all ω ,

$$\sum_{i=1}^n (-\alpha_i)[X_i - P(X_i)] = -X + L(X) \leq -\epsilon,$$

which is a contradiction to P being coherent. \square

In view of Lemma 2, there is no loss of generality in assuming that every finite coherent prevision is defined on a linear space. For the remainder of this paper, we make that assumption.

For each coherent prevision P one can construct a family of coherent conditional previsions $P(X|h)$, defined for each pair, (X, h) with $X \in \mathcal{X}$ and h a nonempty subset of Ω . Coherence places only one restriction on $P(X|h)$, namely that

$$(4) \quad P(h)P(X|h) = P(hX),$$

when the previsions are finite. If $P(h) = 0$ and all previsions are finite, this restriction is vacuous, and $P(X|h)$ is completely arbitrary. de Finetti [2, Section 4.2] says that it is also necessary for

$$(5) \quad \inf_{\omega \in h} X(\omega) \leq P(X|h) \leq \sup_{\omega \in h} X(\omega).$$

If $P(h) > 0$ and the previsions are coherent, then (5) will hold. Although (5) is a useful property for conditional previsions to have in general, it is not required in order to avoid sure loss in de Finetti's sense when $P(h) = 0$. If one is going to impose additional constraints on conditional previsions when $P(h) = 0$ so that they behave more like the cases in which $P(h) > 0$, it might make more sense to assume something along the following lines.

ASSUMPTION 1. *Let h be a nonempty subset of Ω and let \mathcal{L} be a linear space of random variables that includes h . Then*

- $P(\cdot|h)$ is a linear functional on \mathcal{L} ,
- $P(h|h) = 1$, and
- If $X \in \mathcal{L}$ and $X(\omega) \geq 0$ for all $\omega \in h$, then $P(X|h) \geq 0$.

We next prove some useful implications of Assumption 1 including that Assumption 1 implies (5).

LEMMA 3. *Let \mathcal{L} be a linear space of random variables. Let h be a nonempty set such that $h \in \mathcal{L}$. Let $P(\cdot|h)$ be a conditional prevision on \mathcal{L} that satisfies Assumption 1. Let $X \in \mathcal{L}$.*

- If $Xh \in \mathcal{L}$, then $P(Xh|h) = P(X|h)$.
- (5) holds.
- Let z be a real constant. If $X(\omega) = z$ for all $\omega \in h$, then $P(X|h) = z$.

PROOF. For the first claim, let $Y = X - Xh$. Then $Y(\omega) = 0$ for all $\omega \in h$. It follows from the last bullet in Assumption 1 that $P(Y|h) \geq 0$ and $P(Y|h) \leq 0$, hence $P(Y|h) = 0$. By the first bullet in Assumption 1, $P(Xh|h) = P(X|h)$. For the second claim, we prove only the first inequality in (5) as the second follows from applying the first inequality with $-X$ instead of X . If $\inf_{\omega \in h} X(\omega) = -\infty$, the first inequality in (5) is immediate. Because X is real valued, $\inf_{\omega \in h} X(\omega) < \infty$. So, assume that $\inf_{\omega \in h} X(\omega)$ is finite, and call the value z . Then $X(\omega) - zh(\omega) \geq 0$ for all $\omega \in h$. The last bullet in Assumption 1 implies that $P(X - zh|h) \geq 0$. The first and second bullets imply that

$$(6) \quad P(X|h) \geq P(zh|h) = zP(h|h) = z.$$

For the third claim, notice that $Xh = zh \in \mathcal{L}$. Then apply the first claim together with the last two equalities in (6). \square

Dubins [3] relies on Assumption 1 in proving his equivalence between conglomerability and disintegrability. For the remainder of this paper, we assume that Assumption 1 holds for every conditional prevision. The following result shows that, if one starts with coherent previsions, then coherent conditional previsions that satisfy Assumption 1 can be constructed simultaneously for each set h in a partition π .

LEMMA 4. *Assume that P is a finite coherent prevision on a linear space \mathcal{L} of random variables defined on Ω . Let π be a partition of Ω into nonempty sets. For every $h \in \pi$, there exists $P(\cdot|h)$ that satisfies Assumption 1.*

PROOF. Let π be a partition of Ω into nonempty events. For all $h \in \pi$ such that $P(h) = 0$, let $\omega_h \in h$, and define $P(X|h) = X(\omega_h)$ for all $X \in \mathcal{L}$. This clearly satisfies Assumption 1. For those h with $P(h) > 0$, we need to ensure both (4) and Assumption 1. Construct the collection \mathcal{D} of all random variables of the form Xh where $X \in \mathcal{L}$, $h \in \pi$ has $P(h) > 0$, and such that $Xh \notin \mathcal{L}$. If \mathcal{D} is nonempty, well order \mathcal{D} . The fundamental theorem of prevision (e.g., [7, Proposition 6]) allows one to extend a coherent prevision defined on an arbitrary collection, one random variable at a time, to a coherent prevision on a larger collection. Using transfinite induction, we can extend P to $\mathcal{L} \cup \mathcal{D}$ as follows. At each successor ordinal, the fundamental

theorem applies directly. At each limit ordinal, coherence places requirements on finitely many previsions at a time. Since all finite collections of previsions were verified as coherent at earlier stages of the induction, the entire collection is coherent at each limit ordinal. Next, use Lemma 2 to extend P to the linear span of $\mathcal{L} \cup \mathcal{D}$. Finally, regardless of whether or not \mathcal{D} was empty, for each $X \in \mathcal{L}$, define $P(X|h) = P(Xh)/P(h)$, which satisfies Assumption 1. \square

Although the conditional previsions constructed in Lemma 4 satisfy Assumption 1, they are not necessarily the only ones that do so. For the remainder of the paper we assume that, for the partition π under consideration and for each $h \in \pi$, $P(\cdot|h)$ satisfies Assumption 1.

2. Conglomerability and disintegrability. We turn now to precise definitions of conglomerability and disintegrability. Let π be a partition of Ω . That is, π is a collection $\{h : h \in \pi\}$ of mutually disjoint subsets of Ω such that their union is Ω . The conditional prevision of each random variable X given each element h of π is $P(X|h)$. In order to make sense out of the loose phrase “the integral of the conditional prevision equals the marginal prevision,” we need to be precise about what it means to integrate the conditional prevision. We take the approach of Lemma 1. Let $H : \Omega \rightarrow \pi$ be the function defined by $H(\omega)$ equal to that unique $h \in \pi$ such that $\omega \in h$. Let P_H denote the finitely additive Daniell integral induced from P by H . For each function $Z : \pi \rightarrow \mathbb{R}$ defined on π , we use (3) to write

$$(7) \quad \int_{\pi} Z(h)P_H(dh) = \int_{\Omega} Z(H(\omega))P(d\omega) = P[Z(H)].$$

For a random variable X such that $P(X|h)$ is finite for all $h \in \pi$, let $P(X|h)$ be $Z(h)$ in (7). Then (7) becomes

$$(8) \quad \int_{\pi} P(X|h)P_H(dh) = \int_{\Omega} P(X|H(\omega))P(d\omega) = P[P(X|H)].$$

Then (8) is what we mean by the integral of the conditional prevision.

In order to avoid cases in which previsions or conditional previsions are infinite, we deal only with random variables that satisfy the following assumption.

ASSUMPTION 2. *Let $X : \Omega \rightarrow \mathbb{R}$ and let π be a partition of Ω .*

1. $P(|X|) < \infty$,
2. for each $h \in \pi$, $P(X|h)$ is finite, and

3. $P[P(X|H)]$ is finite.

The set \mathcal{Z} of all X that satisfy Assumption 2 is a linear space. Also, if $X \in \mathcal{Z}$ then $P(X|H) \in \mathcal{Z}$.

DEFINITION 3. Let π be a partition of Ω . Let \mathcal{W} be a collection of random variables defined on Ω and that satisfy Assumption 2. P is conglomerable in π with respect to \mathcal{W} if, for each $X \in \mathcal{W}$

$$(9) \quad \inf_{h \in \pi} P(X|h) \leq P(X) \leq \sup_{h \in \pi} P(X|h).$$

P is disintegrable in π with respect to \mathcal{W} if, for each $X \in \mathcal{W}$,

$$(10) \quad \int_{\pi} P(X|h)P_H(dh) = \int_{\Omega} X(\omega)P(d\omega).$$

In view of (8), there is an alternative way to express that P is disintegrable in a partition.

PROPOSITION 1. P is disintegrable in π with respect to \mathcal{W} if and only if, for each $X \in \mathcal{W}$ $P(X) = P[P(X|H)]$.

Dubins [3, Theorem 1] shows that, with respect to the collection \mathcal{X} of bounded random variables, P is conglomerable in π if and only if P is disintegrable in π . The purpose of this note is to extend the equivalence of conglomerability and disintegrability to larger collections of random variables (such as \mathcal{Z}) that satisfy Assumption 2.

Finally, we demonstrate that the equivalence of conglomerability and disintegrability may fail for a collection \mathcal{Y} of random variables that are bounded below, where $\mathcal{X} \subset \mathcal{Y} \subset \mathcal{Z}$. In Section 4, we produce a coherent prevision P and partition π where P is conglomerable in π with respect to \mathcal{Y} , yet where P fails to be disintegrable in π with respect to \mathcal{Y} . In this case, the collection \mathcal{Y} is not a linear space, e.g., when $Y \in \mathcal{Y}$ is an unbounded random variable, then $-Y \notin \mathcal{Y}$.

Readers of [3] will note that the definition of conglomerable in Definition 3 looks different from the corresponding definition of Dubins. Specifically, Dubins' definition is that P is conglomerable in π with respect to the collection \mathcal{W} if

$$(11) \quad \text{for all } X \in \mathcal{W}, P(X|h) \geq 0 \text{ for all } h \in \pi \text{ implies } P(X) \geq 0.$$

Definition 3 is a straightforward generalization of the definition that [2, p. 143] gives for indicators of events. Definition 3 and (11) are equivalent

when $\mathcal{W} = \mathcal{X}$, the collection of all bounded random variables. The proof relies on the fact that \mathcal{X} is a linear space and contains all constants. The two definitions are not necessarily equivalent for every collection that is not a linear space and/or does not contain all constants. Here is an example taken from [3] and discussed further in [5].

EXAMPLE 1. Let $\Omega = \{0, 1\} \times \{1, 2, \dots\}$. Let P be a finitely additive probability such that $P((1, k)) = 2^{-k-1}$ for $k = 1, 2, \dots$, and $P(\{(0, 1), (0, 2), \dots\}) = 1/2$ but $P((0, k)) = 0$ for $k = 1, 2, \dots$. Let \mathcal{W} be the collection of all non-negative random variables. Let $\pi = \{h_1, h_2, \dots\}$ where $h_k = \{(0, k), (1, k)\}$. Because each $X \in \mathcal{W}$ is nonnegative, it follows that $P(X|h) \geq 0$ for all h and $P(X) \geq 0$. Hence, (11) holds. On the other hand, let $X((i, k)) = i$ for all $i = 0, 1$ and $k = 1, 2, \dots$. Then $P(X|h_k) = 1$ for all k while $P(X) = 1/2$ and P is not conglomerable by Definition 3.

In order to maintain the spirit of Dubins' definition when \mathcal{W} is not a linear space or does not contain all constants, we need to strengthen (11).

DEFINITION 4. Let \mathcal{W} be a collection of random variables. Let π be a partition, and let P be a coherent prevision on \mathcal{W} . We say that P is D-conglomerable in π with respect to \mathcal{W} if the following is true. For all $X \in \mathcal{W}$ and all real c ,

- $P(X|h) \leq c$ for all $h \in \pi$ implies $P(X) \leq c$, and
- $P(X|h) \geq c$ for all $h \in \pi$ implies $P(X) \geq c$.

We now show that Definition 4 is equivalent to conglomerability from Definition 3.

LEMMA 5. Let \mathcal{W} be a collection of random variables. Let π be a partition, and let P be a coherent prevision on \mathcal{W} . Then P is conglomerable in π with respect to \mathcal{W} if and only if P is D-conglomerable in π with respect to \mathcal{W} .

PROOF. For the "if" direction, suppose that P is D-conglomerable in π with respect to \mathcal{W} . Let $X \in \mathcal{W}$, and let $c_1 = \inf_{h \in \pi} P(X|h)$ and $c_2 = \sup_{h \in \pi} P(X|h)$. If c_1 is finite, then $P(X|h) \geq c_1$ for all $h \in \pi$ and Definition 4 says that $P(X) \geq c_1$. If c_2 is finite, then $P(X|h) \leq c_2$ for all $h \in \pi$ so that $P(X) \leq c_2$. If $c_1 = -\infty$, then $c_1 \leq P(X)$ is obvious. Similarly, if $c_2 = \infty$, then $P(X) \leq c_2$ is obvious. The only remaining cases are when $c_1 = \infty$ or $c_2 = -\infty$. Suppose that $c_1 = \infty$. Then $c_2 = \infty$ also. We need to show that $P(X) = \infty$. Because $P(X|h) \geq c$ for all real c and all $h \in \pi$, Definition 4

implies that $P(X) \geq c$ for all $c > 0$. Hence $P(X) = \infty$. Similarly, if $c_2 = -\infty$ then $c_1 = -\infty$ also and $P(X|h) \leq c$ for all real c and $h \in \pi$, Definition 4 implies that $P(X) \leq c$ for all real c . Hence $P(X) = -\infty$.

For the “only if” direction, suppose that P is conglomerable in π with respect to \mathcal{W} . Let $X \in \mathcal{W}$. Then Definition 3 implies that

$$\inf_{h \in \pi} P(X|h) \leq P(X) \leq \sup_{h \in \pi} P(X|h).$$

Let $c \in \mathbb{R}$. If $P(X|h) \geq c$ for all $h \in \pi$, then $c \leq \inf_{h \in \pi} P(X|h) \leq P(X)$. Similarly, if $P(X|h) \leq c$ for all $h \in \pi$, then $c \geq \sup_{h \in \pi} P(X|h) \geq P(X)$. \square

3. Extending the equivalence of conglomerability and disintegrability to unbounded variables. Lemma 6 shows that, under Assumption 2, disintegrability implies conglomerability for arbitrary collections.

LEMMA 6. *Let \mathcal{W} be a collection of random variables each of which satisfies Assumption 2. Let π be a partition, and let P be a coherent prevision on \mathcal{W} such that P is disintegrable in π with respect to \mathcal{W} . Then P is conglomerable in π with respect to \mathcal{W} .*

PROOF. Let $H : \Omega \rightarrow \pi$ be defined by $H(\omega)$ equal to the unique $h \in \pi$ such that $\omega \in h$. Let $X \in \mathcal{W}$, and define X' by $X'(\omega) = P(X|H(\omega))$. Then we have defined $P[P(X|H)]$ to be $P(X')$. Because P is coherent,

$$(12) \quad \inf_{\omega \in \Omega} X'(\omega) \leq P(X') \leq \sup_{\omega \in \Omega} X'(\omega).$$

From the definition of X' , we see that

$$(13) \quad \inf_{\omega \in \Omega} X'(\omega) = \inf_{h \in \pi} P(X|h), \text{ and } \sup_{\omega \in \Omega} X'(\omega) = \sup_{h \in \pi} P(X|h).$$

Combining (12) and (13) we get that (9) holds. Since the above argument applies to all $X \in \mathcal{W}$, P is conglomerable in π with respect to \mathcal{W} . \square

In light of Lemma 6, for each partition π and each coherent prevision P , every set \mathcal{W} of random variables satisfying Assumption 2 falls into one of three classes.

DEFINITION 5. *Let P be a coherent prevision, and let π be a partition. Let \mathcal{W} be a collection of random variables that have previsions under P . We say that*

- \mathcal{W} is of Class 0 relative to P and π if P is neither conglomerable nor disintegrable in π with respect to \mathcal{W} .
- \mathcal{W} is of Class 1 relative to P and π if P is conglomerable in π with respect to \mathcal{W} but P is not disintegrable in π with respect to \mathcal{W} .
- \mathcal{W} is of Class 2 relative to P and π if P is both conglomerable and disintegrable in π with respect to \mathcal{W} .

Dubins [3, Theorem 1] can be reexpressed as saying that for each partition π and each coherent prevision P the class \mathcal{X} of bounded random variables is either of Class 0 or of Class 2 but never of Class 1 relative to P and π . In Section 4, we give an example of a coherent prevision P , a partition π , and a collection \mathcal{Y} of random variables such that $\mathcal{X} \subset \mathcal{Y}$ and \mathcal{Y} is of Class 1 relative to P and π . The following result is a straightforward consequence of the class definitions.

PROPOSITION 2. *Let P be a coherent prevision that satisfies Assumption 1, and let π be a partition. Let \mathcal{W} be a collection of random variables that satisfy Assumption 2. If \mathcal{W} is of Class 0 relative to P and π , then every superset of \mathcal{W} is also of Class 0. If \mathcal{W} is of Class 2 relative to P and π , then every subset of \mathcal{W} is also of Class 2.*

Our extension of Dubins' theorem gives a sufficient condition for a collection \mathcal{W} of random variables to not be of Class 1.

THEOREM 1. *Let P be a coherent prevision that satisfies Assumption 1. Let π be a partition of Ω , and let $H : \Omega \rightarrow \pi$ be defined by $H(\omega)$ equal to that unique $h \in \pi$ such that $\omega \in h$. Let \mathcal{W} be a set of real-valued random variables defined on Ω that satisfy Assumption 2. Finally, assume that \mathcal{W} satisfies the following condition:*

$$(14) \quad \text{for every } X \in \mathcal{W}, X - P(X|H) \in \mathcal{W}.$$

Then, with respect to the collection \mathcal{W} , P is conglomerable in π if and only if P is disintegrable in π .

PROOF. Let P be a coherent prevision over the collection \mathcal{W} . We show first that P is both conglomerable and disintegrable in the finest partition Ω with respect to \mathcal{W} . Let $\pi = \Omega$. To see that P is conglomerable in Ω , let $X \in \mathcal{W}$. Assumption 1 implies that $P(X|\omega) = X(\omega)$ and, by coherence of the unconditional prevision P ,

$$\inf_{\omega \in \Omega} X(\omega) \leq P(X) \leq \sup_{\omega \in \Omega} X(\omega).$$

To see that P is disintegrable in Ω , note that $H(\omega) = \{\omega\}$ for all ω , and $P(X|H) = X$ for all $X \in \mathcal{W}$. Hence, we have $P[P(X|H)] = P(X)$, and P is both conglomerable and disintegrable in Ω with respect to \mathcal{W} .

Next, let π be a general partition. By the third claim in Lemma 3, $P[P(X|H)|h] = P(X|h)$ for all $h \in \pi$ and all $X \in \mathcal{W}$. By linearity of conditional prevision, it follows that, for each $h \in \pi$ and $X \in \mathcal{W}$,

$$P[X - P(X|H)|h] = 0.$$

Hence

$$\inf_{h \in \pi} P[X - P(X|H)|h] = 0 = \sup_{h \in \pi} P[X - P(X|H)|h].$$

We have assumed that $X - P(X|H) \in \mathcal{W}$. If P is conglomerable in π , then $P[X - P(X|H)] = 0$, from which it follows that $P(X) = P[P(X|H)]$, so that P is disintegrable in π .

If P is disintegrable in π then Lemma 6 shows that P is conglomerable in π . \square

It is easy to see that the collection \mathcal{Z} of all random variables that satisfy Assumption 2 satisfies the conditions of Theorem 1, and hence is not of Class 1. The key assumption in Theorem 1 is (14). For an arbitrary collection \mathcal{W} that satisfies Assumption 2, define

$$\begin{aligned} \mathcal{W}_- &= \{X - P(X|H) : X \in \mathcal{W}\}, \\ \mathcal{W}_+ &= \mathcal{W} \cup \mathcal{W}_-. \end{aligned}$$

The following results (the second of which is trivial) help to distinguish some collections of random variables by their class.

LEMMA 7. *Let P be a coherent prevision and π a partition. Let H be as defined in Theorem 1. Let \mathcal{W} be a collection of random variables that satisfy Assumption 2. Then*

1. \mathcal{W}_+ satisfies (14),
2. \mathcal{W} is of Class 2 relative to P and π if and only if \mathcal{W}_+ is also of Class 2, and
3. if \mathcal{W} is not of Class 2 relative to P and π , then \mathcal{W}_+ is of Class 0.

PROOF. For part (i), let $X \in \mathcal{W}$ so that $X - P(X|H) \in \mathcal{W}_+$. Also $P[X - P(X|H)|H]$ is identically 0, hence

$$X - P(X|H) - P[X - P(X|H)] \in \mathcal{W}_+.$$

For part (ii), the “if” direction is immediate from Proposition 2. For the “only if” direction, note that for every $Y \in \mathcal{W}_-$ $P(Y|H)$ is identically 0 and $P(Y) = 0$ if \mathcal{W} is of Class 2. For part (iii), Theorem 1 says that \mathcal{W}_+ is either of Class 0 or Class 2. If \mathcal{W} is not of Class 2, then no superset of it, such as \mathcal{W}_+ , can be of Class 2. Hence \mathcal{W}_+ must be of Class 0. \square

PROPOSITION 3. *If P is a countably additive prevision on a class \mathcal{W} of random variables and every element of π has positive probability, then \mathcal{W} is of Class 2 relative to P and π .*

One subtle point concerning Proposition 3 is that P can be a countably additive prevision on the collection of all bounded random variables but fail to be countably additive on a collection that includes unbounded random variables. The example in Section 4 has this property.

4. An example in which the equivalence of conglomerability and disintegrability fails. The following construction is a modification of an example given in [3, Theorem 2]. Let $\Omega = \{(i, k) : i = 1, 2, \dots, k = i, i + 1, \dots\}$ be the set of ordered pairs of positive integers in which the first coordinate is greater than or equal to the second. We use the algebra $\mathcal{A} = 2^\Omega$. Define a countably additive probability P on Ω by $P(\{(i, k)\}) = 2^{-k}$ for each $(i, k) \in \Omega$. It follows that \emptyset is the only subset of Ω with 0 probability. Since P is countably additive over \mathcal{A} , it is also countably additive on the set \mathcal{X} of bounded random variables defined on Ω . By Proposition 3, \mathcal{X} is of Class 2 relative to P and every partition π that we choose to consider.

Next, we extend P to a collection \mathcal{Y} of random variables that are bounded below in such a way that $P(|Y|) < \infty$ for each $Y \in \mathcal{Y}$. Part of the extension relies on the expectation operator

$$E(Y) = \sum_{(i,k) \in \Omega} P(\{(i, k)\})Y((i, k)),$$

which is well-defined for all Y that are bounded below. Also, $E(X) = P(X)$ for all $X \in \mathcal{X}$. As [2, Section 3.12] showed, if Y is unbounded above, it is possible that $P(Y) > E(Y)$. Indeed, unless there exists at least one unbounded Y whose prevision differs from its expectation, P will be countably additive on the collection of all random variables with finite prevision and hence will be both conglomerable and disintegrable in every partition. We call the difference $\beta(Y) = P(Y) - E(Y)$ the *boost function*. (See [8] for a more detail about the boost function.) We have that $\beta(X) = 0$ for every bounded X , and β is a linear functional that is nonnegative for every random variable bounded below.

We begin by extending P to a particular unbounded random variable W , and then to a linear space \mathcal{L} including W and all of \mathcal{X} . We then let \mathcal{Y} be the subcollection of \mathcal{L} consisting of the random variables that are bounded below. The starting random variable is $W(i, k) = k$. We set $P(W) = 14 = E(W) + 10$, so that $\beta(W) = 10$. In order to find a partition in which P is conglomerable but not disintegrable, we need to extend P beyond the linear span of \mathcal{X} and $\{W\}$. We do this by defining β on a larger collection of random variables.

DEFINITION 6. *Let Ω be a non-empty set. A collection p of subsets of Ω is called an ultrafilter if the following conditions hold:*

- *For every subset $A \subseteq \Omega$, either $A \in p$ or $A^C \in p$, but not both.*
- *If $A, B \in p$, then $A \cap B \in p$.*
- *If $A \in p$ and $A \subseteq B$, then $B \in p$.*

The simplest ultrafilters are the *principal* ultrafilters that consist of all subsets that contain a specific element of Ω . All other ultrafilters are called *non-principal*. A proof of the existence of non-principal ultrafilters can be found in [1]. The following fact about non-principal ultrafilters is useful.

LEMMA 8. *Let Ω be an infinite set, and let p be a non-principal ultrafilter. Then every element of p has infinitely many elements. In particular, the complement of every finite set is in p .*

PROOF. Let $A \in p$. Suppose, to the contrary, that A has finitely many elements. Split A into two nonempty subsets $A_1 \cup A_2 = A$. Then either A_1 or $A_1^C \in p$. If $A_1^C \in p$, then $A_2 = A_1^C \cap A \in p$. So either A_1 or A_2 is in p . Repeat this exercise with the set that is in p until one arrives at a set in p with exactly one element. This would make p a principal ultrafilter. To see that the complement of every finite set A is in p , suppose to the contrary that $A^C \notin p$. Then $A \in p$, which contradicts what we have already proved. \square

Let p and q be non-principal ultrafilters on the positive integers, and construct the ultrafilter product $q \cdot p$ on Ω as follows. For each $S \subseteq \Omega$, define $S_i = \{k : (i, k) \in S\}$. We say that $S \in q \cdot p$ if $\{i : S_i \in p\} \in q$. The following result follows easily from elementary properties of non-principal ultrafilters. (See [1, p. 157].)

LEMMA 9. *A necessary (but not sufficient) condition for $S \in q \cdot p$ is that there exist infinitely many i such that $(i, k) \in S$ for infinitely many k . A sufficient (but not necessary) condition for $S \in q \cdot p$ is that for all but*

finitely many i $(i, k) \in S$ for all but finitely many k . In particular, if the supremum over k of the cardinality of $\{i : (i, k) \in S\}$ is finite, then $S \notin q \cdot p$.

PROOF. For the necessary condition, assume that $S \in q \cdot p$, so that $T = \{i : S_i \in p\} \in q$. It follows from Lemma 8 that T has infinitely many elements. Also, for each $i \in T$, S_i has infinitely many elements according to Lemma 8 because $S_i \in p$. For the sufficient condition, assume that for all but finitely many i $(i, k) \in S$ for all but finitely many k . Let $T = \{i : S_i \text{ is the complement of a finite set}\}$. Lemma 8 says that $S_i \in p$ for each $i \in T$. Also, $T \in p$ because T is the complement of a finite set. This implies $S \in q \cdot p$. For the final claim, assume to the contrary that $S \in q \cdot p$. Let n be the supremum over k of the cardinality of $T_k = \{i : (i, k) \in S\}$. Because $S_1, \dots, S_{n+1} \in p$, $T = \bigcap_{i=1}^{n+1} S_i \in p$. Let $k \in T$. Then $k \in S_i$ for $i = 1, \dots, n+1$. But then the cardinality of T_k is at least $n+1$, a contradiction. \square

We use Lemma 9 to determine which of the following three sets is in the ultrafilter $q \cdot p$, as these sets are key to the main example of this section:

$$(15) \quad L = \{(i, k) : k > 2i\}, \quad U = \{(i, k) : k < 2i\}, \quad \text{and} \quad Q = \{(i, k) : k = 2i\}.$$

Then $L \in q \cdot p$ according to the sufficient condition in Lemma 9, while $U \notin q \cdot p$ according to the necessary condition. Also, $Q \notin q \cdot p$, according to the last result in Lemma 9.

We now make use of the ultrafilter $q \cdot p$ to extend β to the collection of all random variables of the form WS where S is the indicator function of an arbitrary subset $S \subseteq \Omega$. Set $\beta(WS) = 10$ if $S \in q \cdot p$ and $\beta(WS) = 0$ otherwise. The prevision P now extends easily to the linear span \mathcal{L} of all of the random variables for which β has been defined. The set \mathcal{L} consists of all random variables of the form

$$(16) \quad Y = X + \alpha WS,$$

where X is bounded, α is real, and S is an indicator of a subset of Ω . Define \mathcal{Y} to be the set of all $Y \in \mathcal{L}$ for which $\alpha \geq 0$ in (16). Because every nonempty event has positive probability, if $Y \in \mathcal{Y}$, then $P(Y|h)$ is finite for every $h \subseteq \Omega$. The following result follows easily from the linearity of β .

PROPOSITION 4. *If $Z \in \mathcal{Y}$ and $\{Z \leq cW\} \in q \cdot p$, then $\beta(Z) \leq 10c$. Hence, if for all $c > 0$ $\{Z > cW\} \notin q \cdot p$, then $\beta(Z) = 0$.*

Next, partition Ω by the values of W . That is, $\pi_W = \{h_k\}_{k=2}^{\infty}$ where $h_k = \{W = k\}$. Each h_k consists of exactly $k-1$ points, i.e., $h_k =$

$\{(k, 1), \dots, (k, k - 1)\}$ for $k = 2, 3, \dots$. The conditional distribution given h_k is uniform over those $k - 1$ points.

THEOREM 2. *With respect to the collection \mathcal{Y} , previsions are conglomerable in π_W but they are not disintegrable in π_W .*

PROOF. First we show that P is not disintegrable in π_W . Let L , U , and Q be as defined in (15). For each k , $U \cap h_k$ and $L \cap h_k$ have the same number of elements and all elements of h_k have the same probability. Hence,

$$P(U) = P(L) = \frac{1}{2}[1 - P(Q)] > 0.$$

Because $L \in q \cdot p$ and $U \notin q \cdot p$, $P(LW) = P(UW) + 10$. Because each $h \in \pi_W$ is a finite set, $P(Z|h) = E(Z|h)$ for each $h \in \pi_W$. Hence, for $k = 2, 3, \dots$,

$$P(UW|h_k) = P(LW|h_k) = \begin{cases} \frac{k}{2} & \text{if } k \text{ is odd,} \\ \frac{k(k-2)}{2(k-1)} & \text{if } k \text{ is even.} \end{cases}$$

Hence,

$$P[P(UW|H)] = P[P(LW|H)],$$

but $P(UW) \neq P(LW)$, and P is not disintegrable in π_W .

Next, we show that with respect to variables in \mathcal{Y} , P is conglomerable in π_W . Let $Z \in \mathcal{Y}$. Recall that $P(Z) = E(Z) + \beta(Z)$, and $\beta(Z) \geq 0$. We know that

$$\inf_{h \in \pi_W} P(Z|h) = \inf_k E(Z|h_k) \leq E(Z) \leq P(Z).$$

What remains is to show that $P(Z) \leq \sup_{h \in \pi_W} P(Z|h)$.

If $\sup_{h \in \pi_W} P(Z|h) = \infty$, the proof is complete. So, assume that

$$\sup_{h \in \pi_W} P(Z|h) = r < \infty.$$

Hence $\sup_k E(Z|h_k) = r$ and $E(Z) \leq r$. Since $Z \in \mathcal{Y}$, there exists $b > -\infty$ such that $b \leq Z(i, k)$ for all i, k . Let $Z' = Z - b$ so that $Z' \geq 0$ and $\beta(Z') = \beta(Z)$. The conditional previsions $\{P(Z'|h_k)\}_{k=2}^{\infty}$ are bounded above by $r - b$ and below by 0. Let $d > 0$. The Markov inequality says that

$$P(Z' > dk|h_k) = P(Z' > dk|h_k) \leq \frac{r - b}{dk}.$$

Recall that the conditional distribution $P(\cdot|h_k)$ is uniform over the $k - 1$ points in h_k . Hence, for each $d > 0$ and all k , at most $(r - b)/d$ out of

the $k - 1$ points in h_k may satisfy $Z'(i, k) > dW(i, k)$. That is, the event $\{Z' > dW\} \cap h_k$ contains of at most $(r - b)/d$ points for each k . By the last result in Lemma 9, we have $\{Z' > dW\} \notin q \cdot p$. Proposition 4 now says that $\beta(Z') = 0$, hence $\beta(Z) = 0$. So $P(Z) = E(Z) \leq r$, as required by conglomerability. \square

5. Discussion. Conglomerability and disintegrability are familiar concepts in the countably additive theory of probability, although the names may not be as familiar as the concepts. The law of total probability or “tower property” of conditional expectations is essentially disintegrability, namely that the mean of a conditional mean is the marginal mean. With disintegrability taken for granted, conglomerability is simply an instance of the property of countably additive expectations that the mean of a random variable lies in the closed convex hull of its range. Of course, the countably additive theory guarantees disintegrability by allowing the conditional probabilities of events to change with the partition on which one conditions. The well-known Borel paradox is a classic example of how this happens. In the countably additive theory [4] illustrates how pervasive the Borel paradox is. If one insists on $P(X|h)$ having a meaning for every random variable X and every nonempty event h , then not even the countably additive theory can guarantee disintegrability in every partition.

As a final note, it is important to keep in mind that the concepts of conglomerability and disintegrability are defined with respect to a collection of random variables. The larger the collection of random variables, the more conditions of the form (9) and (10) that each concept requires. That is, in order for P to be conglomerable in π with respect to a collection \mathcal{W} , (9) must hold for every $X \in \mathcal{W}$. Similarly, for P to be disintegrable in π with respect to \mathcal{W} , (10) must hold for every $X \in \mathcal{W}$. Consider the three collections $\mathcal{X} \subset \mathcal{Y} \subset \mathcal{Z}$ that figure in the results of this paper. That is, \mathcal{X} is the collection of all bounded random variables, \mathcal{Z} is the collection of all random variables that satisfy Assumption 2, and \mathcal{Y} is an intermediate collection such as the collection in Section 4. If P is conglomerable in π with respect to \mathcal{Z} , then \mathcal{Z} is of Class 2 relative to P and π and so are \mathcal{Y} and \mathcal{X} . Similarly, if P is disintegrable in \mathcal{Z} with respect to π , then all three collections are of Class 2. However, the equivalence of conglomerability and disintegrability does not carry over from larger collections to smaller collections. The reason is that \mathcal{Z} might be of Class 0 while \mathcal{Y} is of Class 1 and \mathcal{X} is of Class 2. Indeed, this is precisely what occurs in the example of Section 4.

References.

- [1] COMFORT, W. W. and NEGREPONTIS, S. (1974). *The Theory of Ultrafilters*, Springer-Verlag, Berlin.
- [2] DE FINETTI, B. (1974). *Theory of Probability (2 vols.)*, Wiley, New York.
- [3] DUBINS, L. (1975). Finitely additive conditional probabilities, conglomerability and disintegrations. *Annals of Probability* **3** 89–99.
- [4] KADANE, J., SCHERVISH, M. and SEIDENFELD, T. (1986). Statistical implications of finitely additive probability. In *Bayesian Inference and Decision Techniques* (P. K. Goel and A. Zellner, eds.) 59–76. North- Holland, Amsterdam.
- [5] KADANE, J. B., SCHERVISH, M. J. and SEIDENFELD, T. (1996). Reasoning to a foregone conclusion. *Journal of the American Statistical Association* **91** 1228–1235.
- [6] ROYDEN, H. L. (1968). *Real Analysis*, Macmillan, London.
- [7] SCHERVISH, M. J., KADANE, J. B. and SEIDENFELD, T. (2008). Fundamental theorems of prevision and asset pricing. *International Journal of Approximate Reasoning* to appear.
- [8] SEIDENFELD, T., SCHERVISH, M. J. and KADANE, J. B. (2006). When coherent preferences may not preserve indifference between equivalent random variables. Technical Report 840, Department of Statistics, Carnegie Mellon University.

MARK J. SCHERVISH AND JOSEPH B. KADANE
DEPARTMENT OF STATISTICS
CARNEGIE MELLON UNIVERSITY
PITTSBURGH, PA 15213
E-MAIL: mark@stat.cmu.edu
kadane@stat.cmu.edu

TEDDY SEIDENFELD
DEPARTMENT OF STATISTICS
AND DEPARTMENT OF PHILOSOPHY
CARNEGIE MELLON UNIVERSITY
PITTSBURGH, PA 15213
E-MAIL: teddy@stat.cmu.edu