

Separating Populations with Wide Data: a Spectral Analysis

Avrim Blum^{*}, Amin Coja-Oghlan^{**}, Alan Frieze^{***}, and Shuheng Zhou[†]

Carnegie Mellon University, Pittsburgh, PA 15213, USA

Abstract. In this paper, we consider the problem of partitioning a small data sample drawn from a mixture of k product distributions. We are interested in the case that individual features are of low average quality γ , and we want to use as few of them as possible to correctly partition the sample. We analyze a spectral technique that is able to approximately optimize the total data size—the product of number of data points n and the number of features K —needed to correctly perform this partitioning as a function of $1/\gamma$ for $K > n$. Our goal is motivated by an application in clustering individuals according to their population of origin using markers, when the divergence between any two of the populations is small.

1 Introduction

We explore a type of classification problem that arises in the context of computational biology. The problem is that we are given a small sample of size n , e.g., DNA of n individuals (think of n in the hundreds or thousands), each described by the values of K features or markers, e.g., SNPs (Single Nucleotide Polymorphisms, think of K as an order of magnitude larger than n). Our goal is to use these features to classify the individuals according to their population of origin. Features have slightly different probabilities depending on which population the individual belongs to, and are assumed to be independent of each other (i.e., our data is a small sample from a mixture of k very similar product distributions). The objective we consider is to minimize the total data size $D = nK$ needed to correctly classify the individuals in the sample as a function of the “average quality” γ of the features, under the assumption that $K > n$. Throughout the paper, we use p_i^j and μ_i^j as shorthands for $p_i^{(j)}$ and $\mu_i^{(j)}$ respectively.

Statistical Model: We have k probability spaces $\Omega_1, \dots, \Omega_k$ over the set $\{0, 1\}^K$. Further, the components (features) of $z \in \Omega_t$ are independent and $\Pr_{\Omega_t}[z_i = 1] = p_t^i$ ($1 \leq t \leq k, 1 \leq i \leq K$). Hence, the probability spaces $\Omega_1, \dots, \Omega_k$ comprise the distribution of the features for each of the k populations. Moreover, the input of the algorithm consists of a collection (mixture) of $n = \sum_{t=1}^k N_t$ unlabeled samples, N_t points from Ω_t , and the algorithm is to determine for each data point from which of $\Omega_1, \dots, \Omega_k$ it was chosen. In general we do *not* assume that N_1, \dots, N_t are revealed

^{*} Supported in part by the NSF under grant CCF-0514922.

^{**} Supported by the German Research Foundation under grant CO 646.

^{***} Supported in part by the NSF under grant CCF-0502793.

[†] Supported in part by the NSF under grants CCF-0625879 and CNF-0435382.

to the algorithm; but we do require some bounds on their relative sizes. An important parameter of the probability ensemble $\Omega_1, \dots, \Omega_k$ is the *measure of divergence*

$$\gamma = \min_{1 \leq s < t \leq k} \frac{\sum_{i=1}^K (p_s^i - p_t^i)^2}{K} \quad (1)$$

between any two distributions. Note that $\sqrt{K\gamma}$ measures the Euclidean distance between the means of any two distributions and thus represents their separation. Further, let $N = n/k$ (so if the populations were balanced we would have N of each type) and assume from now on that $kN < K$. Let $D = nK$ denote the size of the data-set. In addition, let $\sigma^2 = \max_{i,t} p_t^i(1 - p_t^i)$ denote the maximum variance of any random bit.

The biological context for this problem is we are given DNA information from n individuals from k populations of origin and we wish to classify each individual into the correct category. DNA contains a series of markers called SNPs, each of which has two variants (alleles). Given the population of origin of an individual, the genotypes can be reasonably assumed to be generated by drawing alleles independently from the appropriate distribution. The following theorem gives a sufficient condition for a balanced ($N_1 = N_2$) input instance when $k = 2$.

Theorem 1. (Zhou 06 [25]) *Assume $N_1 = N_2 = N$. If $K = \Omega(\frac{\ln N}{\gamma})$ and $KN = \Omega(\frac{\ln N \log \log N}{\gamma^2})$ then with probability $1 - 1/\text{poly}(N)$, among all balanced cuts in the complete graph formed among $2N$ sample individuals, the maximum weight cut corresponds to the partition of the $2N$ individuals according to their population of origin. Here the weight of a cut is the sum of weights across all edges in the cut, and the edge weight equals the Hamming distance between the bit vectors of the two endpoints.*

Variants of the above theorem, based on a model that allows two random draws from each SNP for an individual, are given in [3, 25]. In particular, notice that edge weights based on the inner-product of two individuals' bit vectors correspond to the sample covariance, in which case the max-cut corresponds to the correct partition [25] with high probability. Finding a max-cut is computationally intractable; hence in the same paper [3], a hill-climbing algorithm is given to find the correct partition for balanced input instances but with a stronger requirement on the sizes of both K and nK .

A Spectral Approach: In this paper, we construct two simpler algorithms using spectral techniques, attempting to reproduce conditions above. In particular, we study the requirements on the parameters of the model (namely, γ , N , k , and K) that allow us to classify every individual correctly and efficiently with high probability.

The two algorithms CLASSIFY and PARTITION compare as follows. Both algorithms are based on spectral methods originally developed in graph partitioning. More precisely, Theorem 2 is based on computing the singular vectors with the two largest singular values for each of the $n \times K$ input random matrix. The procedure is conceptually simple, easy to implement, and efficient in practice. For simplicity, Procedure Classify assumes the separation parameter γ is known to decide which singular vector to examine; in practice, one can just try both singular vectors as we do in the simulations. Proof techniques for Theorem 2, however, are difficult to apply to cases of multiple populations, i.e., $k > 2$. Procedure Partition is based on computing a rank- k

approximation of the input random matrix and can cope with a mixture of a constant number of populations. It is more intricate for both implementation and execution than CLASSIFY. It does not require γ as an input, while only requires that the constant k is given. We prove the following theorems.

Theorem 2. Let $\omega = \frac{\min(N_1, N_2)}{n}$ and ω_{\min} be a lower bound on ω . Let γ be given. Assume that $K > 2n \ln n$ and $k = 2$. Procedure CLASSIFY allows us to separate two populations w.h.p., when $n \geq \Omega\left(\frac{\sigma^2}{\gamma \omega_{\min} \omega}\right)$, where σ^2 is the largest variance of any random bit, i.e. $\sigma^2 = \max_{i,t} p_t^i(1 - p_t^i)$. Thus if the populations are roughly balanced, then $n \geq \frac{c}{\gamma}$ suffices for some constant c .

This implies that the data required is $D = nK = O(\ln n \sigma^4 / \gamma^2 \omega^2 \omega_{\min}^2)$. Let $P_s = (p_s^i)_{i=1, \dots, K}$, we have

$$\|P_1 - P_2\|_2 = \sqrt{K} \gamma = \sqrt{\sum_{i=1}^K (p_1^i - p_2^i)^2} \geq \frac{\sigma}{\sqrt{\omega_{\min} \omega}} \sqrt{\ln n}. \quad (2)$$

Theorem 3. Let $\omega = \frac{\min(N_1, \dots, N_k)}{n}$. There is a polynomial time algorithm PARTITION that satisfies the following. Suppose that $K > n \log n$ and $n > \frac{C_k \sigma^2}{\gamma \omega}$ for some large enough constant C_k and that $\omega = \Omega(1)$. Then given the empirical $n \times K$ matrix comprising the K features for each of the n individuals along with the parameter k , PARTITION separates the k populations correctly w.h.p.

Summary and Future Direction: Note that unlike Theorem 1, both Theorem 2 and Theorem 3 require a lower bound on n , even when $k = 2$ and the input instance is balanced. We illustrate through simulations to show that this seems not to be a fundamental constraint of the spectral techniques; our experimental results show that even when n is small, by increasing K so that $nK = \Omega(1/\gamma^2)$, one can classify a mixture of two populations using ideas in Procedure CLASSIFY with success rate reaching an ‘oracle’ curve, which is computed assuming that distributions are known, where success rate means the ratio between correctly classified individuals and N . Exploring the tradeoffs of n and K that are sufficient for classification, when sample size n is small, is both of theoretical interests and practical value.

1.1 Related Work

In their seminal paper [21], Pritchard, Stephens, and Donnelly presented a model-based clustering method to separate populations using genotype data. They assume that observations from each cluster are random from some parametric model. Inference for the parameters corresponding to each population is done jointly with inference for the cluster membership of each individual, and k in the mixture, using Bayesian methods.

The idea of exploiting the eigenvectors with the first two eigenvalues of the adjacency matrix to partition graphs goes back to the work of Fiedler [12], and has been used in the heuristics for various NP-hard graph partitioning problems (e.g., [13]). The main difference between graph partitioning problems and the classification problem

that we study is that the matrices occurring in graph partitioning are symmetric and hence diagonalizable, while our input matrix is rectangular in general. Thus, the contribution of Theorem 2 is to show that a conceptually simple and efficient algorithm based on singular value decompositions performs well in the framework of a fairly general probabilistic model, where probabilities for each of the K features for each of the k populations are allowed to vary. Indeed, the analysis of CLASSIFY requires exploring new ideas such as the Separation Lemma and the normalization of the random matrix X , for generating a large gap between top two singular values of the expectation matrix \mathcal{X} and for bounding the angle between random singular vectors and their static correspondents, details of which are included in Section 2 with analysis in full version.

Procedure Partition and its analysis build upon the spectral techniques of McSherry [18] on graph partitioning, and an extension due to Coja-Oghlan [4]. McSherry provides a comprehensive probabilistic model and presents a spectral algorithm for solving the partitioning problem on random graphs, provided that a separation condition similar to (2) is satisfied. Indeed, [18] encompasses a considerable portion of the prior work on Graph Coloring, Minimum Bisection, and finding Maximum Clique. Moreover, McSherry’s approach easily yields an algorithm that solves the classification problem studied in the present paper under similar assumptions as in Theorem 3, provided that the algorithm is given the parameter γ as an additional input; this is actually pointed out in the conclusions of [18]. In the context of graph partitioning, an algorithm that does not need the separation parameter as an input was devised in [4]. The main difference between PARTITION and the algorithm presented in [4] is that PARTITION deals with the asymmetric $n \times K$ matrix of individuals/features, whereas [4] deals with graph partitioning (i.e., a symmetric matrix).

There are two streams of related work in the learning community. The first stream is the recent progress in learning from the point of view of clustering: given samples drawn from a mixture of well-separated Gaussians (component distributions), one aims to classify each sample according to which component distribution it comes from, as studied in [8, 9, 2, 23, 1, 15, 7]. This framework has been extended to more general distributions such as log-concave distributions in [1, 15] and heavy-tailed distributions in [7], as well as to more than two populations. These results focus mainly on reducing the requirement on the separations between any two centers P_1 and P_2 . In contrast, we focus on the sample size D . This is motivated by previous results [3, 25] stating that by acquiring enough attributes along the same set of dimensions from each component distribution, with high probability, we can correctly classify every individual.

While our aim is different from those results, where $n > K$ is almost universal and we focus on cases $K > n$, we do have one common axis for comparison, the ℓ_2 -distance between any two centers of the distributions. In earlier works [9, 2], the separation requirement depended on the number of dimensions of each distribution; this has recently been reduced to be independent of K , the dimensionality of the distribution for certain classes of distributions [1, 15]. This is comparable to our requirement in (2) for the discrete distributions. For example, according to Theorem 7 in [1], in order to separate the mixture of two Gaussians,

$$\|P_1 - P_2\|_2 = \Omega \left(\frac{\sigma}{\sqrt{\omega}} + \sigma \sqrt{\log n} \right) \quad (3)$$

is required. Besides Gaussian and Logconcave, a general theorem: Theorem 6 in [1] is derived that in principle also applies to mixtures of discrete distributions. The key difficulty of applying their theorem directly to our scenario is that it relies on a concentration property of the distribution (Eq. (10) of [1]) that need not hold in our case. In addition, once the distance between any two centers is fixed (i.e., once γ is fixed in the discrete distribution), the sample size n in their algorithms is always larger than $\Omega(\frac{K}{\omega} \log^5 K)$ [1, 15] for log-concave distributions (in fact, in Theorem 3 of [15], they discard at least this many individuals in order to correctly classify the rest in the sample), and larger than $\Omega(\frac{K}{\omega})$ for Gaussians [1], whereas in our case, $n < K$ always holds. Hence, our analysis allows one to obtain a clean bound on n in the discrete case.

The second stream of work is under the PAC-learning framework, where given a sample generated from some target distribution Z , the goal is to output a distribution Z_1 that is close to Z in Kullback-Leibler divergence: $KL(Z||Z_1)$, where Z is a mixture of product distributions over discrete domains or Gaussians [16, 14, 5, 6, 20, 10, 11]. They do not require a minimal distance between any two distributions, but they do not aim to classify every sample point correctly either, and in general require much more data.

2 A Simple Algorithm Using Singular Vectors

As described in Theorem 2, we assume we have a mixture of two product distributions. Let N_1, N_2 be the number of individuals from each population class. Our goal is to correctly classify all individuals according to their distributions. Let $n = 2N = N_1 + N_2$, and refer to the case when $N_1 = N_2$ as the balanced input case. For convenience, let us redefine “ K ” to assume we have $O(\log n)$ blocks of K features each (so the total number of features is really $O(K \log n)$) and we assume that each set of K features has divergence at least γ . (If we perform this partitioning of features into blocks randomly, then with high probability this divergence has changed by only a constant factor for most blocks.) The high-level idea of the algorithm is now to repeat the following procedure for each block of K features: use the K features to create an $n \times K$ matrix X , such that each row $X_i, i = 1, \dots, n$, corresponds to a feature vector for one sample point, across its K dimensions. We then compute the top two left singular vectors u_1, u_2 of X and use these to classify each sample. This classification induces some probability of error f for each individual at each round, so we repeat the procedure for each of the $O(\log n)$ blocks and then take majority vote over different runs. Each round we require $K \geq n$ features, so we need $O(n \log n)$ features total in the end.

In more detail, we repeat the following procedure $O(\log n)$ times. Let $T = \frac{15N}{32} \sqrt{3\omega_{\min}\gamma}$, where ω_{\min} is the lower bound on the minimum weight $\min\{\frac{N_1}{2N}, \frac{N_2}{2N}\}$, which is independent of an actual instance. Let $s_1(X), s_2(X)$ be the top two singular values of X .

Procedure Classify: Given γ, N, ω_{\min} . Assume that $N \gg \frac{1}{\gamma}$,

- Normalization: use the K features to form a random $n \times K$ matrix X ; Each individual random variable $X_{i,j}$ is a *normalized* random variable based on the original Bernoulli r.v. $b_{i,j} \in \{0, 1\}$ with $\Pr[b_{i,j} = 1] = p_1^j$ for $X_i \in P_1$ and $\Pr[b_{i,j} = 1] = p_2^j$ for $X_i \in P_2$, such that $X_{i,j} = \frac{b_{i,j}}{2}$.

- Take top two left singular vectors u_1, u_2 of X , where $u_i = [u_{i,1}, \dots, u_{i,n}]$, $i = 1, 2$.
 1. If $s_2(X) > T = \frac{15N}{32} \sqrt{3\omega_{\min}\gamma}$, use u_2 to partition the individuals with 0 as the threshold, i.e., partition $j \in [n]$ according to $u_{2,j} < 0$ or $u_{2,j} \geq 0$.
 2. Otherwise, use u_1 to partition, with mixture mean $M = \sum_{i=1}^n u_{1,n}$ as the threshold.

Analysis of the Simple Algorithm: Our analysis is based on comparing entries in the top two singular vectors of the normalized random $n \times K$ matrix X , with those of a static matrix \mathcal{X} , where each entry $\mathcal{X}_{i,j} = \mathbf{E}[X_{i,j}]$ is the expected value of the corresponding entry in X . Hence $\forall i = 1, \dots, N_1$, $\mathcal{X}_i = [\mu_1^1, \mu_1^2, \dots, \mu_1^K]$, where $\mu_1^j = \frac{1+p_1^j}{2}$, $\forall j$, and $\forall i = N_1 + 1, \dots, n$, $\mathcal{X}_i = [\mu_2^1, \mu_2^2, \dots, \mu_2^K]$, where $\mu_2^j = \frac{1+p_2^j}{2}$, $\forall j$. We assume the divergence is exactly γ among the K features that we have chosen in all calculations.

The inspiration for this approach is based on the following lemma, whose proof is built upon a theorem that is presented in a lecture note by Spielman [22]. For a $n \times K$ matrix A , let $s_1(A) \geq s_2(A) \geq \dots \geq s_n(A)$ be singular values of A . Let $u_1, \dots, u_n, v_1, \dots, v_n$, be the n left and right singular vectors of X , corresponding to $s_1(X), \dots, s_n(X)$ such that $\|u_i\|_2 = 1, \|v_i\|_2 = 1, \forall i$. We denote the set of n left and right singular vectors of \mathcal{X} with $\bar{u}_1, \dots, \bar{u}_n, \bar{v}_1, \dots, \bar{v}_n$.

Lemma 4. *Let X be the random $n \times K$ matrix and \mathcal{X} its expected value matrix. Let $A = X - \mathcal{X}$ be the zero-mean random matrix. Let θ_i be the angle between two vectors: $[u_i, v_i], [\bar{u}_i, \bar{v}_i]$, where $\|[u_i, v_i]\|_2 = \|[\bar{u}_i, \bar{v}_i]\|_2 = 2$ and $[u, v]$ represents a vector that is the concatenation of two vectors u, v .*

$$\|u_i - \bar{u}_i\|_2 \leq \|[u_i, v_i] - [\bar{u}_i, \bar{v}_i]\|_2 \approx 2\theta_i \approx 2 \sin(\theta_i) \leq \frac{4s_1(A)}{\text{gap}(i, \mathcal{X})}, \quad (4)$$

where $\text{gap}(i, \mathcal{X}) = \min_{j \neq i} |s_i(\mathcal{X}) - s_j(\mathcal{X})|$.

We first bound the largest singular value $s_1(A) = s_1(X - \mathcal{X})$ of $(a_{i,j})$ with independent zero-mean entries, which defines the Euclidean operator norm

$$\|(a_{i,j})\| := \sup \left\{ \sum_{i,j} a_{i,j} x_i y_j : \sum x_i^2 \leq 1, \sum y_i^2 \leq 1 \right\}. \quad (5)$$

The behavior of the largest singular value of an $n \times m$ random matrices A with i.i.d. entries is well studied. Latala [17] shows that the weakest assumption for its regular behavior is boundedness of the fourth moment of the entries, even if they are not identically distributed. Combining Theorem 5 of Latala with the concentration Theorem 6 by Meckes [19] proves Theorem 7 that we need¹.

Theorem 5. (Bounded Norm of Random Matrices [17]) *For any finite $n \times m$ matrix A of independent mean zero r.v.'s $a_{i,j}$ we have, for an absolute constant C ,*

$$\mathbf{E} \|(a_{i,j})\| \leq C \left(\max_i \sqrt{\sum_j \mathbf{E} a_{i,j}^2} + \max_j \sqrt{\sum_i \mathbf{E} a_{i,j}^2} + \left(\sum_{i,j} \mathbf{E} a_{i,j}^4 \right)^{\frac{1}{4}} \right). \quad (6)$$

¹ One can also obtain an upper bound of $O(\sqrt{n+K})$ on $s_1(A)$ using a theorem on by Vu [24], through the construction a $(n+K) \times (n+K)$ square matrix out of A .

Theorem 6. (Concentration of Largest Singular Value: Bounded Range [19]) For any finite $n \times m$, where $n \leq m$, matrix A , such that entries $a_{i,j}$ are independent r.v. supported in an interval of length at most D , then, for all t ,

$$\Pr[|s_1(A) - \mathbb{M}s_1(A)| \geq t] \leq 4e^{-t^2/4D^2}. \quad (7)$$

Theorem 7. (Largest Singular Value of a Mean-zero Random Matrix) For any finite $n \times K$, where $n \leq K$, matrix A , such that entries $a_{i,j}$ are independent mean zero r.v. supported in an interval of length at most D , with fourth moment upper bounded by B , then

$$\Pr[s_1(A) \geq CB^{1/4}\sqrt{K} + 4D\sqrt{\pi} + t] \leq 4e^{-t^2/4} \quad (8)$$

for all t . Hence $\|A\| \leq C_1B^{1/4}\sqrt{K}$ for an absolute constant C_1 .

2.1 Generating a Large Gap in $s_1(\mathcal{X})$, $s_2(\mathcal{X})$

In order to apply Lemma 4 to the top two singular vectors of X and \mathcal{X} through

$$\|u_1 - \bar{u}_1\|_2 \leq \frac{4s_1(X - \mathcal{X})}{|s_1(\mathcal{X}) - s_2(\mathcal{X})|} \quad (9)$$

$$\|u_2 - \bar{u}_2\|_2 \leq \frac{4s_1(X - \mathcal{X})}{\min(|s_1(\mathcal{X}) - s_2(\mathcal{X})|, |s_2(\mathcal{X})|)}, \quad (10)$$

we need to first bound $|s_1(\mathcal{X}) - s_2(\mathcal{X})|$ away from zero, since otherwise, RHSs on both (9) and (10) become unbounded. We then analyze $\text{gap}(2, \mathcal{X}) = \min(|s_1(\mathcal{X}) - s_2(\mathcal{X})|, |s_2(\mathcal{X})|)$.

Let us first define values a, b, c that we use throughout the rest of the paper:

$$a = \sum_{k=1}^K (\mu_1^k)^2, \quad b = \sum_{k=1}^K \mu_1^k \mu_2^k, \quad c = \sum_{k=1}^K (\mu_2^k)^2. \quad (11)$$

For the following analysis, we can assume that $a, b, c \in [K/4, K]$, given that X is normalized in Procedure Classify.

We first show that normalization of X as described in Procedure Classify guarantees that not only $|s_1(\mathcal{X}) - s_2(\mathcal{X})| \neq 0$, but there also exists a $\Theta(\sqrt{NK})$ amount of gap between $s_1(\mathcal{X})$ and $s_2(\mathcal{X})$ in Proposition 8:

$$\text{gap}(\mathcal{X}) := |s_1(\mathcal{X}) - s_2(\mathcal{X})| = \Theta(\sqrt{NK}). \quad (12)$$

Proposition 8. For a normalized random matrix X , its expected value matrix \mathcal{X} satisfies $\frac{4c_0\sqrt{2NK}}{5} \leq \text{gap}(\mathcal{X}) \leq \sqrt{2NK}$, where $c_0 = \frac{|b|\sqrt{ac}}{K(a+c)}$ is a constant, given that $a, b, c \in [K/4, K]$ as defined in (11). In addition,

$$\sqrt{\frac{KN}{4}} \leq s_1(\mathcal{X}) \leq \sqrt{2NK}, \text{ and } \sqrt{\frac{NK}{2}} \leq s_1(\mathcal{X}) + s_2(\mathcal{X}) \leq \sqrt{2NK}. \quad (13)$$

We next state a few important results that justify Procedure Classify. Note that the left singular vectors $\bar{u}_i, \forall i$ of \mathcal{X} are of the form $[x_i, \dots, x_i, y_i, \dots, y_i]^T$:

$$\bar{u}_1 = [x_1, \dots, x_1, y_1, \dots, y_1]^T, \text{ and } \bar{u}_2 = [x_2, \dots, x_2, y_2, \dots, y_2]^T, \quad (14)$$

where x_i repeats N_1 times and y_i repeats N_2 times. We first show Proposition 9 regarding signs of $x_i, y_i, i = 1, 2$, followed by a lemma bounding the separation of x_2, y_2 . We then state the key Separation Lemma that allows us to conclude that least one of top two left singular vectors of X can be used to classify data at each round. It can be extended to cases when $k > 2$.

Proposition 9. *Let b as defined in (11): when $b > 0$, entries x_1, y_1 in \bar{u}_1 have the same sign while x_2, y_2 in \bar{u}_2 have opposite signs.*

Lemma 10. $|x_2 - y_2|^2 \leq \frac{C_{\max}}{2N}$ where $C_{\max} = \left(\sqrt{\frac{1}{\omega_1}} + \sqrt{\frac{1}{\omega_2}}\right)^2 \leq \frac{4}{\omega_{\min}}$; $|x_2|^2 \geq \frac{C_{x \min}}{2N}$ where $C_{x \min} = \frac{\omega_2}{4\omega_1^2 + \omega_1\omega_2}$; $|y_2|^2 \geq \frac{C_{y \min}}{2N}$ where $C_{y \min} = \frac{\omega_1}{4\omega_2^2 + \omega_1\omega_2}$.

Lemma 11. (Separation Lemma) $K\gamma = s_1(\mathcal{X})^2(x_1 - y_1)^2 + s_2(\mathcal{X})^2(x_2 - y_2)^2$.

Proof. Let $\Delta := P_1 - P_2$ as in Theorem 2, and $\mathbf{b} = [1, 0, \dots, 0, -1, 0, \dots, 0]^T$, where 1 appears in the first and -1 appears in the $N_1 + 1^{\text{st}}$ positions. Then $\Delta = X^T \mathbf{b} = [\mu_1^1 - \mu_2^1, \mu_1^2 - \mu_2^2, \dots, \mu_1^K - \mu_2^K]$. Given $\mathcal{X} = s_1(\mathcal{X})\bar{u}_1\bar{v}_1^T + s_2(\mathcal{X})\bar{u}_2\bar{v}_2^T$, we thus rewrite Δ as: $\Delta = \mathcal{X}^T \mathbf{b} = s_1(\mathcal{X})\bar{v}_1\bar{u}_1^T \mathbf{b} + s_2(\mathcal{X})\bar{v}_2\bar{u}_2^T \mathbf{b} = s_1(\mathcal{X})\bar{v}_1(x_1 - y_1) + s_2(\mathcal{X})\bar{v}_2(x_2 - y_2)$. The lemma follows from the fact that $\|\Delta\|_2 = \sqrt{K}\gamma$ and \bar{v}_1, \bar{v}_2 are orthonormal. ■

Combining Proposition 9, Lemma 10, (13), and Lemma 11, we have

Corollary 12. $s_2(\mathcal{X}) \leq \frac{\sqrt{2N}K\gamma}{\sqrt{c_{x \min}} + \sqrt{c_{y \min}}}$, and hence $\text{gap}(2, \mathcal{X}) = \min(s_2(\mathcal{X}), |s_1(\mathcal{X}) - s_2(\mathcal{X})|) = s_2(\mathcal{X})$ for a sufficiently small γ .

Finally, we show that the probability of error at each round for each individual is at most $f = 1/10$, given the sample size n as specified in Theorem 2. Hence by taking majority vote over the different runs for each sample, our algorithm will find the correct partition with probability $1 - 1/n^2$, given that at each round we take a set of $K > n$ independent features. We leave the detailed analysis in full version.

3 The Algorithm PARTITION

As in Section 2, by repeating the partitioning process $\log n$ times, we may restrict our attention to the problem of classifying a constant fraction of the individuals correctly. Let $V = \{1, \dots, n\}$ be the set of all n individuals, and let $\psi : V \rightarrow \{1, \dots, k\}$ be the map that assigns to each individual the population it belongs to. Further, set $V_t = \psi^{-1}(t)$, define $N_t = |V_t|$, $\Gamma = K\gamma$, and $\lambda = \sqrt{K}\sigma$. In addition, let $A = (a_{vi})$ denote the empirical $n \times K$ input matrix. Then the assumption from Theorem 3 can be rephrased as $n_{\min}K\gamma > C_k\lambda^2$.

If $X = (x_{ij})_{1 \leq i \leq n, 1 \leq j \leq K}$ is a $n \times K$ matrix, then we let $\|X\| = \max_{\|\xi\|=1} \|X\xi\|$ signify the operator norm of X , while $\|X\|_F = \left(\sum_{i,j} x_{ij}^2\right)^{\frac{1}{2}}$ denotes the Frobenius

norm. The algorithm PARTITION computes a *rank k approximation* \widehat{A} of the input matrix A . That is, \widehat{A} is a $n \times K$ matrix of rank at most k , and if B is any $n \times K$ matrix of rank at most k , then $\|A - \widehat{A}\| \leq \|A - B\|$. Such an \widehat{A} can be computed in polynomial time via singular value decomposition. Let \widehat{A}_v denote the v -row of \widehat{A} .

Algorithm 13. PARTITION(A, k)

Input: A $n \times K$ matrix A and the parameter k . *Output:* A partition S_1, \dots, S_k of V .

1. Compute a rank k approximation \widehat{A} of A .
- For $j = 1, \dots, 2 \log K$ do
 2. Let $\Gamma_j = K2^{-j}$ and compute $Q^{(j)}(v) = \{w \in V : \|\widehat{A}_w - \widehat{A}_v\|^2 \leq 0.01\Gamma_j^2\}$ for all $v \in V$.
Then, determine sets $Q_1^{(j)}, \dots, Q_k^{(j)}$ as follows: for $i = 1, \dots, k$ do
 3. Pick $v \in V \setminus \bigcup_{l=1}^{i-1} Q_l^{(j)}$ such that $|Q^{(j)}(v) \setminus \bigcup_{l=1}^{i-1} Q_l^{(j)}|$ is maximum.
Set $Q_i^{(j)} = Q^{(j)}(v) \setminus \bigcup_{l=1}^{i-1} Q_l^{(j)}$ and $\xi_i^{(j)} = \frac{1}{|Q_i^{(j)}|} \sum_{w \in Q_i^{(j)}} \widehat{A}_w$.
 4. Partition the entire set V as follows: first, let $S_i^{(j)} = Q_i^{(j)}$ for all $1 \leq i \leq k$.
Then, add each $v \in V \setminus \bigcup_{l=1}^k Q_l^{(j)}$ to a set $S_i^{(j)}$ such that $\|\widehat{A}_v - \xi_i^{(j)}\|$ is minimum.
Set $r_j = \sum_{i=1}^k \sum_{v \in S_i^{(j)}} \|\widehat{A}_v - \xi_i^{(j)}\|^2$.
5. Let J be such that $r^* = r_J$ is minimum. Return $S_1^{(J)}, \dots, S_k^{(J)}$.

The basic idea behind PARTITION is to classify each individual $v \in V$ according to its row vector \widehat{A}_v in the rank k approximation \widehat{A} . That is, two individuals v, w are deemed to belong to the same population iff $\|\widehat{A}_v - \widehat{A}_w\|^2 \leq 0.01\Gamma^2$. Hence, PARTITION tries to determine sets S_1, \dots, S_k such that for any two v, w in the same set S_j the distance $\|\widehat{A}_v - \widehat{A}_w\|$ is small. To see why classifying the individuals according to their corresponding row vectors in \widehat{A} is a good idea, we consider an auxiliary matrix $\mathbb{E} = (\mathbb{E}_{vi})$ with entries $\mathbb{E}_{vi} = p_{\psi(v)}^i$. Thus, the entries of \mathbb{E} equal the expectations of the entries of A .

Lemma 14. *There is a constant $C > 0$ such that $\sum_{v \in V} \|\widehat{A}_v - \mathbb{E}_v\|^2 \leq Ck\lambda^2$ whp.*

Proof. Recall that \widehat{A} and \mathbb{E} both have rank $\leq k$, we obtain

$$\sum_{v \in V} \|\widehat{A}_v - \mathbb{E}_v\|^2 = \|\widehat{A} - \mathbb{E}\|_F^2 \leq 2k\|\widehat{A} - \mathbb{E}\| \leq 8k\|A - \mathbb{E}\|^2 \leq Ck\lambda^2,$$

where the last inequality follows from Theorem 7. ■

Observe that Lemma 14 implies that for *most* v we have $\|\widehat{A}_v - \mathbb{E}_v\|^2 \leq 10^{-6}\Gamma$, say. For letting $z = |\{v : \|\widehat{A}_v - \mathbb{E}_v\|^2 > 10^{-6}\Gamma\}|$, we get $10^{-6}\Gamma z \leq \sum_{v \in V} \|\widehat{A}_v - \mathbb{E}_v\|^2 \leq Ck\lambda^2$, whence $z \ll n_{\min}$ due to our assumption that $n_{\min}\Gamma \gg k\lambda^2$. Thus, most rows of \widehat{A} are close to the corresponding rows of the *expected* matrix \mathbb{E} . Since Γ is not given to the algorithm as an input parameter, PARTITION has to estimate Γ on its own.

To this end, the outer loop goes through $2 \log K$ “candidate values” Γ_j . These values are then used to obtain a partition $Q_1^{(1)}, \dots, Q_1^{(k)}$ in Steps 2–4, which are similar to the algorithm presented in [18]. In addition, Step 4 computes the error parameter r_j . Finally Step 5 outputs the partition that minimizes the error parameter r_j . More precisely, Step 2

uses Γ_j to compute for each $v \in V$ the set $Q(v)$ of elements w such that $\|\widehat{A}_w - \widehat{A}_v\| \leq 0.01\Gamma_j^2$. Then, Step 3 tries to compute “big” disjoint $Q_1^{(j)}, \dots, Q_k^{(j)}$, where each $Q_i^{(j)}$ results from some $Q(v_i)$. Further, Step 4 assigns all elements v not covered by $Q_1^{(j)}, \dots, Q_k^{(j)}$ to that $Q_i^{(j)}$ whose “center vector” $\xi_i^{(j)}$ is closest to \widehat{A}_v .

Thus, we need to show that eventually picking the partition whose error term r_j is minimum yields a good approximation to the ideal partition V_1, \dots, V_k . The basic reason why this is true is that $\xi_i^{(j)}$ should approximate the expectation \mathbb{E}^{V_i} for class V_i well iff $Q_i^{(j)}$ is a good approximation of V_i . Hence, if $Q_1^{(j)}, \dots, Q_k^{(j)}$ is “close” to V_1, \dots, V_k , then $r_j = \sum_{i=1}^k \sum_{v \in S_i^{(j)}} \|\widehat{A}_v - \xi_i^{(j)}\|^2 \approx \|\widehat{A} - \mathbb{E}\|_F^2$ will be about as small as $\|\widehat{A} - \mathbb{E}\|_F^2$ (cf. Lemma 14). Furthermore, Lemma 16 shows that any partition such that r_j is small yields a good approximation of V_1, \dots, V_k . Theorem 3 is an immediate consequence of Lemmas 15 and 16.

Lemma 15. *If $\frac{1}{2}\Gamma \leq \Gamma_j \leq \Gamma$, then $r_j \leq C_0 k^3 \lambda^2$ for a certain constant $C_0 > 0$.*

Lemma 16. *Let S_1, \dots, S_k be a partition and ξ_1, \dots, ξ_k a sequence of vectors such that $\sum_{i=1}^k \sum_{v \in S_i} \|\xi_i - A_v^*\|^2 \leq C_0 k^3 \lambda^2$. Then there is a bijection $\Xi : \{1, \dots, k\} \rightarrow \{1, \dots, k\}$ such that the following holds.*

1. $\|\xi_i - \mathbb{E}^{V_{\Xi(i)}}\|^2 \leq 0.001\Gamma^2$ for all $i = 1, \dots, k$, and
2. $\sum_{i=1}^k |S_i \Delta V_{\Xi(i)}| < 0.001n_{\min}$.

4 Experiments

We illustrate the effectiveness of spectral techniques using simulations. In particular, we explore the case when we have a mixture of two populations; we show that when $NK > 1/\gamma^2$ and $K > 1/\gamma$, either the first or the second left singular vector of X shows an approximately correct partitioning, meaning that the success rate is well above 1/2. The entry-wise expected value matrix \mathcal{X} is: among $K/2$ features, $p_1^i > p_2^i$ and for the other half, $p_1^i < p_2^i$ such that $\forall i, p_1^i, p_2^i \in \{\frac{1+\alpha}{2} + \frac{\epsilon}{2}, \frac{1-\alpha}{2} + \frac{\epsilon}{2}\}$, where $\epsilon = 0.1\alpha$. Hence $\gamma = \alpha^2$. We report results on balanced cases only, but we do observe that unbalanced cases show similar tradeoffs. For each population P , the success rate is defined as the number of individuals that are correctly classified, i.e., they belong to a group that P is the majority of that group, versus the size of the population $|P|$.

Each point on the SVD curve corresponds to an average rate over 100 trials. Since we are interested in exploring the tradeoffs of N, K in all ranges (e.g., when $N \ll K$ or $N \gg K$), rather than using the threshold T in Procedure Classify that is chosen in case both $N, K > 1/\gamma$, to decide which singular vector to use, we try both u_1 and u_2 and use the more effective one to measure the success rate at each trial. For each data point, the distribution of X is fixed across all trials and we generate an independent $X_{2N \times K}$ for each trial to measure success rate based on the more effective classifier between u_1 and u_2 .

One can see from the plot that when $K < 1/\gamma$, i.e., when $K = 200$ and 400 , no matter how much we increase N , the success rate is consistently low. Note that 50/100 of success rate is equivalent to a total failure. In contrast, when N is smaller than $1/\gamma$,

as we increase K , we can always classify with a high success rate, where in general, $NK > 1/\gamma^2$ is indeed necessary to see a high success rate. In particular, the curves for $K = 5000, 2500, 1250$ show the sharpness of the threshold behavior for increasing sample size n from below $1/K\gamma^2$ to above. For each curve, we also compute the best possible classification one could hope to make if one knew in advance which features satisfied $p_1^i > p_2^i$ and which satisfied $p_1^i < p_2^i$. These are the horizontal(ish) dotted lines above each curve. The fact that the solid curves are approaching these information-theoretic upper bounds shows that the spectral technique is correctly using the available information.




Fig. 1. Plots show success rate as a function of N for several values of K , when $\gamma = (0.04)^2$. Each point is an average over 100 trials. Horizontal lines (“oracles”) indicate the information-theoretically best possible success rate for that value of K (how well one could do if one knew in advance which features satisfied $p_1^i > p_2^i$ and which satisfied $p_1^i < p_2^i$; they are not exactly horizontal because they are also an average over 100 runs). Vertical bars indicate the value of N for which $NK = 1/\gamma^2$.

5 Acknowledgments

We thank John Lafferty, Frank McSherry, Roman Vershynin and Larry Wasserman for many helpful discussions on this work.

References

1. D. Achlioptas and F. McSherry. On spectral learning of mixtures of distributions. In *Proceedings of the 18th Annual COLT*, pages 458–469, 2005. (Version in <http://www.cs.ucsc.edu/> op-

tas/papers/).

2. S. Arora and R. Kannan. Learning mixtures of arbitrary gaussians. In *Proceedings of 33rd ACM Symposium on Theory of Computing*, pages 247–257, 2001.
3. K. Chaudhuri, E. Halperin, S. Rao, and S. Zhou. A rigorous analysis of population stratification with limited data. In *Proceedings of the 18th ACM-SIAM SODA*, 2007.
4. A. Coja-Oghlan. An adaptive spectral heuristic for partitioning random graphs. In *Proceedings of the 33rd ICALP*, 2006.
5. M. Cryan. *Learning and approximation Algorithms for Problems motivated by evolutionary trees*. PhD thesis, University of Warwick, 1999.
6. M. Cryan, L. Goldberg, and P. Goldberg. Evolutionary trees can be learned in polynomial time in the two state general markov model. *SIAM J. of Computing*, 31(2):375–397, 2002.
7. A. Dasgupta, J. Hopcroft, J. Kleinberg, and M. Sandler. On learning mixtures of heavy-tailed distributions. In *Proceedings of the 46th IEEE FOCS*, pages 491–500, 2005.
8. S. Dasgupta. Learning mixtures of gaussians. In *Proceedings of the 40th IEEE Symposium on Foundations of Computer Science*, pages 634–644, 1999.
9. S. Dasgupta and L. J. Schulman. A two-round variant of em for gaussian mixtures. In *Proceedings of the 16th Conference on Uncertainty in Artificial Intelligence (UAI)*, 2000.
10. J. Feldman, R. O’Donnell, and R. Servedio. Learning mixtures of product distributions over discrete domains. In *Proceedings of the 46th IEEE FOCS*, 2005.
11. J. Feldman, R. O’Donnell, and R. Servedio. PAC learning mixtures of Gaussians with no separation assumption. In *Proceedings of the 19th Annual COLT*, 2006.
12. M. Fiedler. Algebraic connectivity of graphs. *Czechoslovak Mathematical Journal*, pages 298–305, 1973.
13. P. Fjallstrom. Algorithms for graph partitioning: a survey. Technical report, Linkoping University Electroni Press, 1998.
14. Y. Freund and Y. Mansour. Estimating a mixture of two product distributions. In *Proceedings of the 12th Annual COLT*, pages 183–192, 1999.
15. R. Kannan, H. Salmasian, and S. Vempala. The spectral method for general mixture models. In *Proc. of the 18th Annual COLT*, 2005.
16. M. Kearns, Y. Mansour, D. Ron, R. Rubinfeld, R. Schapir, and L. Sellie. On the learnability of discrete distributions. In *Proceedings of the 26th ACM STOC*, pages 273–282, 1994.
17. R. Latala. Some estimates of norms of random matrices. In *Proceedings of the American Mathematical Society*, volume 133, pages 1273–1282, 2005.
18. Frank McSherry. Spectral partitioning of random graphs. In *Proceedings of the 42nd IEEE Symposium on Foundations of Computer Science*, pages 529–537, 2001.
19. M. Meckes. Concentration of norms and eigenvalues of random matrices. *J. Funct. Anal.*, 211(2):508–524, 2004.
20. E. Mossel and S. Roch. Learning nonsingular phylogenies and hidden markov models. In *Proceedings of the 37th ACM STOC*, 2005.
21. J. K. Pritchard, M. Stephens, and P. Donnelly. Inference of population structure using multilocus genotype data. *Genetics*, 155:954–959, June 2000.
22. D. Spielman. The behavior of algorithms in practice, 2002. Lecture notes.
23. V. Vempala and G. Wang. A spectral algorithm of learning mixtures of distributions. In *Proceedings of the 43rd IEEE FOCS*, pages 113–123, 2002.
24. V. Vu. Spectral norm of random matrices. In *Proceedings of 37th ACM STOC*, pages 423–430, 2005.
25. S. Zhou. *Routing, Disjoint Paths, and Classification*. PhD thesis, Carnegie Mellon University, Pittsburgh, PA, 2006. CMU Technical Report, CMU-PDL-06-109.

A Detailed Analysis for the Simple Algorithm

In this section, we first prove a proposition regarding a, b, c as defined in (11). We next provide the proof to Theorem 7 regarding the largest singular value of $(X - \mathcal{X})$. We then use Lemma 4 to bound the number of individuals that we misclassify at each round in Section A.2. We finish by showing that with high probability, we can correctly classify all individuals by taking majority vote over $O(\log N)$ different runs.

Throughout the rest of the paper, we use X, Y, H to represent random matrices, where $H = XX^T$ and $Y = \begin{bmatrix} 0 & X \\ X^T & 0 \end{bmatrix}$. We use $\mathcal{X}, \mathcal{Y}, \mathcal{H}$ to represent the corresponding static matrices. Let us substitute a, b, c in $\mathcal{H} = \mathcal{X}\mathcal{X}^T$, where the blocks in \mathcal{H} from top to bottom and from left to right are of size: $N_1 \times N_1, N_1 \times N_2, N_2 \times N_1$ and $N_2 \times N_2$ respectively:

$$\mathcal{H} = \mathcal{X}\mathcal{X}^T = \begin{bmatrix} a & \dots & a & b & \dots & b \\ \dots & & & & & \\ a & \dots & a & b & \dots & b \\ b & \dots & b & c & \dots & c \\ \dots & & & & & \\ b & \dots & b & c & \dots & c \end{bmatrix}_{2N \times 2N}. \quad (15)$$

Proposition 17. *For any choices of μ_i^k , $ac \geq b^2$; By definition,*

$$a + c - 2b = \sum_{i=1}^K \alpha_k^2, \text{ where } \alpha_k = |\mu_1^k - \mu_2^k|. \quad (16)$$

Proof. $a + c - 2b = \sum_k \alpha_k^2$ holds by definition.

$$\begin{aligned} ac - b^2 &= \sum_{k=1}^K (\mu_1^k)^2 \sum_{k=1}^K (\mu_2^k)^2 - \left(\sum_{k=1}^K (\mu_1^k \mu_2^k) \right)^2 \\ &= \sum_{k=1}^K (\mu_1^k \mu_1^k)^2 + \sum_{j \neq k} ((\mu_1^k \mu_2^j)^2 + (\mu_1^j \mu_2^k)^2) - \left(\sum_{k=1}^K (\mu_1^k \mu_1^k)^2 + \sum_{j \neq k} 2\mu_1^k \mu_2^k \mu_1^j \mu_2^j \right) \\ &= \sum_{j \neq k} (\mu_1^k \mu_2^j)^2 + (\mu_1^j \mu_2^k)^2 - 2\mu_1^k \mu_2^k \mu_1^j \mu_2^j = \sum_{j \neq k} (\mu_1^k \mu_2^j - \mu_1^j \mu_2^k)^2 \geq 0. \end{aligned}$$

■

Remark 18. Both matrices of \mathcal{X} and $\mathcal{X}\mathcal{X}^T$ have rank at most two. When $ac = b^2$, \mathcal{H} has rank 1.

A.1 Proof of Theorem 7

Proof of Theorem 7. By having an upper bound on both maximum variance and fourth moment of any entry, we have the following corollary of Theorem 5.

Corollary 19. (Largest Singular Value: Bounded Fourth Moment [17]) For any finite $n \times m$, where $n \leq m$, matrix of independent mean zero r.v.'s $a_{i,j}$, such that the maximum variances of any entry is at most σ^2 , and each entry has a finite fourth moment B we have

$$\mathbf{E} \|(a_{i,j})\| \leq C \left(\sigma(\sqrt{m} + \sqrt{n}) + (mnB)^{1/4} \right) \leq CB^{1/4} \sqrt{m} \quad (17)$$

for an absolute constant C .

Remark 20. The requirement that σ^2 is upper bounded is not essential. The conclusion in Corollary 19 works so long as fourth moment is bounded by B .

Let $\mathbb{M}s_1(A)$ be the median of $s_1(A)$. Following a calculation from [19], we have

$$\begin{aligned} |\mathbf{E}[s_1(A)] - \mathbb{M}s_1(A)| &\leq \mathbf{E}[|s_1(A) - \mathbb{M}s_1(A)|] \\ &= \int_0^\infty \mathbf{Pr}[|s_1(A) - \mathbb{M}s_1(A)| \geq t] dt \\ &\leq 4 \int_0^\infty e^{-t^2/4D^2} dt = 4D\sqrt{\pi}, \end{aligned}$$

where $D \leq 1$ for Bernoulli random variables that we consider. This allows us to conclude Theorem 7. \blacksquare

A.2 Correctness of Classification for the Simple Algorithm

We now prove correctness of our algorithm. We first show how to choose T for Procedure Classify. Let B denote the fourth moment bound for a single random variable in the mean zero random matrix $X - \mathcal{X}$; for the type of normalized Bernoulli r.v.s that we care about, \sqrt{B} is in the order of σ^2 , where σ^2 is defined in Theorem 2.

Let $N\gamma$ be a large enough constant. Let $s_1(X - \mathcal{X}) \leq C_0\sqrt{K}$, where $C_0 = C_1B^{1/4}$ as defined in Theorem 7 and let the threshold

$$T = \sqrt{C_3KN\gamma} \geq 15C_0\sqrt{K}, \quad (18)$$

which requires that

$$C_3N\gamma \geq 225C_0^2, \text{ where } C_3 \text{ satisfies (22).} \quad (19)$$

Following Lemma 27, (37), (39), and Proposition 25, we have

$$|s_2(\mathcal{X}) - s_2(X)| \leq s_1(X - \mathcal{X}) \leq C_0\sqrt{K}. \quad (20)$$

We have two cases,

1. When $s_2(X) \leq T$, by Lemma 10 and the fact that $s_2(\mathcal{X}) \leq s_2(X) + s_1(X - \mathcal{X}) \leq T + C_0\sqrt{K} \leq \frac{16T}{15}$, we have

$$s_2(\mathcal{X})^2 |x_2 - y_2|^2 \leq \frac{256T^2}{225} \frac{C_{\max}}{2N} \leq \frac{128C_3K\gamma C_{\max}}{225} \leq \frac{128C_3K\gamma 4}{225\omega_{\min}} \quad (21)$$

for C_{\max} as defined in Lemma 10. We want $s_2(\mathcal{X})^2|x_2 - y_2|^2 \leq \frac{3K\gamma}{4}$. This holds so long as $\frac{128C_3K\gamma C_{\max}}{225} \leq \frac{128C_3K\gamma 4}{225\omega_{\min}} \leq \frac{3K\gamma}{4}$, which is true if

$$C_3 \leq \frac{675\omega_{\min}}{2048}; \text{ thus we take } C_3 = \frac{675\omega_{\min}}{2048} \text{ from this point on.} \quad (22)$$

It follows from Lemma 11 that $s_1(\mathcal{X})^2|x_1 - y_1|^2 \geq \frac{K\gamma}{4}$. Hence by (13)

$$|x_1 - y_1| \geq \frac{\sqrt{K\gamma}}{2s_1(\mathcal{X})} \geq \frac{\sqrt{K\gamma}}{2\sqrt{2NK}} \geq \frac{1}{2}\sqrt{\frac{\gamma}{2N}}. \quad (23)$$

Thus the condition of Theorem 21 holds with $c_2 = \frac{1}{2}$, so long as

$$N\gamma \geq \frac{2048C_1^2\sqrt{B}}{3\omega_{\min}}, \quad (24)$$

due to (19) and (22); This is a weaker condition than (35) for $f < \frac{1}{2}$.

2. When $s_2(X) \geq T$, we have $s_2(\mathcal{X}) \geq s_2(X) - s_1(X - \mathcal{X}) \geq T - C_0\sqrt{K} \geq \frac{14T}{15}$;
This satisfies the condition of Theorem 23, with $c_3 = \frac{14\sqrt{C_3}}{15} = \frac{7}{16}\sqrt{\frac{3\omega_{\min}}{2}}$.

Let us first denote the first singular vector u_1 and its ‘‘noise’’ vector ϵ as follows:

$$u_1^T = (x + \delta_1, \dots, x + \delta_{N_1}, y + \tau_1, \dots, y + \tau_{N_2}), \quad \epsilon^T = (\delta_1, \dots, \delta_{N_1}, \tau_1, \dots, \tau_{N_2}).$$

It turns out that we only need to use the mixture mean

$$M = \frac{\sum_{i=1}^{N_1}(x + \delta_i) + \sum_{i=1}^{N_2}(y + \tau_i)}{2N} \quad (25)$$

to decide which side to put a node, i.e., to partition $j \in [2N]$ according to $u_{1,j} < M$ or $u_{1,j} \geq M$, given that N_1/N_2 is a constant; Misclassifying any entry will contribute $\Omega\left(\frac{\gamma}{2N}\right)$ amount to $\|\bar{u}_1 - u_1\|_2^2$.

Theorem 21. *Assume w.l.o.g. that $N_1 \leq N_2$ and $2N \leq K$. Let $\omega_1 = N_1/2N$ and $\omega_2 = N_2/2N$. Suppose $|x_1 - y_1| \geq c_2\sqrt{\frac{\gamma}{2N}}$ for some constant $c_2 = \frac{1}{2}$. By requiring $N \geq \frac{2048C_1^2\sqrt{B}}{3\gamma\omega_{\min}}$ as in (24), and*

$$N_1 \geq \frac{2c_1^2\sigma^2}{fc_2^2\gamma\omega_1\omega_2}, \text{ or equivalently } 2N \geq \frac{2c_1^2\sigma^2}{fc_2^2\gamma\omega_2\omega_1^2} = \frac{25C_1^2\sqrt{B}}{fc_2^2\gamma\omega_2\omega_1^2}, \quad (26)$$

where $c_1 = \frac{5C_1B^{1/4}}{\sqrt{2c_0}\sigma}$ for C_1 specified in Theorem 7 and c_0 specified in Proposition 8, we can classify the two population using the mixture mean M with the error factor at most f for N_1, N_2 respectively whp.

By Lemma 4 and Theorem 7, we immediately have the following claim.

Claim. For c_1 chosen as in Theorem 21, $\|\epsilon\|_2^2 = \sum_{i=1}^{N_1} \delta_i^2 + \sum_{i=1}^{N_2} \tau_i^2 \leq \frac{c_1^2\sigma^2}{N}$.

Proof. Given that $c_1 = \frac{5C_1 \sqrt[4]{B}}{\sqrt{2}c_0\sigma}$ such that C_1 appears in Theorem 7 and c_0 appears in Proposition 8,

$$\begin{aligned} \sqrt{\sum_{i=1}^{N_1} \delta_i^2 + \sum_{i=1}^{N_2} \tau_i^2} &= \|u_1 - \bar{u}_1\|_2 \sim 2\theta_1 \sim 2\sin(\theta_1) \\ &\leq \frac{4s_1(X - \mathcal{X})}{\text{gap}(1, \mathcal{X})} \leq \frac{4C_1 \sqrt[4]{B}\sqrt{K}}{4c_0\sqrt{2NK}/5} = \frac{c_1\sigma}{\sqrt{N}}. \end{aligned}$$

This allows us to conclude the claim. \blacksquare

We need the following lemma, proof of which appears in Appendix C.

Lemma 22. *Assume that $2N \leq K$ and Condition (26) in Theorem 21, we have*

$$|M - x| \geq \frac{N_2(1 - \sqrt{\gamma})|y - x|}{2N}, \quad |y - M| \geq \frac{N_1(1 - \sqrt{\gamma})|y - x|}{2N}. \quad (27)$$

Proof of Theorem 21. Recall that the largest \bar{u}_1, \bar{u}_2 have the form of $[x, \dots, x, y, \dots, y]$, where x repeats N_1 times and y repeats N_2 times; hence w.l.o.g., assume that $x < y$, we have

$$\forall i, s.t. x + \delta_i > M, \text{ it contributes } \delta_i^2 > |M - x|^2 \geq \frac{N_2^2 c_2^2 \gamma}{8N^3} \text{ to } \|\epsilon\|_2^2, \quad (28)$$

$$\forall i, s.t. y - \tau_i < M, \text{ it contributes } \delta_i^2 > |M - y|^2 \geq \frac{N_2^2 c_2^2 \gamma}{8N^3} \text{ to } \|\epsilon\|_2^2. \quad (29)$$

Hence the total number of entries that goes above M from P_1 , and those goes below M from P_2 can not be too many since their total contribution is upper bounded by $\|\epsilon\|_2^2 = \|u_1 - \bar{u}_1\|_2^2$. Let ℓ_1 be the number of misclassified entries from N_1 , i.e., those described in (28), by Lemma 4,

$$\ell_1 \frac{N_2^2 c_2^2 \gamma}{8N^3} \leq \ell_1 |M - x|^2 \leq \|\epsilon\|_2^2 \leq \frac{c_1^2 \sigma^2}{N}. \quad (30)$$

Thus given that $N_1 \geq \frac{8c_1^2 \sigma^2}{f c_2^2 \gamma} \geq \frac{2c_1^2 \sigma^2}{f \omega_2^2 c_2^2 \gamma}$; hence it suffices to guarantee that $\ell_1 \leq \frac{2c_1^2 \sigma^2}{\omega_2^2 c_2^2 \gamma} \leq f N_1$.

We next bound the number of entries from P_2 that goes below M , which can not be too many either; let ℓ_2 be the number of misclassified entries from P_2 ,

$$\ell_2 \frac{N_1^2 c_2^2 \gamma}{8N^3} \leq \ell_2 |M - y|^2 \leq \|\epsilon\|_2^2 \leq \frac{c_1^2 \sigma^2}{N}, \quad (31)$$

hence by requiring

$$N_2 \geq \frac{2c_1^2 \sigma^2}{f \omega_1^2 c_2^2 \gamma}, \quad (32)$$

it suffices to guarantee that $\ell_2 \leq \frac{2c_1^2\sigma^2}{\omega_2^2 c_2^2 \gamma} \leq fN_2$.

Condition (32) is equivalent to

$$N_1 = \frac{N_2\omega_1}{\omega_2} \geq \frac{2c_1^2\sigma^2}{f\omega_1\omega_2 c_2^2 \gamma}, \quad (33)$$

Thus by requiring

$$N_1 \geq \frac{2c_1^2\sigma^2}{fc_2^2\gamma\omega_2\omega_1}, \quad (34)$$

we have satisfied all requirements. \blacksquare

Combining Lemma 4 and Corollary 12, we have

Claim. Given that $s_2(\mathcal{X}) \geq c_3\sqrt{KN\gamma}$, $\|u_2 - \bar{u}_2\|_2^2 \leq \frac{16s_1(X-\mathcal{X})^2}{s_2(\mathcal{X})^2} \leq \frac{16C_0^2K}{c_3^2KN\gamma}$.

This allows us to prove the following theorem. Let the classification error factor be the number of misclassified individuals from one group over total amount of people in that group.

Theorem 23. Assume $N_1 \leq N_2$ and $2N \leq K$. Let $\omega_1 = N_1/2N$ and $\omega_2 = N_2/2N$. Let $s_2(\mathcal{X}) \geq c_3\sqrt{KN\gamma}$, where $c_3 = \frac{7}{16}\sqrt{\frac{3\omega_{\min}}{2}}$ and ω_{\min} is the minimum possible weight allowed by the algorithm. By requiring

$$2N \geq \frac{360C_0^2}{\omega_{\min}f\gamma} \left(\frac{\omega_2}{\omega_1} + 1 \right) = \Theta \left(\frac{\sigma^2}{f\gamma\omega_{\min}\omega_1} \right), \quad (35)$$

we can classify the two population using 0 to separate components in u_2 , with error factor at most f for both P_1, P_2 whp.

Proof. Let ℓ_1, ℓ_2 be the number of misclassified entries from P_1 and P_2 respectively; they each contribute at least $\frac{C_{x \min}}{2N}$, and $\frac{C_{y \min}}{2N}$ amount to $\|u_2 - \bar{u}_2\|_2^2$, and hence by Claim A.2,

$$\ell_1 \frac{C_{x \min}}{2N} \leq \|u_2 - \bar{u}_2\|_2^2 \leq \frac{16C_0^2K}{c_3^2KN\gamma} \leq \frac{16C_0^2}{c_3^2N\gamma}. \quad (36)$$

Hence $\ell_1 \leq \frac{32C_0^2}{c_3^2\gamma C_{x \min}} \leq fN_1$ given that $N \geq \frac{16C_0^2}{c_3^2f\gamma} \left(4\frac{\omega_1}{\omega_2} + 1 \right)$.

Similarly, by Claim A.2, we have $\ell_2 \frac{C_{y \min}}{2N} \leq \|u_2 - \bar{u}_2\|_2^2$ and thus $\ell_2 \leq \frac{32C_0^2}{c_3^2\gamma C_{y \min}} \leq \frac{N_2}{f}$ so long as $N \geq \frac{16C_0^2}{c_3^2f\gamma} \left(4\frac{\omega_2}{\omega_1} + 1 \right)$; the bound on $2N$ follows by plugging in $c_3 = \frac{7}{16}\sqrt{\frac{3\omega_{\min}}{2}}$. \blacksquare

Finally,

Theorem 24. Given a set of $n \geq \Omega \left(\frac{\sigma^2}{\gamma f \omega_{\min}} \right)$ individuals, by trying Procedure Classify for $\log n$ rounds, with probability of error at each round for each individual being $f = 1/10$, where each round we take a set of $K > n$ independent features, and by taking majority vote over the different runs for each sample, our algorithm will find the correct partition with probability $1 - 1/n^2$.

Proof. A sample is put in the wrong side with a probability $1/10$ at each round. Let \mathcal{E}_i be the event that sample i is misclassified for more than $\log n$ times, thus $\Pr[\mathcal{E}_i] = \left(\frac{1}{10}\right)^{\log n} \leq 1/n^{3.32}$; hence by union bound, with probability $1 - 1/n^2$, none of the $2N$ individuals is misclassified. ■

B More Proofs for the Simple Algorithm Classify

B.1 Proof of Lemma 4

Let $u_1, \dots, u_n, v_1, \dots, v_n$ be the n left and right singular vectors of X , corresponding to $s_1(X) \geq s_2(X) \geq \dots \geq s_n(X)$, we have for $\forall i, \|u_i\|_2 = 1, \|v_i\|_2 = 1$ such that $X^T u_i = s_i(X)v_i$ and $X v_i = s_i(X)u_i$.

Before we prove Lemma 4, given an $n \times K$ matrix X , where $n < K$, let us first define $H = X X^T$ and a block matrix

$$Y = \begin{bmatrix} 0 & X \\ X^T & 0 \end{bmatrix}_{(2N+K) \times (2N+K)}. \quad (37)$$

Recall that singular values of a real $n \times K$ matrix X are exactly the nonnegative square roots of the n largest eigenvalues of $H = X X^T$, i.e, $s_i(X) = \sqrt{\lambda_i(H)}, \forall i = 1, \dots, n$, given that

$$H u_i = X X^T u_i = s_i(X) X v_i = s_i^2(X) u_i. \quad (38)$$

Hence the left singular vectors u_1, \dots, u_n of X are eigenvectors of H corresponding to $\lambda_i(H) = s_i^2(X)$.

We next show that the first n eigenvalues of Y and their corresponding eigenvectors:

$$Y \begin{bmatrix} u_i \\ v_i \end{bmatrix} = \begin{bmatrix} 0 & X \\ X^T & 0 \end{bmatrix} \begin{bmatrix} u_i \\ v_i \end{bmatrix} = \begin{bmatrix} X v_i \\ X^T u_i \end{bmatrix} = \begin{bmatrix} s_i(X) u_i \\ s_i(X) v_i \end{bmatrix} = s_i(X) \begin{bmatrix} u_i \\ v_i \end{bmatrix}, \quad (39)$$

and hence

Proposition 25. *The largest n eigenvalues of Y are $s_1(X), \dots, s_n(X)$ with corresponding eigenvectors $[u_i, v_i], \forall i = 1, \dots, n$, where $u_i, v_i, \forall i$, are left and right singular vectors of X corresponding to $s_i(X)$.*

In fact both $\pm s_i(X)$ are eigenvalues of Y , which is irrelevant. *Proof of Lemma 4.* We first state a theorem, whose statement appears in a lecture note by Spielman [22], with a slight modification (off by a factor on RHS). Our proof for this theorem is included here for completeness. It is known that for any real symmetric matrix, there exist a set of n orthonormal eigenvectors.

Theorem 26. (Modified Version of Spielman [22]) *For A and M being two symmetric matrices and $E = M - A$. Let $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$ be eigenvalues of A , with orthonormal eigenvectors v_1, v_2, \dots, v_n and let $\lambda_1(M) \geq \lambda_2(M) \geq \dots \geq$*

$\lambda_n(M)$ be eigenvalues of M and w_1, w_2, \dots, w_n be the corresponding orthonormal eigenvectors of M , with $\theta_i = \angle(v_i, w_i)$. Then

$$\theta_i \sim \sin(\theta_i) \leq \frac{\|E - \Delta_i I\|_2}{\text{gap}(i, A)} \leq \frac{\|E\|_2 + |\Delta_i|}{\text{gap}(i, A)} \leq \frac{2\|E\|_2}{\text{gap}(i, A)} \quad (40)$$

where $\text{gap}(i, A) = \min_{j \neq i} |\lambda_i(A) - \lambda_j(A)|$ and $\Delta_i = \lambda_i(M) - \lambda_i(A)$.

Let us apply Theorem 26 to the symmetric matrix Y in (37). In particular, we only compare the first n eigenvectors of Y of \mathcal{Y} . For the numerator of RHS of (40), we have $E = Y - \mathcal{Y}$, and $\|E\|_2 = \|Y - \mathcal{Y}\|_2 = s_1(Y - \mathcal{Y})$ by a derivation similar to (39), where eigenvectors of E are concatenations of left and right singular vectors of $X - \mathcal{X}$; For the denominator, we have by Proposition 25, $\text{gap}(i, \mathcal{Y}) = \min_{j \neq i} |\lambda_i(\mathcal{Y}) - \lambda_j(\mathcal{Y})| = \min_{j \neq i} |s_i(\mathcal{X}) - s_j(\mathcal{X})|$. ■

We first prove the following claim.

Claim. For any symmetric $n \times n$ matrix A , let $\lambda_i, \forall i = 1, \dots, n$ be eigenvalues of A with orthonormal eigenvectors v_1, v_2, \dots, v_n , for all $y \perp v_i$,

$$\|(A - \lambda_i)y\|_2 \geq \min_{j \neq i} |\lambda_i - \lambda_j| \|y\|_2. \quad (41)$$

Proof. Let us first assume $y \perp v_i$ and write $y = \sum_{j=1, j \neq i}^n c_j v_j$, thus we have $\|y\|_2 = \sqrt{\sum_{j=1, j \neq i}^n c_j^2}$ and

$$\begin{aligned} \|(A - \lambda_i)y\|_2 &= \left\| \sum_{j=1, j \neq i}^n c_j (A - \lambda_i)v_j \right\|_2 \\ &= \left\| \sum_{j=1, j \neq i}^n c_j (\lambda_j - \lambda_i)v_j \right\|_2 \\ &= \sqrt{\sum_{j=1, j \neq i}^n c_j^2 |\lambda_j - \lambda_i|^2} \\ &\geq \min_{j \neq i} |\lambda_i - \lambda_j| \sqrt{\sum_{j=1, j \neq i}^n c_j^2} = \min_{j \neq i} |\lambda_i - \lambda_j| \|y\|_2. \end{aligned}$$

■

Proof of Theorem 26. Let us construct a vector y that is orthogonal to v_i as follows:

$$y = w_i - (v_i^T w_i)v_i \quad (42)$$

By Claim B.1, we have

$$\|(A - \lambda_i(A))y\|_2 \geq \min_{j \neq i} |\lambda_i(A) - \lambda_j(A)| \|y\|_2, \quad (43)$$

and hence

$$\|y\|_2 \leq \frac{\|(A - \lambda_i(A))y\|_2}{\min_{j \neq i} |\lambda_i(A) - \lambda_j(A)|} \quad (44)$$

On the other hand,

$$\begin{aligned} \|(A - \lambda_i(A))y\|_2 &= \|(A - \lambda_i(A))(w_i - (v_i^T w_i)v_i)\|_2 \\ &= \|(A - \lambda_i(A))w_i\|_2 \\ &= \|(M - E - \lambda_i(A))w_i\|_2 \\ &= \|(\lambda_i(M) - \lambda_i(A))w_i - Ew_i\|_2 \\ &= \|(\Delta_i I - E)w_i\|_2 \leq \|E - \Delta_i I\|_2 \\ &\leq \|E\|_2 + |\Delta_i| \end{aligned}$$

Finally, given that $\|w\|_2 = 1$,

$$\begin{aligned} \sin(\theta_i) &= \frac{\|y\|_2}{\|w\|_2} \leq \frac{\|(A - \lambda_i(A))y\|_2}{\min_{j \neq i} |\lambda_i(A) - \lambda_j(A)|} \\ &\leq \frac{\|E\|_2 + |\Delta_i|}{\text{gap}(i, A)}. \end{aligned}$$

Lemma 27. $\forall i = 1, \dots, n, |\Delta_i| \leq \|E\|_2$.

Proof. Let S_j be a subspace of dimension j . Recall the following definitions of λ_i for a matrix:

$$\lambda_i(M) = \inf_{S_{N-i+1}} \sup_{x \in S_{N-i+1}, \|x\|_2=1} x^T M x. \quad (45)$$

In the following, let S_{N-i+1}^v be the subspace that is orthogonal to the subset of orthonormal eigenvectors v_1, \dots, v_{i-1} of symmetric matrix A . Note that this is the $N - i + 1$ dimensional subspace that achieves the minimum of the maximum of $v^T A v$ over all unit-length vectors v in the particular subspace.

$$\begin{aligned} \lambda_i(M) &= \inf_{S_{N-i+1}} \sup_{x \in S_{N-i+1}, \|x\|_2=1} x^T M x \leq \sup_{x \in S_{N-i+1}^v, \|x\|_2=1} x^T M x \\ &\leq \sup_{x \in S_{N-i+1}^v, \|x\|_2=1} x^T (A + E)x \\ &\leq \sup_{v \in S_{N-i+1}^v, \|v\|_2=1} v^T A v + \sup_{x \in \mathbb{R}^n, \|x\|_2=1} |x^T E x| \\ &= \lambda_i(A) + \|E\|_2. \end{aligned}$$

For the other direction, let S_{N-i+1}^w be the subspace that is orthogonal to the subset of orthonormal eigenvectors w_1, \dots, w_{i-1} of symmetric matrix M . Note that this is the $N - i + 1$ dimensional subspace that achieves the minimum of the maximum of $w^T M w$ over all unit-length vectors w in the particular subspace.

$$\begin{aligned}
\lambda_i(A) &= \inf_{S_{N-i+1}} \sup_{x \in S_{N-i+1}, \|x\|_2=1} x^T A x \leq \sup_{x \in S_{N-i+1}^w, \|x\|_2=1} x^T A x \\
&\leq \sup_{x \in S_{N-i+1}^w, \|x\|_2=1} x^T (M + (-E))x \\
&\leq \sup_{w \in S_{N-i+1}^w, \|w\|_2=1} w^T M w + \sup_{x \in \mathbb{R}^n, \|x\|_2=1} x^T (-E)x \\
&\leq \sup_{w \in S_{N-i+1}^w, \|w\|_2=1} w^T M w + \sup_{x \in \mathbb{R}^n, \|x\|_2=1} |x^T (-E)x| \\
&= \lambda_i(M) + \|E\|_2,
\end{aligned}$$

where $\|E\|_2 = \|-E\|_2$.

Thus we have $-\|E\|_2 \leq \lambda_i(M) - \lambda_i(A) \leq \|E\|_2$, and hence $|\Delta_i| \leq \|E\|_2$. ■

Therefore, $\sin(\theta_i) \leq \frac{\|E\|_2 + |\Delta_i|}{\text{gap}(i, A)} \leq \frac{2\|E\|_2}{\text{gap}(i, A)}$. ■

B.2 Some Propositions Regarding the Static Matrices

For static matrix $\mathcal{H} = \mathcal{X}\mathcal{X}^T$ and $\mathcal{Y} = \begin{bmatrix} 0 & \mathcal{X} \\ \mathcal{X}^T & 0 \end{bmatrix}$, we define

$$\begin{aligned}
\text{gap}(\mathcal{H}) &= |\lambda_1(\mathcal{H}) - \lambda_2(\mathcal{H})|, \\
\text{gap}(\mathcal{Y}) &= |\lambda_1(\mathcal{Y}) - \lambda_2(\mathcal{Y})| = \frac{\text{gap}(\mathcal{H})}{\lambda_1(\mathcal{Y}) + \lambda_2(\mathcal{Y})},
\end{aligned}$$

Proposition 28. For static matrix \mathcal{Y} , let

$$\text{gap}(\mathcal{Y}) = |\lambda_1(\mathcal{Y}) - \lambda_2(\mathcal{Y})| = \frac{\text{gap}(\mathcal{H})}{\lambda_1(\mathcal{Y}) + \lambda_2(\mathcal{Y})}, \quad (46)$$

we have

$$\begin{aligned}
\sqrt{\max\{N_1 a, N_2 c\}} &\leq \lambda_1(\mathcal{Y}) \leq \sqrt{N_1 a + N_2 c} \\
\sqrt{N_1 a + N_2 c} &\leq \lambda_1(\mathcal{Y}) + \lambda_2(\mathcal{Y}) \leq \sqrt{2(N_1 a + N_2 c)} \\
\sqrt{\frac{N_1 N_2 (ac - b^2)}{N_1 a + N_2 c}} &\leq \lambda_2(\mathcal{Y}) \leq \sqrt{\frac{2N_1 N_2 (ac - b^2)}{N_1 a + N_2 c}}
\end{aligned}$$

Thus we have

$$\text{gap}(\mathcal{Y}) = \Theta \left(\frac{\text{gap}(\mathcal{H})}{\sqrt{N_1 a + N_2 c}} \right) = \Theta \left(\frac{\sqrt{(N_1 a + N_2 c)^2 - 4N_1 N_2 (ac - b^2)}}{\sqrt{N_1 a + N_2 c}} \right).$$

Proof. We first show the following:

Proposition 29. For static matrix $\mathcal{H} = \mathcal{X}\mathcal{X}^T$ as in (15), Let $\lambda_1(\mathcal{H}), \lambda_2(\mathcal{H})$ be the non-zero eigenvalues of \mathcal{H} , and denote $\text{gap}(\mathcal{H}) = |\lambda_1(\mathcal{H}) - \lambda_2(\mathcal{H})|$.

$$\lambda_1(\mathcal{H}) = \frac{N_1a + N_2c + \sqrt{(N_1a - N_2c)^2 + 4N_1N_2b^2}}{2}, \quad (47)$$

$$\lambda_2(\mathcal{H}) = \frac{N_1a + N_2c - \sqrt{(N_1a - N_2c)^2 + 4N_1N_2b^2}}{2}, \quad (48)$$

$$|N_1a - N_2c| \leq \text{gap}(\mathcal{H}) \leq N_1a + N_2c, \quad (49)$$

where $\lambda_2(\mathcal{H}) = 0$, when $ac = b^2$ and $\text{gap}(\mathcal{H}) = N_1a + N_2c$.

Proof. Let $\mathcal{H} = \mathcal{X}\mathcal{X}^T$. Rank of \mathcal{H} is at most 2. Therefore there exists at most two non-zero eigenvalues λ_1, λ_2 for \mathcal{H} , with corresponding nonzero eigenvectors v_1, v_2 being constant on each population. This is true because if we multiply $\mathcal{H} = \mathcal{X}\mathcal{X}^T$ by a permutation matrix P to exchange two rows among the same population, we have $P\mathcal{H}v_i = \lambda_i P v_i, \forall i = 1, 2$; given that $P\mathcal{H}v_i = \mathcal{H}v_i$, we deduce that $Pv_i = v_i$ for non-zero λ_i . Hence v_i must be constant on each population.

Let the top two eigenvector v_1, v_2 be of form $[x, \dots, x, y, \dots, y]$, where x repeats N_1 times and y repeats N_2 times; Note that they corresponds to \bar{u}_1 and \bar{u}_2 of \mathcal{X} following a derivation similar to (38).

We thus have the following equations:

$$N_1ax + N_2by = \lambda x, \quad (50)$$

$$N_1bx + N_2cy = \lambda y, \quad (51)$$

which can be written in a matrix form:

$$\begin{bmatrix} N_1a - \lambda & N_2b \\ N_1b & N_2c - \lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 0 \quad (52)$$

Given that

$$\begin{bmatrix} x \\ y \end{bmatrix} \neq 0,$$

the matrix is not one-to-one and therefore

$$D \begin{bmatrix} N_1a - \lambda & N_2b \\ N_1b & N_2c - \lambda \end{bmatrix} = 0.$$

By solving $(N_1a - \lambda)(N_2c - \lambda) - N_1N_2b^2 = 0$, we get $\lambda_1(\mathcal{H}), \lambda_2(\mathcal{H})$ and $\text{gap}(\mathcal{H})$.

We next derive an upper bound on $\text{gap}(\mathcal{H})$.

$$\begin{aligned} \text{gap}(\mathcal{H}) &= \sqrt{(N_1a - N_2c)^2 + 4N_1N_2b^2} \\ &= \sqrt{(N_1a + N_2c)^2 - 4N_1N_2ac + 4N_1N_2b^2} \\ &\leq \sqrt{(N_1a + N_2c)^2} \\ &\leq N_1a + N_2c, \end{aligned}$$

where $a, c \geq 0$ and $ac \geq b^2$ as in Proposition 17.

It is easy to see that

$$\text{gap}(\mathcal{H}) \geq |N_1a - N_2c|, \quad (53)$$

given that $b^2 \geq 0$. ■

Thus we have

$$\begin{aligned} \max\{N_1a, N_2c\} &\leq \lambda_1(\mathcal{H}) \leq N_1a + N_2c, \\ 0 &\leq \lambda_2(\mathcal{H}) \leq \min\{N_1a, N_2c\}, \\ \lambda_1(\mathcal{H}) + \lambda_2(\mathcal{H}) &= N_1a + N_2c, \\ \lambda_1(\mathcal{H})\lambda_2(\mathcal{H}) &= N_1N_2(ac - b^2), \end{aligned}$$

Given that two largest eigenvalues of \mathcal{Y} , $\lambda_1(\mathcal{Y}) = \sqrt{\lambda_1(\mathcal{H})}$ and $\lambda_2(\mathcal{Y}) = \sqrt{\lambda_2(\mathcal{H})}$ for $\mathcal{Y} = \begin{bmatrix} 0 & \mathcal{X} \\ \mathcal{X}^T & 0 \end{bmatrix}$, by Proposition 29 and the following fact:

$$\sqrt{(\lambda_1(\mathcal{Y}))^2 + \lambda_2(\mathcal{Y})^2} \leq \lambda_1(\mathcal{Y}) + \lambda_2(\mathcal{Y}) = \sqrt{2(\lambda_1(\mathcal{Y}))^2 + \lambda_2(\mathcal{Y})^2}, \quad (54)$$

we get all inequalities. ■

B.3 Proofs of Proposition 8 and 9

Proof of Proposition 8. We rewrite Proposition 28 given that, for a normalized \mathcal{X} , $\text{gap}(\mathcal{H}) \geq \frac{8c_0NK}{5}$, as Proposition 31 and $\lambda_j(\mathcal{Y}) = s_j(\mathcal{X})$. In particular,

$$\begin{aligned} \text{gap}(\mathcal{X}) = \text{gap}(\mathcal{Y}) &= \frac{\text{gap}(\mathcal{H})}{\lambda_1(\mathcal{Y}) + \lambda_2(\mathcal{Y})} \\ &\geq \frac{\text{gap}(\mathcal{H})}{\sqrt{N_1a + N_2c}} \geq \frac{8c_0NK}{5\sqrt{2NK}} \\ &\geq \frac{4\sqrt{2NK}}{5}. \end{aligned}$$

For the upper bound on $\text{gap}(\mathcal{X})$, we have that

$$\begin{aligned} \text{gap}(\mathcal{X}) = \text{gap}(\mathcal{Y}) &= \frac{\text{gap}(\mathcal{H})}{\lambda_1(\mathcal{Y}) + \lambda_2(\mathcal{Y})} \\ &\leq \frac{N_1a + N_2c}{\sqrt{N_1a + N_2c}} \\ &\leq \sqrt{N_1a + N_2c} \leq \sqrt{2NK} \end{aligned}$$

■

Definition 30. For our application, we have $\forall k, 1 \geq p_1^k, p_2^k \geq 0$, and

$$\mathcal{X} = \begin{bmatrix} \frac{1+p_1^1}{2} & \frac{1+p_1^2}{2} & \cdots & \frac{1+p_1^K}{2} \\ \cdots & \cdots & \cdots & \cdots \\ \frac{1+p_1^1}{2} & \frac{1+p_1^2}{2} & \cdots & \frac{1+p_1^K}{2} \\ \frac{1+p_2^1}{2} & \frac{1+p_2^2}{2} & \cdots & \frac{1+p_2^K}{2} \\ \cdots & \cdots & \cdots & \cdots \\ \frac{1+p_2^1}{2} & \frac{1+p_2^2}{2} & \cdots & \frac{1+p_2^K}{2} \end{bmatrix}_{2N \times K} \quad (55)$$

It is easy to see that with this normalized random matrix, $\lambda_1(\mathcal{H}) = \lambda_2(\mathcal{H})$ is not possible, given that $a, b, c \in [K/4, K]$; furthermore, $\text{gap}(\mathcal{H}) = \Theta(NK)$ as in the Proposition 31.

Proposition 31. Given $\mathcal{H} = \mathcal{X}\mathcal{X}^T$ and a, b, c as in (11) for any expected value mean matrix \mathcal{X} , which is not necessarily normalized,

$$\text{gap}(\mathcal{H}) = \sqrt{(N_1a - N_2c)^2 + 4N_1N_2b^2} \geq \frac{8c_0NK}{5}, \quad (56)$$

where $c_0 = \frac{|b|\sqrt{ac}}{K(a+c)}$.

Hence for a normalized \mathcal{X} , $\text{gap}(\mathcal{H}) = \Theta(NK)$ given that $a, b, c \in [K/4, K]$.

Proof. For a tighter lower bound of $\text{gap}(\mathcal{H})$ than the obvious $|N_1a - N_2c|$, let us assume w.l.o.g. that $N_2c \geq N_1a$. Thus we have

$$N_2 \geq 2N \frac{a}{a+c} \quad (57)$$

We differentiate two cases:

- Balanced case: $N_1a \geq \frac{4}{25}N_2c$.
- Imbalanced case: $N_1a \leq \frac{4}{25}N_2c$.

For balanced case: we have $N_1 \geq \frac{4}{25} \frac{N_2c}{a}$ and hence

$$\begin{aligned} \text{gap}(\mathcal{H}) &\geq \sqrt{4N_1N_2b^2} \geq \frac{4N_2|b|}{5} \sqrt{\frac{c}{a}} \\ &\geq \frac{8N|b|}{5} \frac{a}{a+c} \sqrt{\frac{c}{a}} \geq \frac{8N|b|}{5} \frac{\sqrt{ac}}{a+c} \\ &= \frac{8c_0NK}{5}, \end{aligned}$$

where $N_2 \geq 2N \frac{a}{a+c}$ as in (57).

For the imbalanced case, given that $\sqrt{ac} \geq |b|$ by Proposition 17,

$$\begin{aligned} \text{gap}(\mathcal{H}) &\geq \sqrt{(N_1a - N_2c)^2} \geq \frac{21}{25}N_2c \\ &\geq \frac{42}{25} \frac{Nac}{a+c} \geq \frac{8}{5} \frac{N|b|\sqrt{ac}}{a+c} \\ &\geq \frac{8c_0NK}{5}. \end{aligned}$$

Finally, for a normalized random matrix X and its \mathcal{X} , we have c_0 being a constant and combing with the upper bound of $\text{gap}(\mathcal{H}) \leq N_1a + N_2c \leq 2NK$ concludes that $\text{gap}(\mathcal{H}) = \Theta(NK)$. ■

Proof of Proposition 9. By (38), \bar{u}_1, \bar{u}_2 are the first and second eigenvectors of \mathcal{H} corresponding to $\lambda_1(\mathcal{H})$ and $\lambda_2(\mathcal{H})$. Let x, y be entries that correspond to P_1, P_2 respectively in the first or second eigenvectors of \mathcal{H} . By (50) and (51), we have

$$\frac{y}{x} = \frac{\lambda - N_1a}{N_2b} = \frac{N_1b}{\lambda - N_2c}.$$

In addition, given any $b \neq 0$, we have $\text{gap}(\mathcal{H}) > |N_1a - N_2c|$ and hence $\lambda_1(\mathcal{H}) > \max\{N_1a, N_2c\} > \lambda_2(\mathcal{H})$. Therefore, for $b > 0$, $\frac{y}{x} \geq 0$ for first eigenvector and ≤ 0 for v_2 . and for $b < 0$, it is the opposite. ■

B.4 Proof of Lemma 10

Proof of Lemma 10. We first show that $|x_2|, |y_2|$ are within a constant factor of each other, given that $\omega_1/\omega_2 = \frac{N_1}{N_2}$ is a constant.

Proposition 32. For a normalized \mathcal{X} , where $N_1, N_2, a, b \neq 0$, x_2, y_2 in the second top left singular vector \bar{u}_2 satisfy

$$\frac{2N_2}{N_1} \geq \frac{|x_2|}{|y_2|} \geq \frac{N_2}{2N_1} \quad (58)$$

Proof. By (50) and given the upper bound on $\text{gap}(\mathcal{H})$ in (49),

$$\frac{|y_2|}{|x_2|} = \frac{N_1a - \lambda_2}{N_2b} = \frac{N_1a - N_2c + \text{gap}(\mathcal{H})}{2N_2b} \leq \frac{N_1a}{N_2b}, \quad (59)$$

and hence $\frac{|x_2|}{|y_2|} \geq \frac{N_2b}{N_1a}$. By (51) and (49), we have

$$\frac{|x_2|}{|y_2|} = \frac{N_2c - \lambda_2}{N_1b} = \frac{N_2c - N_1a + \text{gap}(\mathcal{H})}{2N_1b} \leq \frac{N_2c}{N_1b} \quad (60)$$

We finish the proof by observing that

$$\frac{1}{2} \leq \frac{a}{b} \leq 2, \quad \frac{1}{2} \leq \frac{c}{b} \leq 2, \quad (61)$$

due to the fact that $\frac{1}{2} \leq \frac{\mu_i^j}{\mu_2^j} \leq 2, \forall j = 1, \dots, K$ for $\mu_i^j \in [1/2, 1]$ in a normalized \mathcal{X} , and the following lemma:

Lemma 33. If $0 < c_{\min} \leq \frac{a_i}{b_i} \leq c_{\max}, \forall i = 1, \dots, n$, where $a_i, b_i > 0$, then $c_{\min} \leq \frac{\sum_{i=1}^n a_i}{\sum_{i=1}^n b_i} \leq c_{\max}$.

Proof.

$$c_{\min} = \frac{\sum_{i=1}^n c_{\min} b_i}{\sum_{i=1}^n b_i} \leq \frac{\sum_{i=1}^n a_i}{\sum_{i=1}^n b_i} \leq \frac{\sum_{i=1}^n c_{\max} b_i}{\sum_{i=1}^n b_i} = c_{\max} \quad (62)$$

■
■

Let $x = x_2$ and $y = y_2$. By Proposition 32, $|y| \leq \frac{2N_1|x|}{N_2}$ and

$$1 = N_1x^2 + N_2y^2 \leq N_1x^2 + N_2 \left(\frac{2|x|N_1}{N_2} \right)^2 \leq x^2 \left(\frac{4N_1^2 + N_1N_2}{N_2} \right), \quad (63)$$

hence for $C_{x \min} = \frac{\omega_2}{4\omega_1^2 + \omega_1\omega_2}$,

$$x^2 \geq \frac{\omega_2}{4\omega_1^2 + \omega_1\omega_2} \frac{1}{2N}. \quad (64)$$

Looking in the other direction, by Proposition 32, $|x| \leq \frac{2|y|N_2}{N_1}$,

$$1 = N_1x^2 + N_2y^2 \leq N_2y^2 + N_1 \left(\frac{2|y|N_2}{N_1} \right)^2 \leq y^2 \left(\frac{N_2N_1 + 4N_2^2}{N_1} \right),$$

and hence for a given $C_{y \min} = \frac{\omega_1}{4\omega_2^2 + \omega_1\omega_2}$,

$$|y|^2 \geq \frac{\omega_1}{4\omega_2^2 + \omega_1\omega_2} \frac{1}{2N}. \quad (65)$$

On the other hand, by Proposition 32, we have $|y| \geq \frac{N_1|x|}{2N_2}$, we have

$$1 = N_1x^2 + N_2y^2 \geq N_1x^2 + N_2 \left(\frac{|x|N_1}{2N_2} \right)^2 \geq x^2 \left(\frac{N_1^2 + 4N_1N_2}{4N_2} \right),$$

and thus

$$x^2 \leq \frac{4\omega_2}{\omega_1^2 + 4\omega_1\omega_2} \frac{1}{2N}. \quad (66)$$

Looking in the other direction, by Proposition 32, $|x| \geq \frac{|y|N_2}{2N_1}$,

$$1 = N_1x^2 + N_2y^2 \geq N_2y^2 + N_1 \left(\frac{|y|N_2}{2N_1} \right)^2 \geq y^2 \left(\frac{4N_2N_1 + N_2^2}{4N_1} \right),$$

and hence $|y|^2 \leq \frac{4\omega_1}{\omega_2^2 + 4\omega_1\omega_2} \frac{1}{2N}$. Hence we have that

$$\begin{aligned} |x - y|^2 &= (|x| + |y|)^2 \leq \left(\sqrt{\frac{4\omega_2}{\omega_1^2 + 4\omega_1\omega_2} \frac{1}{2N}} + \sqrt{\frac{4\omega_1}{\omega_2^2 + 4\omega_1\omega_2} \frac{1}{2N}} \right)^2 \\ &\leq \frac{1}{2N} \left(\sqrt{\frac{4\omega_2}{\omega_1^2 + 4\omega_1\omega_2}} + \sqrt{\frac{4\omega_1}{\omega_2^2 + 4\omega_1\omega_2}} \right)^2, \end{aligned}$$

and $C_{\max} = (\sqrt{\frac{1}{\omega_1}} + \sqrt{\frac{1}{\omega_2}})^2$. Hence

$$C_{\max} \leq \left(\sqrt{\frac{4\omega_2}{\omega_1^2 + 4\omega_1\omega_2}} + \sqrt{\frac{4\omega_1}{\omega_2^2 + 4\omega_1\omega_2}} \right)^2 \leq \left(\sqrt{\frac{1}{\omega_1}} + \sqrt{\frac{1}{\omega_2}} \right)^2.$$

■

C Proof of Lemma 22

Recall that the largest left singular vectors u_1, u_2 has the form of $[x, \dots, x, y, \dots, y]$, where x repeats N_1 times and y repeats N_2 times. *Proof* of Lemma 22. Let us define the following random variables,

$$\delta = \frac{1}{N_1} \sum_{i=1}^{N_1} \delta_i, \quad \tau = \frac{1}{N_2} \sum_{i=1}^{N_2} \tau_i, \quad (67)$$

such that by Claim A.2,

$$|\delta| = \left| \frac{1}{N_1} \sum_{i=1}^{N_1} \delta_i \right| \leq \frac{1}{N_1} \sum_{i=1}^{N_1} |\delta_i| \leq \frac{\sqrt{N_1 \sum_{i=1}^{N_1} \delta_i^2}}{N_1} \leq \frac{c_1 \sigma}{\sqrt{N_1 N}}$$

$$|\tau| = \left| \frac{1}{N_2} \sum_{i=1}^{N_2} \tau_i \right| \leq \frac{1}{N_2} \sum_{i=1}^{N_2} |\tau_i| \leq \frac{\sqrt{N_2 \sum_{i=1}^{N_2} \tau_i^2}}{N_2} \leq \frac{c_1 \sigma}{\sqrt{N_2 N}}$$

and hence

$$\max(|N_1 \delta|, |N_2 \tau|) \leq \frac{c_1 \sigma \sqrt{N_2}}{\sqrt{N}} \quad (68)$$

given that we always assume that $N_2 > N_1$. A natural classifier to separate individuals would be: $\frac{x+y}{2}$ when we use u_1 ; but we do not have access to x and y . Recall that

$$M = \frac{\sum_{i=1}^{N_1} (x + \delta_i) + \sum_{i=1}^{N_2} (y + \tau_i)}{2N} = \frac{N_1 x + N_2 y}{2N} + \frac{N_1 \delta + N_2 \tau}{2N}.$$

We are now ready to show that when N_1, N_2 are large enough, we see enough separation between the mixture sample mean and both x and y . We first prove the following claims.

Claim. $xN_1\delta + yN_2\tau = -\frac{\|\epsilon\|_2^2}{2}$.

Proof. This claim is obvious given that $\|u_1\|_2 = \|\bar{u}_1\|_2 = 1$, and \bar{u}_1, u_1, ϵ all being real vectors,

$$\|u_1\|_2^2 = \|\bar{u}_1\|_2^2 + \|\epsilon\|_2^2 + 2 \langle \bar{u}_1, \epsilon \rangle = \|\bar{u}_1\|_2^2 + \|\epsilon\|_2^2 + 2xN_1\delta + yN_2\tau. \quad (69)$$

■

We next use $\frac{1}{\sqrt{2N}} |xN_1\delta + yN_2\tau|$ to obtain a bound on $|\frac{N_1\delta + N_2\tau}{2N}|$, given that

$$\frac{1}{\sqrt{2N}} |xN_1\delta + yN_2\tau| \leq \frac{\|\epsilon\|_2^2}{2\sqrt{2N}} \leq \frac{c_1^2\sigma^2}{2N\sqrt{2N}}; \quad (70)$$

Claim. Let $N_1 \leq N_2$, and $\omega_1 = \frac{N_1}{2N}$ and $\omega_2 = \frac{N_2}{2N}$, and given that $N_1 \geq \max\left(\frac{2c_1^2\sigma^2}{c_2\gamma}, \frac{8c_1^2\sigma^2\omega_2}{\gamma\omega_1}\right)$, we have

$$\left| \frac{N_1\delta + N_2\tau}{2N} \right| \leq \frac{N_1 |y - x| \sqrt{\gamma}}{2N} \quad (71)$$

Proof. We next derive a bound on $\frac{N_1\delta + N_2\tau}{2N}$. By Separation Lemma 11, we have $|x - y| = c_2\sqrt{\frac{\gamma}{2N}}$ for a constant $c_2 = 1/2$, and thus we have $\frac{\max(x,y)}{\sqrt{2N}} > \frac{1}{2N}$. Therefore,

$$\begin{aligned} \frac{|xN_1\delta + yN_2\tau|}{\sqrt{2N}} &= \frac{|\max(x,y)(N_1\delta + N_2\tau) + (x - \max(x,y))N_1\delta + (y - \max(x,y))N_2\tau|}{\sqrt{2N}} \\ &\geq \frac{|\max(x,y)|(N_1\delta + N_2\tau)|}{\sqrt{2N}} - \frac{|x - y| \max(|N_1\delta|, |N_2\tau|)}{\sqrt{2N}} \end{aligned}$$

Thus we have, given (68), (70) and (72),

$$\begin{aligned} \left| \frac{N_1\delta + N_2\tau}{2N} \right| &\leq \frac{|\max(x,y)|}{\sqrt{2N}} |(N_1\delta + N_2\tau)| \\ &\leq \frac{|xN_1\delta + yN_2\tau|}{\sqrt{2N}} + \frac{|x - y| \max(|N_1\delta|, |N_2\tau|)}{\sqrt{2N}} \\ &\leq \frac{c_1^2\sigma^2}{2N\sqrt{2N}} + \frac{c_2\sqrt{\gamma}}{2N} c_1\sigma\sqrt{\frac{N_2}{N}} \\ &\leq \frac{N_1c_2\gamma}{2N\sqrt{2N}} \leq \frac{N_1|y - x|\sqrt{\gamma}}{2N}, \end{aligned}$$

where

$$\frac{c_1^2\sigma^2}{2N\sqrt{2N}} < \frac{N_1c_2\gamma}{4N\sqrt{2N}}, \text{ holds so long as } N_1 \geq \frac{2c_1^2\sigma^2}{c_2\gamma}, \text{ and} \quad (72)$$

$$\frac{c_2\sqrt{\gamma}c_1\sigma}{2N} \sqrt{\frac{N_2}{N}} < \frac{N_1c_2\gamma}{4N\sqrt{2N}}, \text{ holds so long as } N_1 \geq \frac{8c_1^2\sigma^2\omega_2}{\gamma\omega_1}, \text{ so that} \quad (73)$$

$$N_1 \geq \frac{2\sqrt{2}c_1\sigma\sqrt{N_2}}{\sqrt{\gamma}}, \quad (74)$$

Both conditions are guaranteed by (26) in Theorem 21. ■

This allows us to conclude that

$$\begin{aligned} \left| \frac{N_1x + N_1\delta + N_2y + N_2\tau}{2N} - \frac{N_1x + N_2y}{2N} \right| &= \left| \frac{N_1\delta + N_2\tau}{2N} \right| \\ &\leq \left(\frac{\min\{N_1, N_2\}\sqrt{\gamma}}{2N} \right) |x - y|. \end{aligned}$$

Given that $|y - x| = c_2\sqrt{\gamma}/\sqrt{2N}$ as shown in the Separation Lemma 11, we have

$$\begin{aligned} \left| \frac{N_1x + N_1\delta + N_2y + N_2\tau}{2N} - x \right| &= \left| \frac{N_2(y - x)}{2N} + \frac{N_1\delta + N_2\tau}{2N} \right| \\ &\geq \left| \frac{N_2(y - x)}{2N} \right| - \left| \frac{N_1\delta + N_2\tau}{2N} \right| \\ &\geq \frac{N_2|y - x|}{2N} - \frac{\min\{N_1, N_2\}\sqrt{\gamma}|y - x|}{2N} \\ &\geq \frac{(1 - \sqrt{\gamma})N_2|y - x|}{2N}, \end{aligned}$$

and similarly,

$$\begin{aligned} \left| y - \frac{N_1x + N_1\delta + N_2y + N_2\tau}{2N} \right| &= \left| \frac{N_1(y - x)}{2N} - \frac{N_1\delta + N_2\tau}{2N} \right| \\ &\geq \left| \frac{N_1(y - x)}{2N} \right| - \left| \frac{N_1\delta + N_2\tau}{2N} \right| \\ &\geq \frac{N_1|y - x|}{2N} - \frac{\min\{N_1, N_2\}\sqrt{\gamma}|y - x|}{2N} \\ &\geq \frac{(1 - \sqrt{\gamma})N_1|y - x|}{2N}. \end{aligned}$$

■

D Proofs of Procedure Partition

D.1 Proof of Lemma 15

Suppose that $\frac{1}{2}\rho \leq \rho_j \leq \rho$. To ease up the notation, we omit the superscript j ; thus, we let $S_i = S_i^{(j)}$, $Q_i = Q_i^{(j)}$ for $1 \leq i \leq k$, and $Q(v) = Q^{(j)}(v)$ for $v \in V$ (cf. Steps 2–4 of `Partition`). The following lemma, whose proof we postpone to Section D.2, shows that there is a permutation π such that ξ_i is “close” to $\mathbb{E}^{V_{\pi(i)}}$ for all $1 \leq i \leq k$, and that the sets Q_i are “not too small”. Here we let $\mathbb{E}^{V_t} = (p_t^l)_{l=1, \dots, K}$. In addition, let $\rho = \sqrt{T}$ and $\rho_j = \sqrt{T_j}$.

Lemma 34. *Suppose that $\frac{1}{2}\rho \leq \rho_j \leq \rho$. There is a bijection $\pi : \{1, \dots, k\} \rightarrow \{1, \dots, k\}$ such that for each $1 \leq i \leq k$ we have $|Q_i| \geq \frac{1}{2}|V_{\pi(i)}|$ and $\|\xi_i - \mathbb{E}^{V_{\pi(i)}}\|^2 \leq 0.1\rho^2$.*

In the sequel, we shall assume without loss of generality that the map π from Lemma 34 is just the identity, i.e., $\pi(i) = i$ for all i . Bootstrapping on the estimate $\|\xi_i - \mathbb{E}^{V_i}\|^2 \leq 0.1\rho^2$ for $1 \leq i \leq k$ from Lemma 34, we derive the following stronger estimate.

Corollary 35. *For all $1 \leq i \leq k$ we have $\|\xi_i - \mathbb{E}^{V_i}\|^2 \leq 100|Q_i|^{-1} \sum_{v \in Q_i} \|\widehat{A}_v - \mathbb{E}_v\|^2$.*

Proof. By the Cauchy-Schwarz inequality,

$$\|\xi_i - \mathbb{E}^{V_i}\| = |Q_i|^{-1} \left\| \sum_{v \in Q_i} \widehat{A}_v - \mathbb{E}^{V_i} \right\| \leq |Q_i|^{-1/2} \left[\sum_{v \in Q_i} \|\widehat{A}_v - \mathbb{E}^{V_i}\|^2 \right]^{1/2}. \quad (75)$$

Furthermore, as $\|\xi_i - \mathbb{E}^{V_i}\|^2 \leq 0.1\rho^2$ by Lemma 34, for all $v \in Q_i \setminus V_i$ we have

$$\|\widehat{A}_v - \mathbb{E}^{V_i}\|^2 \leq 2(\|\widehat{A}_v - \xi_i\|^2 + \|\xi_i - \mathbb{E}^{V_i}\|^2) \leq \rho^2/3, \quad (76)$$

because the construction of Q_i in Step 3 of `Partition` ensures that $\|\widehat{A}_v - \xi_i\|^2 \leq 0.01\rho^2$. Hence, as $\|\mathbb{E}_v - \mathbb{E}^{V_i}\|^2 \geq \rho^2$, (76) implies that $\|\widehat{A}_v - \mathbb{E}_v\| \geq 0.1\|\widehat{A}_v - \mathbb{E}^{V_i}\|$. Therefore, the assertion follows from (75). ■

Corollary 36. *For all $v \in S_i \setminus V_i$ we have $\|\widehat{A}_v - \xi_i\| \leq 3\|\widehat{A}_v - \mathbb{E}_v\|$.*

Proof. Let $i \neq l$ and consider a $v \in S_i \cap V_l$. We shall establish below that

$$\|\widehat{A}_v - \xi_i\| \leq \|\widehat{A}_v - \xi_l\|. \quad (77)$$

Then by Lemma 34 $\|\widehat{A}_v - \xi_i\| \leq \|\widehat{A}_v - \mathbb{E}_v\| + \|\mathbb{E}_v - \xi_i\| \leq \|\widehat{A}_v - \mathbb{E}_v\| + \rho/3$, and thus $\rho \leq \|\mathbb{E}_v - \mathbb{E}^{V_i}\| \leq \|\widehat{A}_v - \xi_i\| + \|\xi_i - \mathbb{E}^{V_i}\| + \|\widehat{A}_v - \mathbb{E}_v\| \leq 2\|\widehat{A}_v - \mathbb{E}_v\| + \frac{2}{3}\rho$. Consequently, we obtain $\|\widehat{A}_v - \mathbb{E}_v\| \geq \frac{1}{6}\rho$, so that the assertion follows from the estimate

$$\|\widehat{A}_v - \xi_i\| \stackrel{(77)}{\leq} \|\widehat{A}_v - \xi_l\| \leq \|\widehat{A}_v - \mathbb{E}_v\| + \|\mathbb{E}_v - \xi_l\| \stackrel{\text{Lemma 34}}{\leq} \|\widehat{A}_v - \mathbb{E}_v\| + \frac{\rho}{3} \leq 3\|\widehat{A}_v - \mathbb{E}_v\|.$$

Finally, we prove (77). If $v \in S_i \cap V_l \setminus Q_i$, then the construction of S_i in Step 4 of `Partition` guarantees that $\|\widehat{A}_v - \xi_i\| \leq \|\widehat{A}_v - \xi_l\|$, as claimed. Thus, assume that $v \in Q_i \cap V_l$. Then

$$\begin{aligned} \|\widehat{A}_v - \xi_i\| &\leq 0.15\rho && \text{[by the definition of } Q_i \text{ in Step 3 of } \text{Partition}], \\ \max\{\|\xi_i - \mathbb{E}^{V_i}\|, \|\xi_l - \mathbb{E}_v\|\} &\leq \frac{1}{3}\rho && \text{[by Lemma 34],} \\ \|\mathbb{E}^{V_i} - \mathbb{E}_v\| &\geq \rho && . \end{aligned}$$

Therefore, if $\|\widehat{A}_v - \xi_l\| < \|\widehat{A}_v - \xi_i\|$, then we would arrive at the contradiction

$$\begin{aligned} \rho &\leq \|\mathbb{E}^{V_i} - \mathbb{E}_v\| \leq \|\mathbb{E}^{V_i} - \xi_i\| + \|\mathbb{E}_v - \xi_l\| + \|\xi_i - \xi_l\| \\ &\leq \frac{2}{3}\rho + \|\widehat{A}_v - \xi_i\| + \|\widehat{A}_v - \xi_l\| < \frac{2}{3}\rho + 2\|\widehat{A}_v - \xi_i\| \leq 0.99\rho. \end{aligned}$$

Thus, we conclude that $\|\widehat{A}_v - \xi_l\| \geq \|\widehat{A}_v - \xi_i\|$, thereby completing the proof. ■

Proof of Lemma 15. Since $|Q_i| \geq \frac{1}{2}|V_i|$ by Lemma 34, we have the estimate

$$\begin{aligned} \sum_{i=1}^k \sum_{w \in S_i \cap V_i} \|\widehat{A}_w - \xi_i\|^2 &\leq 2 \sum_{i=1}^k \sum_{w \in S_i \cap V_i} \left[\|\widehat{A}_w - \mathbb{E}_w\|^2 + \|\mathbb{E}_w - \xi_i\|^2 \right] \\ &\stackrel{\text{Cor. 35}}{\leq} 2\|\widehat{A} - \mathbb{E}\|_F^2 + 200 \sum_{i=1}^k \frac{|S_i \cap V_i|}{|Q_i|} \sum_{v \in Q_i} \|\widehat{A}_v - \mathbb{E}_v\|^2 \leq 500\|\widehat{A} - \mathbb{E}\|_F^2. \end{aligned} \quad (78)$$

Furthermore, by Corollary 36

$$\sum_{i=1}^k \sum_{v \in S_i \setminus V_i} \|\widehat{A}_v - \xi_i\|^2 \leq 9 \sum_{i=1}^k \sum_{v \in S_i \setminus V_i} \|\widehat{A}_v - \mathbb{E}_v\|^2 \leq 9\|\widehat{A} - \mathbb{E}\|_F^2. \quad (79)$$

Since $\|\widehat{A} - \mathbb{E}\|_F^2 \leq c_k \lambda^2$ by Lemma 14, the bounds (78) and (79) imply the assertion. \blacksquare

D.2 Proof of Lemma 34

Proof of Lemma 34. For $1 \leq i \leq k$ we choose $\pi(i)$ so that $|Q_i \cap V_{\pi(i)}|$ is maximum. We shall prove below that for all $1 \leq l \leq k$ we have

$$\|\xi_l - \mathbb{E}^{V_{\pi(l)}}\|^2 \leq 0.1\rho^2, \quad (80)$$

$$|Q_l| \geq \max\{|V_i| : i \in \{1, \dots, k\} \setminus \pi(\{1, \dots, l-1\})\} - 0.01n_{\min}, \quad (81)$$

$$|Q_l \cap V_{\pi(l)}| \geq |Q_l| - 0.01n_{\min}. \quad (82)$$

These three inequalities imply the assertion. To see that π is a bijection, let us assume that $\pi(l) = \pi(l')$ for two indices $1 \leq l < l' \leq k$. Indeed, suppose that $l = \min \pi^{-1}(l)$. Then $|Q_l| \geq |V_{\pi(l)}| - 0.01n_{\min}$ by (81), and thus $|V_{\pi(l)} \setminus Q_l| \leq 0.1n_{\min}$ by (82). Therefore, we obtain the contradiction

$$0.99n_{\min} \stackrel{(81)}{\leq} |Q_{l'}| \stackrel{(82)}{\leq} 1.1|Q_{l'} \cap V_{\pi(l)}| \leq 1.1|V_{\pi(l)} \setminus Q_l| \leq 0.11n_{\min}.$$

Finally, as π is bijective, (81) entails that $|Q_l| \geq 0.9|V_{\pi(l)}|$ for all $1 \leq l \leq k$. Hence, due to (82) we obtain $|Q_l \cap V_l| \geq 0.9|Q_l| \geq \frac{1}{2}|V_{\pi(l)}|$, as desired.

The remaining task is to establish (80)–(82). We proceed by induction on l . Thus, let us assume that (80)–(82) hold for all $l < L$; we are to show that then (80)–(82) are true for $l = L$ as well. As a first step, we establish (81). To this end, consider a class V_i such that $i \notin \pi(\{1, \dots, L-1\})$ and let $Z_i = \{v \in V_i : \|\widehat{A}_v - \mathbb{E}_v\|^2 \leq 0.001\rho^2\}$. Then $0.001\rho^2(|V_i| - |Z_i|) \leq \sum_{v \in V_i \setminus Z_i} \|\widehat{A}_v - \mathbb{E}_v\|^2 \leq \|\widehat{A} - \mathbb{E}\|_F^2 \leq c_k \lambda^2$, whence the assumption on $\rho^2 = \Gamma$ yields

$$|Z_i| \geq |V_i| - 0.01n_{\min}. \quad (83)$$

Moreover, for all $v \in Z_i$ we have

$$Q(v) = \{w \in V : \|\widehat{A}_v - \widehat{A}_w\|^2 \leq 0.01\rho^2\} \supset Z_i. \quad (84)$$

In addition, let $w \in Q_l$ for some $l < L$; since our choice of i ensures that $v \in V_i \neq V_{\pi(l)}$, we have

$$\rho \leq \|\mathbb{E}^{V_{\pi(l)}} - \mathbb{E}_v\| \leq \|\mathbb{E}_v - \widehat{A}_v\| + \|\widehat{A}_w - \widehat{A}_v\| + \|\xi_l - \widehat{A}_w\| + \|\xi_l - \mathbb{E}^{V_{\pi(l)}}\| \quad (85)$$

Now, the construction in Step 3 of `Partition` ensures that $\|\widehat{A}_w - \xi_l\| \leq 0.1\rho$. Furthermore, $\|\xi_l - \mathbb{E}^{V_{\pi(l)}}\| \leq \rho/3$ by induction (cf. (80)), and $\|\widehat{A}_w - \mathbb{E}_v\| \leq 0.1\rho$, because $v \in Z_i$. Hence, (85) entails that $\|\widehat{A}_w - \widehat{A}_v\| > 0.1\rho$, so that $w \notin Q(v)$. Consequently, (84) yields

$$Z_i \cap Q_l = \emptyset \text{ for all } l < L. \quad (86)$$

Finally, let v_L signify the element chosen by Step 3 of `Partition` to construct Q_L . Then by construction $|Q_L| = |Q(v_L) \setminus \bigcup_{l=1}^{L-1} Q_l| \geq |Q(v) \setminus \bigcup_{l=1}^{L-1} Q_l|$. Therefore,

$$|Q_L| \geq |Q(v) \setminus \bigcup_{l=1}^{L-1} Q_l| \stackrel{(84), (86)}{\geq} |Z_i| \stackrel{(83)}{\geq} |V_i| - 0.01n_{\min}.$$

As this estimate holds for all $i \notin \pi(\{1, \dots, L-1\})$, (81) follows.

Thus, we know that Q_L is “big”. As a next step, we prove (82), i.e., we show that Q_L “mainly” consists of vertices in $V_{\pi(L)}$. To this end, let $1 \leq i \leq k$ be such that $\|\mathbb{E}^{V_i} - \widehat{A}_{v_L}\|$ is minimum. Let $Y = Q_L \setminus V_i$. Then for all $w \in Y$ we have $\|\mathbb{E}_w - \widehat{A}_{v_L}\| \geq \|\mathbb{E}^{V_i} - \widehat{A}_{v_L}\|$. Further, since $\rho \leq \|\mathbb{E}_w - \mathbb{E}^{V_i}\| \leq \|\mathbb{E}_w - \widehat{A}_{v_L}\| + \|\mathbb{E}^{V_i} - \widehat{A}_{v_L}\| \leq 2\|\mathbb{E}_w - \widehat{A}_{v_L}\|$, we conclude that $\|\mathbb{E}_w - \widehat{A}_{v_L}\|^2 \geq \frac{1}{4}\rho^2$. On the other hand, as $w \in Q_L$, we have $\|\widehat{A}_w - \widehat{A}_{v_L}\|^2 \leq 0.01\rho^2$. Therefore, we obtain $\|\widehat{A}_w - \mathbb{E}_w\|^2 \geq 0.1\rho^2$ for all $w \in Y$, so that

$$0.1|Y|\rho^2 \leq \sum_{w \in Y} \|\widehat{A}_w - \mathbb{E}_w\|^2 \leq \|\widehat{A} - \mathbb{E}\|_F^2 \stackrel{\text{Lemma 14}}{\leq} c_k \lambda^2. \quad (87)$$

Hence, due to our assumption on $\rho^2 = \Gamma$, (87) yields that $|Y| < 0.01n_{\min}$. Consequently, (81) entails that $|V_i \cap Q_L| \geq 0.99|Q_L|$, so that $i = \pi(L)$. Hence, we obtain $|Q_L \cap V_{\pi(L)}| = |Q_L \cap V_i| = |Q_L \setminus Y| \geq |Q_L| - 0.01n_{\min}$, thereby establishing (82).

Finally, to show (80), we note that by construction $\|\xi_L - \widehat{A}_{v_L}\|^2 \leq 0.01\rho^2$ and $\|\widehat{A}_w - \widehat{A}_{v_L}\|^2 \leq 0.01\rho^2$ for all $w \in Q_L \cap V_{\pi(L)}$ (cf. Step 3 of `Partition`). Therefore,

$$\begin{aligned} & |Q_L \cap V_{\pi(L)}| \cdot \|\mathbb{E}_{\pi(L)} - \xi_L\|^2 \\ & \leq 3 \sum_{w \in Q_L \cap V_{\pi(L)}} \|\xi_L - \widehat{A}_{v_L}\|^2 + \|\widehat{A}_w - \widehat{A}_{v_L}\|^2 + \|\widehat{A}_w - \mathbb{E}_{\pi(L)}\|^2 \\ & \leq 0.06\rho^2 |Q_L \cap V_{\pi(L)}| + 3\|\widehat{A} - \mathbb{E}\|_F^2 \\ & \stackrel{\text{Lemma 14}}{\leq} 0.06\rho^2 |Q_L \cap V_{\pi(L)}| + 3c_k \lambda^2. \end{aligned} \quad (88)$$

Since $|Q_L \cap V_{\pi(L)}| \geq 0.9n_{\min}$ due to (81) and (82), (88) entails that $\|\mathbb{E}_{\pi(L)} - \xi_L\|^2 \leq 0.07\rho^2 + \frac{6c_k \lambda^2}{n_{\min}} \leq 0.1\rho^2$. Thus, (80) follows. \blacksquare

D.3 Proof of Lemma 16

Proof of Lemma 16. Set $S_{ab} = S_a \cap V_b$ for $1 \leq a, b \leq k$. Moreover, for each $1 \leq a \leq k$ let $1 \leq \pi(a) \leq k$ be such that $\|\mathbb{E}^{V_{\pi(a)}} - \xi_a\|$ is minimum. Then for all $b \neq \pi(a)$ we have

$$\rho \leq \|\mathbb{E}^{V_{\pi(a)}} - \mathbb{E}^{V_b}\| \leq \|\mathbb{E}^{V_{\pi(a)}} - \xi_a\| + \|\mathbb{E}^{V_b} - \xi_a\| \leq 2\|\mathbb{E}^{V_b} - \xi_a\|, \quad (89)$$

so that $\|\mathbb{E}^{V_b} - \xi_a\| \geq \rho/2$. Therefore, by our assumption that $\sum_{i=1}^k \sum_{v \in S_i} \|\xi_i - \widehat{A}_v\|^2 \leq C_0 k^3 \lambda^2$, we have

$$\begin{aligned} \frac{\rho^2}{4} \sum_{a=1}^k \sum_{1 \leq b \leq k: b \neq \pi(a)} |S_{ab}| &\leq \sum_{a,b=1}^k |S_{ab}| \cdot \|\mathbb{E}^{V_b} - \xi_a\|^2 \\ &\leq 2 \sum_{a,b=1}^k \sum_{v \in S_{ab}} \|\mathbb{E}_v - \widehat{A}_v\|^2 + \|\widehat{A}_v - \xi_a\|^2 \\ &\leq 2\|\widehat{A} - \mathbb{E}\|_F^2 + 2 \sum_{a,b=1}^k \sum_{v \in S_{ab}} \|\widehat{A}_v - \xi_a\|^2 \\ &\stackrel{\text{Lemma 14}}{\leq} 4C_0 k^3 \lambda^2 + 2C_0 k^3 \lambda^2 \leq C_0^2 k^3 \lambda^2. \end{aligned} \quad (90)$$

Hence,

$$\sum_{a=1}^k |S_a \triangle V_{\pi(a)}| = \sum_{1 \leq a, b \leq k: b \neq \pi(a)} 2|S_{ab}| \leq \frac{8C_0^2 k^3 \lambda^2}{\rho^2} \leq 0.001 n_{\min}. \quad (91)$$

Combining (90) and (91), we obtain $\frac{n_{\min}}{2} \|\mathbb{E}^{V_{\pi(a)}} - \xi_a\|^2 \leq |S_a \cap V_{\pi(a)}| \cdot \|\mathbb{E}^{V_{\pi(a)}} - \xi_a\|^2 \leq C_0^2 k^3 \lambda^2$, whence

$$\|\mathbb{E}^{V_{\pi(a)}} - \xi_a\|^2 \leq \frac{2C_0^2 k^3 \lambda^2}{n_{\min}} \leq 0.001 \rho^2 \quad \text{for all } 1 \leq a \leq k. \quad (92)$$

Thus, we have established the first two parts of the lemma. In addition, observe that (91) implies that π is bijective (because the sets S_1, \dots, S_k are pairwise disjoint and $|V_a| \geq n_{\min}$ for all $1 \leq a \leq k$). Finally, the third assertion follows from the estimate

$$\begin{aligned} \sum_{a,b=1}^k |S_{ab}| \cdot \|\mathbb{E}^{V_{\pi(a)}} - \mathbb{E}^{V_{\pi(b)}}\|^2 &\leq 2 \sum_{a,b=1}^k |S_{ab}| (\|\mathbb{E}^{V_{\pi(a)}} - \xi_a\|^2 + \|\mathbb{E}^{V_{\pi(b)}} - \xi_a\|^2) \\ &\stackrel{(89)}{\leq} 8 \sum_{a,b=1}^k |S_{ab}| \cdot \|\mathbb{E}^{V_{\pi(b)}} - \xi_a\|^2 \\ &\stackrel{(90)}{\leq} 8C_0^2 k^3 \lambda^2 \leq 0.001 \rho^2 n_{\min}. \end{aligned}$$

■