

1994

Universal bridge free graphs

Martin Goldstern
Carnegie Mellon University

Menachem Kojman

Follow this and additional works at: <http://repository.cmu.edu/math>

This Technical Report is brought to you for free and open access by the Mellon College of Science at Research Showcase @ CMU. It has been accepted for inclusion in Department of Mathematical Sciences by an authorized administrator of Research Showcase @ CMU. For more information, please contact research-showcase@andrew.cmu.edu.

NOTICE WARNING CONCERNING COPYRIGHT RESTRICTIONS:
The copyright law of the United States (title 17, U.S. Code) governs the making of photocopies or other reproductions of copyrighted material. Any copying of this document without permission of its author may be prohibited by law.

UNIVERSAL BRIDGE FREE GRAPHS

by

Martin Goldstern
Institute of Algebra
Technical University
Vienna, Austria

and

Menachem Kojman
Department of Mathematics
Carnegie Mellon University
Pittsburgh, PA 15213, USA

Research Report No. 94-171₂
August, 1994

510.6
C28R
94-171

University Libraries
Carnegie Mellon University
Pittsburgh PA 15213-3890

UNIVERSAL BRIDGE FREE GRAPHS

July 1994

Martin Goldstern

Institute of Algebra

Technical University

Vienna, Austria

`goldstrn@email.tuwien.ac.at`

Menachem Kojman

Department of Mathematics

Carnegie Mellon University

Pittsburgh, PA 15213, USA

`kojman@andrew.cmu.edu`

ABSTRACT. We prove that there is no countable universal B_n -free graph for all n and that there is no countable universal graph in the class of graphs omitting all cycles of length at most $2k$ for $k \geq 2$.

University Libraries
Carnegie Mellon University
Pittsburgh, PA 15213-0350

§0 Introduction

Several papers have addressed the problem of existence of a universal element among all countable graphs omitting given finite subgraphs (see [KP] and the comprehensive bibliography there, and most recently [CK] and [KP1]).

Given a graph F we say that a graph G is F -free if F is not isomorphic to a subgraph of G . A countable F -free graph G^* is universal (strongly universal) in the class of all countable F -free graphs if every countable F -free graph is isomorphic to a subgraph (an induced subgraph) of G^* .

In [CK], Cherlin and Komjath raise the problem of determining for which finite trees T there exists a universal countable T -free graphs. In this paper we describe an infinite set of finite trees B_n , which we call bridges, and show that for no n is there a universal countable B_n -free graph.

In [CK] it is proved that for all $n \geq 4$ there is no universal countable C_n -free graph (where C_n is a cycle of length n). In [KMP] it is proved, on the other hand, that a strongly universal countable graph exists among all countable graphs that omit all odd cycles of length at most $2k + 1$. What if we intersect some of those classes, say look at all graphs omitting C_3, C_4, C_5, C_6 ? We show here, using an idea of S. Mozes, that when all cycles of length at most $2k$ are to be excluded (for $k \geq 2$), then there is no universal countable graph.

0.1 Problem: Is there a countable universal graph in the class of graphs omitting all cycles of length at most $2k + 1$ (for $k \geq 2$)? More generally, for what sets $F \subseteq \mathbb{N}$ does the class of graphs omitting $\{C_n : n \in F\}$ have a countable universal element?

Following [KP] we make the following definition:

0.2 Definition: Let \mathcal{G} be a class of graphs.


- (i) The complexity $\text{cp}(\mathcal{G})$ of \mathcal{G} is the minimal cardinality κ of a set I of graphs in \mathcal{G} with the property that every member in \mathcal{G} is embedded as an induced subgraph into at least one of the members of I .
- (ii) The weak complexity $\text{wcp}(\mathcal{G})$ is defined by omitting the word “induced” from the definition of $\text{cp}(\mathcal{G})$.

Notation: We denote the vertex degree of v in a graph G by $\deg_G(v)$. The *length* of a path is the number of edges in the path. For all m let K_m denote a complete graph with m vertices k_1, \dots, k_m , let P_m be a simple path of length m with vertices p_0, \dots, p_m and edges (p_i, p_{i+1}) for $i \leq m$ and let C_m denote a cycle of length m . Let us call a simple path h_0, h_1, \dots, h_k in a graph G a *highway* iff $\deg_G(v_i) = 2$ for all $0 < i < k$.


0.3 Advice: Drive carefully.

§1 Bridge-free graphs

1.1 Definition: A finite graph with $n+5$ vertices is called an n -bridge iff it is isomorphic to $B_n = \langle V, E \rangle$ where $V = \{a, b, c, x_1, x_2, \dots, x_n, d, e\}$ and $E = \{(a, c), (b, c), (x_n, d), (x_n, e)\} \cup \{(x_i, x_{i+1}) : 1 \leq i < n\}$



1.2 Definition: Let us call a graph D a *dead end* if it is isomorphic to K_{n+3} to which a simple path P_{n+1} is freely adjoined by identifying k_{n+3} with p_0 .



1.3 Claim: A dead end is B_n -free and if a dead end D with vertices $k_1, \dots, k_{n+3} = p_0, x_1, \dots, p_{n+1}$ is a subgraph of a B_n -free graph G then $\deg_D(v) = \deg_G(v)$ for all vertices $v \in D$ except maybe $v = p_{n+1}$.

Proof: : Suppose first that for some $i \leq n$ there is an edge (p_i, y) in G which is not an edge of D . If $y \notin D$, by labeling y as a , labeling p_i as c and p_{i+1} as b it is possible to label vertices of D as x_i ($i \leq n$) and as d, e to produce a copy of B_n .


Suppose, then, that (p_i, p_j) is an edge not among the edges of D . Without loss of generality, $j \geq i + 2$. Now label p_j as a , label P_i as c and p_{i+1} as b . Again, a copy of B_n is easily found.

The remaining possibility for an edge of the form (p_i, y) is that $y = k_j$ for some $j \leq n+3$. Here we distinguish two subcases. First, $j = 1$ (and, of course, $i > 1$). Labeling p_1 as a , k_1 as c and p_1 as b , the remaining $n+2$ vertices of K_{n+3} complete the three labeled ones to make a copy of B_n .

Second, $j \neq 1$. In this case if $i > 1$ label k_j as c , label k_1 as x_n and label p_1 as e . The remaining vertices of K_{n+3} serve as x_i for $1 \leq i < n$ and as b . If, however, $i = 1$, label p_1 as c , label p_2 as a and label k_j as b . Again, a copy of B_n is found.

We show next that the $\deg_D(k_i)$ is preserved. Suppose that (k_i, y) is an edge in G which is not an edge in D . We already proved that $y \neq p_j$ for all $j \leq n$. Therefore, either $y \notin D$ or $y = p_{n+1}$. If $i = 1$ label p_1 as a , label y as b and k_1 as c ; otherwise label k_i as c , y as a , p_1 as e and k_1 as x_n . In both cases a copy of B_n results ☺ 1.3

1.4 Definition: Let us call a graph T a *drive through* if it is isomorphic to the graph obtained as follows: Let k_1, \dots, k_{n+2} be the vertices of a copy of K_{n+2} . For $1 < i < n+2$ adjoin freely to k_i a copy of a dead end by identifying p_{n+1} in that copy with k_i . To k_1 connect a vertex l by an edge and to k_{n+2} connect a vertex r by an edge. Call l the *left exit* of T and call r the *right exit* of T .



1.5 Claim: A drive through T is B_n -free and if T is a subgraph of a B_n -free graph G then $\deg_T(v) = \deg_G(v)$ for all vertices $v \in T$ except l and r .

Proof: Suppose to the contrary that B_n is a subgraph of a drive through T . As $\deg_{B_n}(c) = \deg_{B_n}(x_n) = 3$, both a and x_n are either in a copy of K_{n+3} or in the copy of K_{n+2} . Both cannot be in the same copy of K_{n+3} because the minimum of distances of x_i to c and x_n is smaller than $n+1$, and all the points satisfying this would be in the same dead end as x_n and c , contrary to claim 1.3. Similarly, c and x_n are not both in the copy of K_{n+2} .

Also, c and x_n cannot be in different copies of K_{n+3} , or in a copy of K_{n+3} and in the copy of K_{n+2} because the distance between x_n and c would be greater than n . We conclude that T is B_n -free.


Suppose that T is a subgraph of a B_n free graph G . By claim 1.3 we know that $\deg_T(v) = \deg_G(v)$ for all vertices v in the dead ends except those which are also in the copy of K_{n+2} . Suppose that for some vertex k_i in the copy of K_{n+2} there is an edge (k_i, y) in G which is not an edge of T . Label y as a . If $i = 1$ or $i = n + 2$ label l or r respectively as b . Label k_i as c . Label n of the remaining k_i as x_1, \dots, x_n . Label the last remaining k_i as d . If this i is 1 or $n + 1$ label l or r respectively as e . Otherwise label as e the vertex p_n in the dead end adjoined to k_i . This yields a copy of B_n . ☺ 1.5

For every $\epsilon \in \omega^2$ we construct a connected B_n -free graph G_ϵ as follows.

Let $T^\epsilon(m)$ for $m \in \mathbb{N}$ and $\epsilon \in \omega^2$ be disjoint copies of a drive through. Let $l^\epsilon(m)$ and $r^\epsilon(m)$ be the left and right exits of $T^\epsilon(m)$. Let D^ϵ be a copy of D with vertices $k_1^\epsilon, \dots, k_{n+3}^\epsilon = p_0^\epsilon, \dots, p_{n+1}^\epsilon$. Let $H^\epsilon(m)$ be a simple path of length $n - 1 + \epsilon(m)$ with vertices $h_0^\epsilon(m), \dots, h_{n-1+\epsilon(m)}^\epsilon(m)$.

Adjoin D^ϵ to $l^\epsilon(0)$ by setting $p_{n+1}^\epsilon = l^\epsilon(0)$. Connect $r^\epsilon(m)$ to $l^\epsilon(m + 1)$ by $H^\epsilon(m)$ by setting $r^\epsilon(m) = h^\epsilon(0)$ and $l^\epsilon(m + 1) = h_{n-1+\epsilon(m)}^\epsilon(m)$. (If $n = 1$ then when $\epsilon(m) = 0$ we identify $r^\epsilon(m)$ with $l^\epsilon(m + 1)$.)

Let $G_\epsilon = D^\epsilon \cup \bigcup T^\epsilon(m) \cup \bigcup H^\epsilon(m)$.



Let us observe that all highways in G_ϵ are either of length $n + 1$ or of length $n + 2$. All highways that have an end of degree $n + 3$ are of length $n + 1$ except a unique highway — the one containing $l^\epsilon(0)$ — which is of length $n + 2$. Let us denote this highway by $H(\epsilon)$.

1.6 Claim: The graph G_ϵ is B_n -free and if G_ϵ is a subgraph of a B_n -free graph G then the vertex degree of every vertex $v \in G_\epsilon$ in G_ϵ equals the degree of v in G .

Proof: : A similar argument to that in 1.5 shows that G_ϵ is B_n -free. Suppose now that $G_\epsilon \subseteq G$ and that G is B_n -free. By 1.5 we already know for $\deg_{G_\epsilon}(v) = \deg_G(v)$ for each

vertex $v \in T^\epsilon(m)$ except $l^\epsilon(m), r^\epsilon(m)$. If, however, $\deg_G(v) > \deg_{G_\epsilon}(v)$ when v is on one of the highways of G_ϵ , there must be some $y \in G \setminus G_\epsilon$ such that (v, y) is an edge of G and a copy of B_n is easily produced. ☺ 1.6

1.7 Corollary: For every $\epsilon \in \omega^2$ and every connected B_n -free graph G , if $G_\epsilon \subseteq G$ then $G_\epsilon = G$.

Proof: Suppose that $y \in G \setminus G_\epsilon$. By connectedness of G we may assume that y is connected by an edge to a vertex of G_ϵ . This contradicts 1.6

1.8 Claim: If $\epsilon \neq \nu$ are two members of ω^2 then G_ϵ and G_ν are not isomorphic.

Proof: Suppose that $f : G_\epsilon \rightarrow G_\nu$ is an isomorphism. We show that $\epsilon = \nu$. Clearly, f maps every highway in G_ϵ onto some highway in G_ν .

The highway $H(\epsilon)$ has to be mapped by f onto $H(\nu)$, both being the unique highways in their respective graphs of length $n+2$ with an end of degree $n+3$. As $l^\epsilon(0)$ is connected by an edge to the end of $H(\epsilon)$ that has degree $n+2$, we conclude that $f(l^\epsilon(0)) = l^\nu(0)$. We argue by induction on m that $H^\epsilon(m)$ is mapped by f onto the $H^\nu(m)$ and that $f(l^\epsilon(m+1)) = l^\nu(m+1)$.

If $m = 0$, we already showed that $f(l^\epsilon(0)) = l^\nu(0)$. Therefore $f(r^\epsilon(0)) \neq l^\nu(0)$. Also, $f(r^\epsilon(0))$ cannot lie on any of the highways in $T^\nu(0)$ which are part of a dead end, because both ends of $H^\epsilon(0)$ have degree $n+2$. Therefore necessarily $f(r^\epsilon(0)) = r^\nu(0)$ and consequently $H^\epsilon(0)$ is mapped by f onto the $H^\nu(0)$, with $f(l^\epsilon(1)) = l^\nu(1)$.

Similarly, if f maps $H^\epsilon(m)$ onto $h^\nu(m)$ with $f(l^\epsilon(m+1)) = l^\nu(m+1)$, it follows that f maps $H^\epsilon(m+1)$ onto $h^\nu(m+1)$ with $f(l^\epsilon(m+2)) = l^\nu(m+2)$.

As for all m we have established that $n-1 + \epsilon(m) = n-1 + \nu(m)$, we have shown that $\epsilon = \nu$. ☺ 1.8

1.9 Theorem: There is no universal B_n -free graph. In fact, the weak complexity of the class of countable B_n -free graphs equals 2^{\aleph_0} .

Proof: Suppose that $\{G_\alpha : \alpha \in I\}$ is a collection of less than 2^{\aleph_0} many countable B_n free graphs. By splitting each graph to its connected components we assume that each G_α is connected. Suppose that for every $\epsilon \in \omega^2$ the graph G_ϵ constructed above is isomorphic

to a subgraph of G_α for some $\alpha \in I$. By corollary 1.7 and the assumption just made, each G_ϵ is isomorphic to G_α for some $\alpha \in I$. By the pigeon hole principle there is a single G_α which is isomorphic to uncountably many G_ϵ . This contradicts claim 1.8 ☺ 1.9

§2 Graphs without short cycles

In this section we show that the class of all graphs omitting all cycles of length at most $2k$ ($k \geq 2$) has no countable universal element.

2.1 Definition: Let S_k be the following graph: For five vertices $\{x_i : i \in Z_5\}$ indexed cyclically connect x_i to x_{i+1} by a simple path $x_i, y_{i,1}, \dots, y_{i,k-1}, x_{i+1}$.

2.2 Claim: If f_1 and f_2 are two embeddings of S_k into a graph G omitting all cycles of length at most $2k$ ($k \geq 2$) and $f_1(x_i) = f_2(x_i)$ for $i \in Z_5$ then $f_1 = f_2$.

Proof: : Suppose for simplicity that f_1 is the inclusion, and suppose that $f_2 \neq f_1$. Let i be the least such that among $\{y_{i,j} : j < k\}$ there is a vertex v for which $v \neq f_2(v)$ and let $j(0)$ be the least such that $y_{i,j} \neq f_2(y_{i,j(0)})$. Let $j(1)$ be the the least $j > j(0)$ such that $x_{i,j(1)} = f_2(x_{i,j(1)})$. Now $x_{i,j(0)-1}, x_{i,j(0)}, \dots, x_{i,j(1)}, f_2(x_{i,j(0)}), \dots, f_2(x_{i,j(1)-1})$ forms a cycle of length $\leq 2k$ in G , contrary to the assumption. ☺

Let us define an infinite graph U by induction. For every natural m let $S(m)$ be a copy of S_k with vertices $x_i^m, y_{i,j}^m$ ($i \in Z_5, 1 \leq j < k$).

Let $U(0) = S(0)$. Suppose that $U(m)$ is defined and $S(m) \subseteq U(m)$. To obtain $U(m+1)$ adjoin freely $S(m+1)$ of to $S(m)$ by identifying x_i^{m+1} with with $y_{2i,[(k+1)/2]}^m$. Let $U = \bigcup U(m)$

2.3 Claim: (i) The graph U contains no cycles of length $\leq 2k$, and $\deg_U(v) \leq 3$ for all $v \in U$. (ii) If f_1 and f_2 are two embeddings of U into a graph G which contains no cycles of length at most $2k$ and $f_1(x_i^0) = f_2(x_i^0)$ for $i \in Z_5$ then $f_1 = f_2$.

Proof: : (i) is clear. Suppose that f_1, f_2 are as stated. Using claim 2.2 inductively one sees that $f_1 = f_2$ ☺ 2.3

Let us choose, by induction on m , vertices v_m in U such that the distance in U between v_m and v_{m+1} is at least $2k+1$. For every $\epsilon \in \omega^2$ let us construct a graph U_ϵ as follows:

let $u(m)$ be distinct vertices not in U . Connect $v(m)$ to $v(m+1)$ by an edge if $\epsilon(m) = 1$ and connect $u(m)$ by edges to $v(m), v(m+1)$ otherwise. Let $U_\epsilon = U \cup \{u(m) : m \in M\}$.

It is not hard to verify that each U_ϵ omits all cycles of length at most $2k$.

2.4 Theorem: For all $k \geq 2$ there is no universal countable graph in the class of all graphs omitting all cycles of length at most $2k$. In fact, the weak complexity of the class of all such countable graphs is 2^{\aleph_0} .

Proof: Suppose to the contrary that $\{G_\alpha : \alpha \in I\}$ is a set of countable graphs, each omitting all cycles of length at most $2k$, with the property that every countable graph omitting all cycles of length at most $2k$ is isomorphic to a subgraph of G_α for at least one $\alpha \in I$, and assume that $|I| < 2^{\aleph_0}$. Fix an embedding f_ϵ of U_ϵ to some $G_{\alpha(\epsilon)}$. By the pigeon hole principle there is a single $\alpha \in I$ which equals $\alpha(\epsilon)$ for all $\epsilon \in A$ for some uncountable set $A \subseteq {}^\omega 2$. For each $\epsilon \in A$ let $\eta(\epsilon) = \langle f_\epsilon(x_i^0) : i \in \mathbb{Z}_5 \rangle$. As there are only countably many finite sequences of vertices in G_α , there are different $\epsilon, \nu \in A$ with $\eta(\epsilon) = \eta(\nu)$. But then it follows by claim 2.3 that $f_\epsilon \upharpoonright U = f_\nu \upharpoonright U$. Let m be such that $\epsilon(m) \neq \nu(m)$. The vertices $f_\epsilon(v(m)), f_\epsilon(u(m)), f_\nu(v(m+1))$ span a copy of C_3 in G_α , contrary to the assumption that G_α contains no cycles of length at most $2k$. ☺ 2.4

References

- [CK] G. Cherlin and P. Komjáth, *There is No Universal Countable Pentagon-Free graph*, Journal of Graph Theory 18 (4) (1994) 337–341
- [KP] Peter Komjáth and János Pach, *Universal elements and the complexity of certain classes of infinite graphs*, Discrete Mathematics 95 (1991) 255–270
- [KP1] P. Komjáth and J. Pach, *The complexity of a class of infinite graphs*, Cominatorica 14(1) (1994) 121–125
- [KMP] P. Komjáth, A. Mekler and J. Pach, *Some Universal Graphs*, Israel Journal of Mathematics 64 (1988) 158–168