Carnegie Mellon University
Research Showcase @ CMU

Department of Mathematical Sciences | Mellon College of Science

1991

A minimization problem involving variation of the domain

Dacorogna
Carnegie Mellon University

Irene Fonseca

Follow this and additional works at: http://repository.cmu.edu/math

Recommended Citation
.

This Technical Report is brought to you for free and open access by the Mellon College of Science at Research Showcase @ CMU. It has been accepted for inclusion in Department of Mathematical Sciences by an authorized administrator of Research Showcase @ CMU. For more information, please contact research-showcase@andrew.cmu.edu.

NOTICE WARNING CONCERNING COPYRIGHT RESTRICTIONS:
The copyright law of the United States (title 17, U.S. Code) governs the making of photocopies or other reproductions of copyrighted material. Any copying of this document without permission of its author may be prohibited by law.

NAMS
91-8

A MINIMIZATION PROBLEM INVOLVING VARIATION OF THE DOMAIN

by

Bernard Dacorogna
Department of Mathematics
Ecole Polytechnique Federale de Lausanne
1015 Lausanne, Switzerland

and

Irene Fonseca
Department of Mathematics
Carnegie Mellon University
Pittsburgh, PA 15213

Research Report No. 91-99-NAMS-8

January 1991

University Libraries
Carnegie Mellon University
Pittsburgh, PA 15213-3890

A MINIMIZATION PROBLEM INVOLVING VARIATION OF THE DOMAIN

BERNARD DACOROGNA
Department of Mathematics
Ecole Polytechnique Federate de Lausanne
1015 Lausanne, Switzerland

and

IRENE FONSECA
Department of Mathematics, Carnegie Mellon University
Pittsburgh, Pennsylvania 15213, U. S. A.

Table of Contents :

1. Introduction 1
2. The case W(X) = IIXII^r 4
3. Existence and regularity results 10
4. Non existence results 19
5. Qualitative properties 22
Acknowledgment 27
References 27

1. INTRODUCTION.

DAVINI [4] and DAVINI & **PARRY** [5, 6] introduced a model for slightly defective crystals where non-elastic defect-preserving deformations are called *neutral* and generally they involve some kind of rearrangement representing the slip mechanisms of the classic phenomenological plasticity theories. Neutral deformations can be factorized into components which are exclusively elastic at the macroscopic level or exclusively slip at the microscopic level. Essentially, a neutral change of state of a perfect crystal corresponds to a lattice matrix

L(u(x)) = Vu(x){Vv(x)}-i,

where u is the elastic deformation of the reference configuration *Q Q* [R^3 into u(Q) and v represents the slip or plastic deformation with det Vv = 1 a. e. in *Cl.*

Taking the viewpoint that equilibria correspond to some kind of variational principle, in DAVINI & PARRY [4, 5] and in FONSECA & PARRY [10] the implications of including in the class of admissible variations the neutral changes of state were analyzed. Precisely, FONSECA & PARRY [10] considered the minimization of the total stored energy functional

$$E(u, v) := \int_{\Omega} fw(Vu(x)\{Vv(x)\}^{-1})\, dx \qquad (1.1)$$

where W represents the strain energy density in the class of admissible pairs

& := {(u, v) G W ^ ^ D R ^ W ^{1} * - ^ ^ ^{3}) ! u = uo on 3Q, det Vu > 0 and det Vv = 1 a. e. in Q}.

Of course, *S*& includes the elastic deformations in the case where v is the identity map. Formally, minimizing E(.,.) in *s*& involves variations of the reference domain ; indeed, setting co := u (v^1) the integral (1.1) becomes

$$\int_{(\Omega)} W(Vco(y))dy.$$

As it is well known, the bulk energy W for ordered materials is not quasiconvex (see ERICKSEN [7], , FONSECA [8], KINDERLEHRER [11]) and so, the functional E(.,.) is not lower semicontinuous. Hence, we cannot use the direct methods of the calculus of variations in order to obtain existence of minimizers of the energy and in general, such minimizers exist only in the generalized sense. Using the parametrized probability measures of YOUNG and the theory of

University Libraries
Carnegie Mellon University
Pittsburgh, PA 15213-3890

compensated compactness of MURAT & TARTAR (see TARTAR [12]), FONSECA & PARRY [10] examined the behavior of minimizing sequences for defective crystals and their state functions.

In this paper we study the existence and regularity properties for minimizers of (1.1) where W satisfies some convexity assumption. It should be pointed out immediatly that the direct methods of the calculus of variations fail to apply to this problem. Indeed, sequential weak lower semicontinuous of E(.,.) (see Propositions 3.8 and 3.10) is not sufficient to ensure existence of minimizers. Precisely, setting

$$W(X) = \|X\|^r$$

where $\|X\|^2 := \sum_{i,j=1}^{N} X_{ij}^2$, we shall establish that minimizers exist if and only if $r \geq N$ (See Theorem 2.2 and Proposition 4.1). This is in sharp contrast with the usual Dirichlet problem of minimizing

$$\inf\{ \int_\Omega \|\nabla u(x)\|^r \, dx \mid u = u_0 \text{ on } \partial\Omega,\ u \in W^{1,r}(\Omega)\}$$

which has solutions for every $r > 1$. Surprisingly the problem behaves in fact very similarly to

$$\text{(Q)} \qquad \inf \{ \int_\Omega |\det \nabla u(x)|^{r/N} \, dx \mid u = u_0 \text{ on } \partial\Omega\}$$

(cf. Corollary 2.5 and Proposition 4.3). This is in agreement with the continuum theory for elastic crystals where it can be shown that, due to the crystallographic material symmetries, the relaxation of the bulk energy depends only on the determinant of the deformation gradient (see ERICKSEN [7], FONSECA [8], KINDERLEHRER [11]).

Another interesting feature of this problem is that, under some convexity-type hypotheses on W satisfied by $W(X) = \|X\|^r$, $r \geq N$, there are solutions (u^*, v^*) verifying

$$\nabla u^*(x)\{\nabla v^*(x)\}^{-1} = X_0 \quad \text{for every } x \in \Omega,$$

where X_0 is a constant matrix. In the case where $W(X) = \|X\|^r$, $r \geq N$, it turns out that $X_0 = \lambda R$ where R is an orthogonal transformation and $\lambda^N = \text{meas } u_0(\Omega) / \text{meas } \Omega$.

2. THE CASE $W(X) = ||X||^r$.

Although the results obtained in this section are srtictly included on the next, we present them beforehand for the sake of clarity. We start by introducing some notations.

<u>Notations</u> : i) $M^{N\times N}$ denotes the set of NxN matrices and if $X \in M^{N\times N}$ then adj X denotes the matrix of cofactors. In particular, if A is invertible then

$$A^{-1} = \frac{(\mathrm{adj}X)T}{\det X}$$

and (2.1)

$$\langle X, \mathrm{adj}\, X\rangle = N \det X$$

where

$$\langle X, Y\rangle := \sum_{i,j=1}^{N} X_{ij}\, Y_{ij}.$$

ii) Let $Q \subset \mathbb{R}^N$ be a bounded, open set with strongly Lipschitz boundary 3Q. If $k \geq 1$ is an integer and if $0 \leq a \leq 1$ then by $\mathrm{Diff}^{k,a}(\overline{\Omega})$ we mean the set of diffeomorphisms $u : \overline{\Omega} \to \overline{u(Q)}$ such that u, $u^{-1} \in C^{k,a}(\overline{Q}, \overline{u(Q)})$, where $C^{k,a}$ stands for the usual set of Hölder continuous functions. In the case k = ~ we shall write $\mathrm{Diff}^{\infty}(\overline{Q})$.

iii) With the above notations, if $u_0 \in \mathrm{Diff}^{k,a}(\overline{Q})$ with det $Vu_0 > 0$ in $\overline{Q}$ is given we let

$$\mathcal{A}_{k,a} := \{(u, v) \in \mathrm{Diff}^{k,a}(\overline{Q}) \times \mathrm{Diff}^{k,a}(\overline{Q}) \mid u = u_0 \text{ on } \partial Q \text{ and } \det Vv(x) = 1 \text{ in } \Omega\}$$

and in the case k = «> we write simply $\mathcal{A}_{\infty}$. Finally, for $r \geq 1$ consider the problem

(P) $\inf \{E(u, v) := \int_{\Omega} \|Vu(x)(Vv(x))^{-1}\|^r dx \mid (u, v) \in \mathcal{A}_{k,a}\}$.

Remark 2.1. We note that, formally, problem (P) reduces to the minimization of functional where both the domain and the deformation are varying. Indeed, if v was invertible then

$$E(u, v) = \int_{v(\Omega)} \|Vu(v^{-1}(y))Vv^{-1}(y)\|^r dy$$

$$= \int_{\Omega} \|Vu^*(y)\|^r dx,$$

where $u^*(y) := u(v^{-1}(y))$.

Theorem 2.2

Let k ≥ 1 be an integer, 0 < a < 1, let *Cl Q* [R^N be a bounded, open set with $C^{k+3,a}$ boundary and let uo e Diff^$^a(\overline{Q})$, det Vuo > 0 in $\overline{Q}$. Then (P) attains its minimum at every (u*, v*) € *S#±$_a$* such that

Vu*(x)(Vv*(x))-^{1} = *X* R in Q

where X^N = meas uo(n)/meas *Q* and R is an orthogonal transformation. Thus

inf {E(u, v) I (u, v) G *sa$_{Ka}$*} =E(u*, v*) = N^ meas (Q)i*/N (meas $u_o(Q))^{r/N}$.

The proof of Theorem 2.2 is based on the following lemma.

Lemma 2,3

If A G $M^{N\times N}$ then IIAII^N ≥ NW Idet Al. Furthermore, the equality holds if and only if A = A,R, for some *X e* IR and some orthogonal transformation R.

Proof. If det A = 0, then the inequality is trivially valid. Suppose that det A > 0. Using the polar decomposition, we can write

A = RU,

where U = U^7 > 0 and R is an orthogonal transformation, i. e. R^T R = R R^T = 11, det R = 1. Thus, U = Q^T diag(?ii,..., XN) Q, where Q is an orthogonal transformation and A,i,..., ^N > 0 and so

$$HAH = IIRUII = HUH = (\sum_{i=1}^{N} X\hat{\ })^{1/2}. \tag{2.2}$$

As In is a concave function, we have

$$\ln(\det A) = \ln\left[\prod_{i=1}^{N} \lambda_i\right] = \frac{1}{2}\sum_{i=1}^{N} \ln(\lambda_i^2) \le \frac{N}{2}\ln\left(\frac{1}{N}\sum_{i=1}^{N}\lambda_i^2\right) \tag{2.3}$$

hence, by (2.2) and (2.3)

$$\det A \le \left(\sum_{i=1}^{N} \lambda_i^2/N\right)^{N/2} = \frac{\|A\|^N}{N^{N/2}}.$$

Finally, if det A < 0 choose R^f to be an orthogonal transformation such that det R' = -1. Then, as det $(R^!A) > 0$, by the first part of the proof we have

$$\|A\|^N = \|R'A\|^N \geq N^N \wedge \text{Idet } (R'A)\text{I} = N^N \text{« Idet AL}$$

Due to the strict concavity of the logarithmic function, it follows immediatly from (2.3) that equality holds if and only if $X\backslash = X2 = — = \wedge N»^{\cdot m}$ which case A is proportional to an orthogonal transformation.

Remark 2.4.

By abuse of language we shall call a matrix A such that $\|A\|^N = N^{N/2}$ Idet Al *harmonic.* In $[R^2$, a matrix A such that $\|A\|^2 = 2$ Idet Al is of the form

either $[._{ba}J$ or $[_b$ J.

Proof of Theorem 2.2. If (u*, v*) *e s4*$\wedge_a$ then, as det Vv*(x) = 1 in ft, as r ≥ N by Lemma 2.3 and by Hölder's inequality we have

$$E(u^*,v^*) := \int \|Vu^*(x)(Vv^*(x)r^1H^r dx$$

$$\geq \text{meas}(\Omega)^{1-r/N} \left(\int_\Omega \|\nabla u^*(x)(\nabla v' Hx))''''^1 \wedge dx\right)^{/N}$$

$$\geq \text{meas}(\Omega)^{1-r/N} \left(N^{N/2} \int_\Omega \text{Idet } Vu^*(x)\text{l } dx\right)^{/N}$$

$$\geq \text{meas}(\Omega)^{1-r/N} \left(N^{N/2} \int_\Omega \text{det } Vu^*(x)\, dx\right)^{r/N}$$

$$= N^{r/2} \text{meas}(\Omega)^{1-r/N} \int_\Omega \text{detVuo}(x)dx\Big)^{r/N}. \tag{2.4}$$

Let

$$\hat{x} := \frac{\text{meas } Q}{\text{meas } u_o(Q)} \wedge v_o := x^{\wedge 1/M} u_o,$$

As vo e Diff*-$^0\wedge$), meas VQ(Q) = meas *Q* and since 3Q e $C^{k+3}>^a$, by Theorem 1 in DACOROGNA & MOSER [3] there exists v € Diffc$\wedge\overline{Q}$, $\overline{v_o(Q)}$) such that

$$\begin{cases} \text{det } Vv(x) = 1 & \text{in } Q. \\ v(x) = v_o(x) & \text{on } BQ \end{cases}$$

and define

$$u := \frac{v}{\lambda^{1/N}} \in C^{k,\alpha}(\overline{\Omega}).$$

Clearly

$$u = uo \text{ on } \partial\Omega$$

and we have

$$E(u,v) := \int_\Omega \|\nabla u(x)(\nabla v(x))^{-1}\|^r \, dx$$

$$= \int_\Omega \|\nabla u(x)(\nabla v(x))^{-1}\|^r \, dx$$

$$= \frac{N^{r/2} \text{meas } (fl)}{\lambda^{r/N}}$$

$$= N^{r/2} \text{ meas } (Q)^{1-\wedge} (\text{meas } u_o(Q))^{r/N}$$

which, together with (2.4), finishes the proof.

Corollary 2.5

Under the hypotheses of Theorem 2.2, and in particular if $r \geq N$, then

$$\inf \{E(u, v) \text{ I } (u, v) \text{ G } £0_{tot}\} = N^{r/2} \inf \left\{ \int_\Omega \text{Idet } Vu(x)l^{r/N} \, dx \text{ I } u = \wedge \text{ on } dQ \text{ and } u \in \text{Diff}^{k,\alpha}(\overline{\Omega}) \right\}.$$

Moreover, if (u*, v*) is a solution then there exist a rotation R(.) and a scalar *X(.)* such that

$$Vu^*(x) = X(x) \, R(x) Vv^*(x) \quad \text{for every } x \text{ e } Q .$$

Proof. As det Vuo > 0, for all u eDiffc$^a(\overline{Q})$ with u = uo on 3Q we have

$$\int_a \text{Idet } Vu(x)l \, dx \geq \int_{JQ} \text{det } Vu(x) \, dxl = \int_{Jft} \text{det } Vu_o(x) \, dx \geq a$$

where

$$a := \inf \{\int_\Omega \text{Jldet } Vu(x)l \, dx \text{ I } u = uo \text{ on } dQ \text{ and } u \text{ € } \text{Diff}^{\wedge a}(\overline{Q})\}.$$

Thus, by Theorem 2.2 we obtain

$$\inf \{E(u, v) \text{ I } (u, v) \text{ G } S\&^\wedge a) = N'tf \text{ measCQ})^{1-\wedge} aJt''. \tag{2.5}$$

On the other hand, as $r \geq N$ using Hölder's inequality we deduce that

$$a^{r/N} = \inf \left\{ \left[\int_\Omega |\det Vu(x)|\, dx \right]^{r/N} \;\; I\, u \wedge \text{on } dQ \text{ and } u \in \mathrm{Diff}^{k,\alpha}(\overline{\Omega}) \right\}$$

$$\leq \mathrm{meas}(\Omega)^{r/N-1} \inf \left\{ \int_\Omega |\det Vu(x)|^{r/N}\, dx \;\; I\, u = UQ \text{ on } dQ \text{ and } u \in \mathrm{Diff}^{k,\alpha}(\overline{\Omega}) \right\}$$

which, together with (2.5) implies that

$$\inf \{E(u, v) \;\; I\, (u, v) \in W_{k,a}\} \leq N^{r/2} \inf \left\{ \int_\Omega |\det Vu(x)|^{r/N}\, dx \;\; I\, u = UQ \text{ on } 8Q \text{ and } u \in \mathrm{Diff}^{k,\alpha}(\overline{Q}) \right\}$$

and the reverse inequality follows immediatly from Lemma 2.3.

Finally, by Lemma 2.3 if (u*, v*) is a solution then

$$\int_\Omega |\det Vu^*(x)|^{r/N}\, dx \geq \inf \left\{ \int_\Omega |\det Vu(x)|^{r/N}\, dx \;\; I\, u = UQ \text{ on } BQ \text{ and } u \in \mathrm{Diff}^{k,\alpha}(\overline{f}) \right\}$$

$$= N^{-r/2} \inf \{E(u, v) \;\; I\, (u, v) \in W_{k,a}\}$$

$$= N^{-r/2} \int_{\Omega} \|Vu^*(x)(Vv^*(x))^{-1}\|^r dx$$

$$\geq \int_\Omega |\det Vu^*(x)|^{r/N} dx$$

and so

$$\int_\Omega \{|\det Vu^*(x)|^{r/N} - N^{-r/2} \|Vu^*(x)(Vv^*(x))^{-1}\|^r\}\, dx = 0 \qquad (2.6)$$

which, together with Lemma 2.3, implies that

$$\|Vu^*(x)\, (Vv^*(x))^{-1}\|^N = N^{N/2} |\det (Vu^*(x)\, (Vv^*(x))^{-1})| \quad \text{a. e. in } \Omega.$$

Thus

$$Vu^*(x) = X(x)\, R(x) Vv^*(x) \quad \text{a. e. in } Q$$

for some rotation R(.) and some scalar *X(.)*. From (2.6), Theorem 2.2 and using Hölder's inequality we deduce that

$$(\mathrm{meas}\, \Omega)^{1-r/N} \left(\int_\Omega |\det Vu^*(x)|^{r/N}\, dx \right)^{N/r} =$$

$$= N^{-r/2} \left[(\mathrm{meas}\, \Omega)^{1+r/N} \inf\{E(u, v) \;\; I\, (u, v) \in W_{k,a}\} \right]^m$$

$$= \int_{\Omega} \det Vu_0(x)\, dx$$

$$= \int_\Omega \det Vu^*(x)\, dx \leq \int_\Omega |\det Vu^*(x)|\, dx$$

$$\leq (\mathrm{meas}\, \Omega)^{1-N/r} \left[\int_\Omega |\det Vu^*(x)|^{r/N} dx \right]^{N/r}.$$

Hence

$$\int_Q [\det Vu^*(x)dx - |\det Vu^*(x)|]dx = 0$$

which implies that det Vu* > 0 in *Q*.

If uo is affine then we can obtain existence of minimizers under less restrictive hypothesis on *BQ*, namely

Proposition 2.6.

Let *Q* be a bounded, open set with Lipschitz boundary. Let uo(x) = Ax + b where A € $M^{N\times N}$ with det A > 0 and b € $[R^N$. Then (P) admits a solution (u, v) *e S&**.

This result relies on the fact that any affine deformation is harmonic up to a volume preserving transformation. Precisely

Lemma 2.7.

If det A * 0 then there exists a matrix B such that det B = 1 and $\|AB\|^N = N^N \# |\det A|$.

Proof. Suppose that det A > 0. As in the proof of Lemma 2.3, we can write

A = RQ^Tdiag(^i,..., ^N)Q> where R and Q are orthogonal transformations and *X*\,..., *X*^ > 0. Set

B:=QTdiag(pi,...,p_N)

where

$$p_i := \frac{(\lambda_1 \cdots \lambda_N)^{1/N}}{\lambda_i}.$$

Then det B = 1 and

$\|AB\|_N = (\lambda_1 \cdots \lambda_N)^{1/N} \|A\|_N = (\lambda_1 \cdots \lambda_N)^{1/N} \|A\|^N = N^{N/2} \det A$.

If det A < 0, it suffices to multiply A by an orthogonal transformation R^1 with det R' = -1 and to apply the previous case to the matrix R^fA.

Proof of Proposition 2.6. Setting u* : = uo and **Vv*** := $\mathbf{B^{-1}}$**,** as in the proof of Theorem 2.2 it follows that (u*, v*) is a solution for (P).

3. EXISTENCE AND REGULARITY RESULTS.

Now we show that the results of Section 2 can be generalized in the following way.

Hypothesis (H): Let W : M^{NxN} -»[0, +<») be continuous and such that there exist *Xe* OR and Xo e M^{NxN} with

$$\det X_o = \frac{\text{meas } u_0(Q)}{\text{meas} Q}$$

and

$$W(X) - X \det X \geq W(X_0) - X \det X_o \quad \text{for every } X \in M^{N*N}.$$

Remark 3.1. (i) In some sense the *X* appearing in (H) can be seen as a Lagrange multiplier.

(ii) If W $\in C^1(M^{NxN})$ then

$$\begin{cases} \dfrac{\partial W}{\partial X}(X_0) = \lambda \operatorname{adj} X_0 \\[2ex] \lambda := \dfrac{\left\langle -\dfrac{\partial W}{\partial X}(X_0),\ X\$ \right\rangle \ \text{meas}(Q)}{N \,\text{meas}\, i_Q(Q.)}. \end{cases} \tag{3.1}$$

Indeed, as Xo is a minimum for W(X) - *X* det X we have

$$0 = \wedge(W(X) - X \det X)|_{X_o}$$
$$= \frac{\partial W}{\partial X}(X_0) - \wedge \operatorname{adj} X_0$$

and so, by (2.1) and (H)

$$\langle \wedge X_0), X_0 \rangle = X N \det X_0$$
$$= \lambda N \frac{\text{meas up}(Q)}{\text{meas} Q}$$

which proves (3.1).

Consider the problem

(P) minimize in $\mathcal{A}_{k,\alpha}$ the functional

$$E(u, v) := \int_\Omega W(\nabla u(x)(\nabla v(x))^{-1})\, dx$$

where the class of admissible pairs is defined by

$$\mathcal{A}_{k,\alpha} := \{(u, v) \in \mathrm{Diff}^{k,\alpha}(\overline{\Omega}) \times \mathrm{Diff}^{k,\alpha}(\overline{\Omega}) |\ u = u_0 \text{ on } \partial\Omega \text{ and } \det \nabla v(x) = 1 \text{ a. e. in } \Omega\}$$

and, as in Section 2, $u_0 \in \mathrm{Diff}^{k,\alpha}(\overline{\Omega})$ is such that $\det \nabla u_0 > 0$ in $\overline{\Omega}$.

Before stating the main result of this section, we give examples of functions satisfying the condition (H).

Proposition 3.2

The following functions $W : M^{N\times N} \to \mathbb{R}$ verify (H).

i) Let $N \geq 2$, let $g : \mathbb{R}^2 \to \mathbb{R}$ be convex, C^1 and is increasing with respect to the first variable and set

$$W(X) = g(\|X\|^N, \det X).$$

In particular, (H) holds for

$$W(X) = \|X\|^r \text{ if and only if } r \geq N.$$

ii) Let $N = 2$ and let

$$W(X) = \sum_{i,j,k,l=1,2} a_{ijkl}\, X_{ij}\, X_{kl}$$

with $a_{ijkl} = a_{klij}$ and W a strictly rank one convex function, i. e.

$$W(\lambda\otimes\mu) \geq \alpha\, \|\lambda\otimes\mu\|^2$$

for some $\alpha > 0$ and for all $\lambda, \mu \in \mathbb{R}^2$, where $(\lambda\otimes\mu)_{ij} := \lambda_i\, \mu_j$ for $i, j = 1, 2$.

Remark 3.3.

There are other examples of functions satisfying (H), namely for $N = 2$
$W(X) = \|X\|^4 - 2(\det X)^2$ and $W(X) = \frac{1}{2}(X_{11}^2 + X_{12}^2 + X_{21}^2) + \frac{1}{4}X_{22}^4$.

Proof of Proposition 3.2. i) Set

XQ := £R with ^N = meas uo(£2)/ meas(Q) and R a rotation.

As g = g(t, s) is convex we have

W(X) - W(Xo) ≥ |f(HXoll^N, det Xo)(HXII^N - IIXoll^N) +^IIXo1l^N, det Xo)(det X - det X_o),

and so, as -^ ≥ 0 and since by Lemma 2.3

HX‖N ≥ NN/2 detx and IIX0H^N = N^N# det Xo,

we conclude that

W(X) - W(X0) ≥ X(det X - det X_o)

where

X := NN/2 ^(IIX0II^N, det X_o) + 3f<IIXoll^N, det X_o).

If W(X) = IIXII^r then (H) is violated if r < N (see also Proposition 4.1). Indeed, in this case (3.1) reduces to

rIIX0II^r-2 X0 = λ adj X0

and, as Xo must be parallel to its adjugate matrix, Xo is a harmonic matrix and

Xo = % R for some % € [R, % > 0 and some rotation R.

If in (H) we set X = p R, with p € [R, p > £, then we obtain

$$p^r N^{1/2} - £^r N^{1/2} > X(\$F - £^N) \tag{3.2}$$

and so, either *X* ≤ 0 and then (3.2) fails for p < £, or *X* > 0 and (3.2) is false for p large enough,

(ii) Since W is rank one convex and as N = 2, then W is polyconvex (see DACOROGNA [2]) and so

$$\sup\{\wedge \wedge l \det Y < O\} \leq \inf\{\wedge : l \det Y > 0\}. \tag{3.3}$$

Choose

$$\lambda = \inf\{\frac{W(Y)}{\det Y} \mid \det Y > 0\} \tag{3.4}$$

and observe that the infimum is attained. Indeed, since W is quadratic there is no loss of generality in assuming that a minimizing sequence has norm 1 and so, up to the extraction of a subsequence, we have

$$\frac{W(Y_n)}{\det Y_n} \to \lambda,\ Y_n \to X,\ \det Y_n > 0 \text{ and } \|Y_n\| = 1. \tag{3.5}$$

Note that $\det X > 0$ otherwise $X = a \otimes b$ for some unit vectors a, b, and using the strict rank one convexity and (3.5) we would have for every $\varepsilon > 0$

$$\frac{W(Y_n)}{\varepsilon + \det Y_n} \to \frac{W(X)}{\varepsilon + \det X} \geq \frac{\alpha}{\varepsilon}.$$

Letting $\varepsilon \to 0^+$ we would obtain

$$\frac{W(Y_n)}{\det Y_n} \to +\infty$$

which contradicts (3.3), (3.4), (3.5). Hence $\det X > 0$ and setting

$$X_0 := \xi X \text{ where } \xi^2 := \frac{\text{meas } u_0(\Omega)}{\text{meas } \Omega} \frac{1}{\det X} = \frac{\lambda}{W(X)} \frac{\text{meas } u_0(\Omega)}{\text{meas } \Omega}$$

it follows from (3.3) that

$$W(Y) - \lambda \det Y \geq 0 = W(X_0) - \lambda \det X_0.$$

and

$$\det X_0 = \frac{\text{meas } u_0(\Omega)}{\text{meas } \Omega}.$$

Theorem 3.4.

Let $k \geq 1$ be an integer, $0 < \alpha < 1$, Ω a bounded, open set with $C^{k+3,\alpha}$ boundary and let $u_0 \in \text{Diff}^{k,\alpha}(\overline{\Omega})$ with $\det \nabla u_0 > 0$ in $\overline{\Omega}$. If (H) holds then (P) admits a solution $(u^*, v^*) \in \mathscr{A}_{k,\alpha}$ such that

$$\nabla u^*(x)\, (\nabla v^*(x))^{-1} = X_0 \quad \text{for every } x \in \Omega$$

and

$$\inf \{ \int_\Omega W(\nabla u(x)(\nabla v(x))^{-1})\, dx \mid (u, v) \in \mathscr{A}_{k,\alpha} \} = W(X_0) \text{ meas } \Omega.$$

Remarks 3.5.

(i) As it will become clear in Section 4, in some sense the condition (H) is optimal to guarantee existence of solution.

(ii) The set $S\&\wedge_a$ of admissible pairs of functions (u, v) was chosen so as to give immediatly a regularity result as well as existence of solution.

(iii)If

$$W(X) = \|X\|^r$$

for r ≥ N, by Proposition 3.2 i) we can take Xo = XR where R is a rotation and

$$y_N = \frac{\text{meas up}(Q)}{\text{meas } Cl}$$

Then, according to Theorem 3.4 we can find a minimizer (u*, v*) such that

$$Vu^*(x)(Vv^*(x))^{-1} = X\,R \quad \text{a. e. in } Q$$

and the minimum value of the energy functional is given by

$$W(X_0)\text{ meas } Q = \|XR\|^T \text{ meas } Q$$

$$= X^T N^r \# \text{ meas } Cl$$

$$= N\wedge \text{ meas } Cl^1\wedge (\text{meas uo(ft)})^{r/N}$$

which is in agreement with Theorem 2.2 and Corollary 2.5.

Before giving the proof of Theorem 3.4 we state a theorem which is proved exactly as the preceding one but requires less regularity on *dQ* (see also Proposition 2.6).

Theorem 3.6.

Let *Q*. be a bounded, open, Lipschitz domain, let uo(x) = Ax + b where A **e** $M^{N\times N}$ and b € $[R^N$ and assume that W satisfies (H). If det A * 0 then (P) admits a solution (u*, v*) € *Stf** with u* = uo on *dQ* and det Vv*(x) = 1 in Q. Furthermore, if det A = 0 and if W(X) -> 0 when 11X11 -> 0 then (P) has no solution unless A = 0.

Proof of Theorem 3.4.

Let X_o **e** $M^{N\times N}$ be a matrix for which (H) holds. By Theorem 1 in DACOROGNA & MOSER [3], we find a mapping u* **e** Diffc^$\bar{Q}$) such that

$$\begin{cases} \det Vu^*(x) = \det X_o & \text{in } Q \\ u^*(x) = UQ(X) & \text{on } 3Q. \end{cases}$$

Setting

$$v^* := \wedge\ u^*.$$

we have

$$\det V\, v^*(x) = 1 \text{ in } Q,\ Vu^*(x)\,(Vv^*(x))\text{-!} = Xo \text{ in } Q,$$

and, by (H), if (u, v) € £0k,a we have

$$\mathbf{JwtVuCxXVvtx))\text{-}^{1})\ dx \geq}$$

$$\mathbf{d}$$

$$\mathbf{fw(Vu^*(x)(Vv^*(x))\text{-}^{1})\ dx + X\ J[\det Vu(x) - \det Vu^*(x)]\ dx.} \qquad \mathbf{(3.6)}$$

As u = u* on *dQ*. we obtain

$$\mathbf{f}\ (\det Vu(x)\text{-}\det Vu^*(x))dx = O$$

which, together with (3.6) implies that

$$\mathbf{f\ W(Vu(x)\ (VvCx))''^{1})\ dx \geq f\ W(Vu^*(x)(Vv^*(x)r^{1})dx = meas(Q)W(X_0).}$$

The proof of Theorem 3.6 requires the following lemma.

Lemma 3.7.

If det A = 0 then there exists a family of matrices $B_£$ such that det $B_£$= 1 and $\|AB_£\|$ -» 0 as $\varepsilon \to 0^+$.

Proof. Using the polar decomposition for A we can assume that

$$A = R\ \mathrm{diag}(\wedge i, \ldots, X\text{-}_u \ldots, \lambda_N)$$

where R is a rotation and *X{* = 0. Set

$$B_£ := \mathrm{diag}(bi(e),\ldots, b_N(e))$$

where

$$b_j(\varepsilon) := \begin{cases} 1 & \text{if } \lambda_j = 0, \\ \varepsilon/\lambda_j & \text{if } \lambda_j \neq 0 \end{cases}$$

if $j \neq i$ and

$$b_i(\varepsilon) := \frac{1}{b_1(\varepsilon)\ldots b_{i-1}(\varepsilon) b_{i+1}(\varepsilon)\ldots b_N(\varepsilon)}.$$

Clearly det $B_e = 1$ and $\|AB_E\| \leq (N-l)^{1/2}e$.

Proof of Theorem 3.6. Suppose that det A * 0. Setting u* := uo and v*(x) := sign (det A) XQ^1 U*, by (H) it foUows that

det Xo = Idet Al

and so

det Vv* = 1 in Q.

As in the proof of Theorem 3.4, we conclude that

inf {J\V(Vu(x)(Vv(x))-^{1}) dx I (u, v) e ^ $_{t\ a}$ } = W(X_0) meas Q = E(u*, v*).

Finally, if det A = 0 with A * 0, by Lemma 3.7 consider a sequence {B_e} such that det $B_£$= 1 and $\|AB_E\|$ -> 0 as e -» 0+. Setting $u_£$ = uo and Vv_E = $B''^{1}_{£}$ we obtain

$E(u_e, v_£)$ = meas (Q) W($AB_£$) -^ 0.

It is clear that in this case (P) admits no solution since, if

E(u*, v*) = inf {E(u, v) I (u, v) e *S#$_{Ka}$*} = 0,

then Vu*(x) (Vv*(x))"^{1} = 0 in Q, i. e. Vu*(x) = 0 in £1 Hence u* must be constant and as A * 0, this is in contradiction with the condition u* = uo on $\partial\Omega$.

Finally, we conclude this section with a result on the weak lower semicontinuity of E(.,.). However, we insist that this property is not sufficient to ensure existence since, in general, no weak compactness can be obtained for the minimizing sequences regardless of the coercivity of W.

Proposition 3.8

Let $Q \subset \mathbb{R}^N$ be a bounded, open set. Let $p \geq 1$, $q \geq N$ and $1/p + (N-l)/q \leq 1$. If

$(u_\pounds, v_\pounds) \to (u, v)$ weakly in $W^{1\wedge} \times W^{1\wedge}$

and if $\det Vv_e = 1$ a. e. in Cl then $\det Vv = 1$ a. e. in Q, and

$Vu_e(x)\ (Vve(x))^{-1} \to Vu(x)\ (Vv(x))^{-i}$ weakly in IA

Consequently, if $W : M^{N\times N} \to [0, -H^*>)$ is convex then

$$\int_\Omega W(\nabla u(x)(\nabla v(x))^{-1})\, dx \leq \liminf_{\mathcal{M}} JwCVueCxXVv^\wedge x))^{-1})\, dx.$$

Conjecture.

In Proposition 3.8 we used the convexity of W to obtain the weak lower semicontinuity of the energy functional E(.,.). As, formally, $Vu\ (Vv)^{-1}$ is the gradient of $u \circ v^1$, we conjecture that if W is quasiconvex and if (u_e, v_e) converges weakly to (u, v) then

$$\int_d w(Vu(x)(Vv(x))^{-1})\, dx \leq \liminf_d \mathbf{Jw(Vu_e(x)(Vv_e(x))^{-i})}\, dx.$$

Proof of Proposition 3.8. As $\det Vv = 1$ a. e. in **£2,** we have

$(Vu(x)\ (Vv(x))^{-i})ij = (Vu(x)\ (\mathrm{adj}\ Vv(x))T)_{ij} = m(x).\ ^\wedge(x)$

where r|i is the gradient of the i* component of u and £j is the j^A row of adj Vv. Hence

curl r|i = 0 and div £j = 0

and by the div-curl lemma (see TARTAR [12]) we conclude that if $p \geq 1$, $q \geq N$ and if $1/p + (N-l)/q \leq l$ then

$Vu_\pounds(x)\ (Vv^\wedge x))^{-1} \to Vu(x)\ (Vv(x))^{-1}$ weakly in L^1 (3.7)

whenever

- $(u_\pounds, v_\pounds) \to (u, v)$ weakly in $W^{1*} \times W^{1\wedge}$.

Finally, if W is a convex, nonegative function then by (3.7) the functional (u,v) -» E(u,v) is lower semicontinuous (see DACOROGNA [2]).

Remark 3.9.

Let $p \geq 1$, $q \geq N$ and $1/p + (N-l)/q \leq 1$ and consider the class of admissible pairs to be given by

$$\mathcal{A}_{p,q} := \{(u, v) \in W^{1,p}(Q; \mathbb{R}^N) \times W^{1,q}(Q; \mathbb{R}^N) \mid u = u_0 \text{ on } \partial Q, \det \nabla v(x) = 1 \text{ a. e. in } Q$$

$$\text{and } \int_\Omega v(x)\, dx = 0 = \int_\Omega x\, dx\}.$$

Suppose that W is convex, $W(X) \geq C_1 |X|^r - C_0$ with $C_1 > 0$. By Proposition 3.8 (P) has a solution in $\mathcal{A}_{p,q}$ if there is a minimizing sequence $\{(u_\varepsilon, v_\varepsilon)\}$ bounded in $W^{1,p} \times W^{1,q}$. Suppose that $r \geq p \frac{q}{q-1}$ and that $\{v_\varepsilon\}$ is bounded in $W^{1,q}$. Let s be such that

$$\frac{q}{q-1} \leq s \leq \frac{r}{p}.$$

By Hölder's inequality

$$\int_\Omega |\nabla u_\varepsilon(x)|^p dx = \int_\Omega |\nabla u_\varepsilon(x)| |\det \nabla v_\varepsilon(x)|^p dx$$

$$\leq \left(\int_\Omega |\nabla u_\varepsilon(x)|^r dx\right)^{1/s} \left(\int_\Omega |\det \nabla v_\varepsilon(x)|^{s'} dx\right)^{1/s'}$$

$$\leq \text{Const.} \left(\int_\Omega |\nabla u_\varepsilon(x)|^r dx\right)^{1/r} \left(\int_\Omega |\det \nabla v_\varepsilon(x)|^q dx\right)^{1/q}$$

and so $\{u_\varepsilon\}$ is bounded in $W^{1,p}$. We conclude that if there exists a minimizing sequence $\{(u_\varepsilon, v_\varepsilon)\}$ where $\{v_\varepsilon\}$ is bounded in $W^{1,q}$ then (P) admits a solution in $\mathcal{A}_{p,q}$.

We next show that the set of solutions is weakly closed.

Proposition 3.10.

Let Q be an open, bounded, Lipschitz domain, let W be a convex function, let $u_0 \in W^{1,p}(\Omega; \mathbb{R}^N)$ and let $p \geq 1$, $q \geq N$, $1/p + (N-l)/q \leq 1$ and $r \geq 1$. If $\{(u_n, v_n)\}$ is a sequence of solutions of (P) in $\mathcal{A}_{VA}$ and if (u_n, v_n) converges weakly to (u, v) in $W^{1,p} \times W^{1,q}(Q; \mathbb{R}^N) \times W^{1,q}(Q; \mathbb{R}^N)$ then (u, v) is also a solution of (P).

Proof. As $q \geq N$ standard results imply that

adj Vv_n converges weakly in $L^{\wedge -1}$) to adj Vv.

Moreover, as

det Vv_n converges in the sense of distributions to det Vv,

we must have

det $Vv = 1$ a. e. in Q.

and so

$(u, v) \in Stf_{p,q}$.

Finally, using the div-curl lemma we deduce that

$Vu_n(x)\ (Vvn(x))^{-1} \to Vu(x)\ (Vv(x))^{-1}$ weakly in L^1

and as W is convex we conclude that

$$\int_d W(Vu(x)(Vv(x))^{-1})\,dx \leq \liminf \int_d W(Vu_n(x)(Vv_n(x))^{-1})\,dx$$
$$= \inf\{E(u, v) \mid (u, v) \in \wedge_{p,q}\}.$$

4. NON EXISTENCE RESULTS.

In this section we present two types of non existence results showing that, despite the resemblance of our problem to the classic Dirichlet problem of minimizing $\int_\Omega |Vu|^p$, problem (P) is in fact very different in nature. It turns out to be much closer to

(Q) $\inf\{\int_\Omega |\det Vu(x)|^p\,dx \mid u = u_0 \text{ on } \partial\Omega,\ u \in \mathbf{Diff}^{k,\alpha}(\bar{\Omega})\}$

as already seen in Corollary 2.5 and as it will be illustrated bellow. Indeed, restricting our attention to

$W(X) = \|X\|^r, r>1,$

Theorem 3.6 provides a first type of non existence result. Namely, if $u_0(x) = Ax$ for some $A \in M^{N\times N}$, $A \neq 0$ with $\det A = 0$, then (P) does not admit a solution. This is in sharp contrast with the minimization of $\int_\Omega |Vu|^p$.

We have seen in Theorem 2.2 and Corollary 2.5 that if $r \geq N$ then (P), as well as (Q) (with $p = r/N \geq 1$), admit solutions. Now we show the second type of non existence result, proving that if $r < N$ then neither (P) (see Proposition 4.1) nor (Q) (see Proposition 4.3) have solutions.

Proposition 4.1.

Let $Q = \{x \in \mathbb{R}^2 \mid \|x\| < 1\}$, let $u_0(x) = x$ and let $0 < r < 2 = N$. Then

$$\inf\{ \int_Q |Vu(x)(Vv(x))^{-1}|^r dx \mid (u, v) \in W^{1,\infty}(Q) \times W^{1,\infty}(Q),\ u = u_0 \text{ on } \partial Q,\ \det Vv = 1 \text{ a. e. in } Q\} = 0$$

and hence the infimum is not attained.

Remarks 4.2.

i) In order to avoid some technicalities, in the previous proposition we considered u and v in $W^{1,\infty}(\Omega)$. However, the result remains valid if instead we assume that the admissible pairs $(u, v) \in \mathrm{Diff}^{k,\alpha}(\bar{\Omega}) \times \mathrm{Diff}^{k,\alpha}(\bar{\Omega})$.

ii) Similarly, we take the boundary condition $u_0(x) = x$ just for the sake of illustration, since it could be replaced by any $u_0 \in \mathrm{Diff}^{k,\alpha}(\bar{\Omega})$.

Proof of Proposition 4.1. Using polar coordinates we define

$$u_n(x, y) := \begin{cases} \frac{1}{\varepsilon}(x, y) & \text{if } r \in (0, \varepsilon) \\ \frac{1}{r}(x, y) & \text{if } r \in (\varepsilon, 1), \end{cases}$$

where $\varepsilon := (2n)^{k/(k-2)} \to 0$, and

$$v_n(x, y) := \frac{r}{2n}(-\cos 2n\theta, \sin 2n\theta)$$

where $r = \sqrt{x^2 + y^2}$. A direct computation gives

$$\det Vv_n(x) = 1,$$

$$\|\nabla u_n(x)(\nabla v_n(x))^{-1}\|^k = \begin{cases} \frac{1}{\varepsilon^k}(1 \quad , f. , \quad n < r < e \\ \frac{1}{r^k}\left(\frac{1}{2n}\right)^{k/2} & \text{if } e < r < 1, \end{cases}$$

and

$$E(u», V_n) = n(2n + \wedge)^{k/2} (2n)^{-r} + \pi\frac{2(2n)^{-k/2}}{2 - r}(1 - (2n)^{-k}) \to \text{Oas } n -» +\infty.$$

Finally, we conclude this section with a similar result on problem (Q).

Proposition 4.3.

If an e $C^{3,a}$ for some $0 < a < 1$, if uo € DifiP^C$\overline{Q}$) with det Vuo > 0 in $\overline{Q}$ and if $0 < \beta <$ 1 then for all $p \geq 1$

$$\inf_{J} \int_{\Omega} |\det Vu(x)|^p \, dx \mid u = UQ \text{ on } dQ \text{ and } u \in \mathrm{Diff}^{1>a}(\overline{Q})| = 0$$

and thus the infimum is not attained[1].

Proof. Let xo e Q and let $\overline{B}$(xo, 2e) C Q. Let φ_n be a family of smooth functions such that $0 \leq \varphi \leq 1$ and

$$\varphi_n(t) = \begin{cases} \backslash & \text{if } t \leq 1 \\ 0 & \text{if } t \geq e^{1/n} \end{cases}$$

and define

$$\ldots) := \varphi_n\left(\left|\frac{x - x_0}{\varepsilon}\right|^2\right)\left(1 + \left|\frac{x - x_0}{\varepsilon}\right|^{2n}\right) + 1 - \varphi_n\left(\left|\frac{x - x_0}{\varepsilon}\right|^2\right).$$

Clearly, $f_n \geq 0$, f_n are smooth and $f_n \geq 1$. In addition,

$$G\int_a f_n(x)dx \quad \int_J \quad \int_{\wedge ix-x_o i<\varepsilon} \quad \left|\frac{x - x_0}{\varepsilon}\right|^{2n} dx \quad \int_{yJo} \quad \left|\int^{f^1} R^{N-1} R^{2n} dx\right)^{\beta}$$

---

[1] rhe same result holds for u e W^{1*}, with $p \geq PN$.

$$= \text{Const.} \frac{1}{(N + 2n)^\beta} \tag{4.1}$$

and

$$\int_\Omega f_n(x)^\beta \, dx \le \text{Const.} + \int_{|x-x_0|<\varepsilon} \left|\frac{x - x_0}{\varepsilon}\right|^{2n\beta} dx$$
$$= \text{Const.} + \text{Const.} \frac{1}{N + 2n\beta},$$

and so, from (4.1) we conclude that

$$\lim_{n \to \infty} \frac{\int_\Omega f_n(x)^\beta \, dx}{\left(\int_\Omega f_n(x) \, dx\right)^\beta} \le \lim_{n \to \infty} \text{Const.}\left(1 + \frac{1}{N + 2n\beta}\right)(N + 2n)^\beta = 0. \tag{4.2}$$

Using Theorem 1 in DACOROGNA & MOSER [3], we find a sequence $u_n \in \text{Diff}^{1,\alpha}(\overline{\Omega})$ such that

$$\begin{cases} \det \nabla u_n(x) = \dfrac{\text{meas } u_0(\Omega)}{\int_\Omega f_n(x) \, dx} f_n(x) & \text{in } \Omega \\ \\ u_n(x) = u_0(x) & \text{if } x \in \partial\Omega. \end{cases}$$

From (4.2) it follows that

$$\lim_{n \to \infty} \int_\Omega |\det \nabla u_n(x)|^\beta \, dx = \text{meas } u_0(\Omega)^\beta \lim_{n \to \infty} \frac{\int_\Omega f_n(x)^\beta \, dx}{\left(\int_\Omega f_n(x) \, dx\right)^\beta} = 0.$$

5. QUALITATIVE PROPERTIES.

We remark that if (P) has one solution then, if $\partial\Omega$ is sufficiently smooth[2], there are uncountably many solutions. In fact, if

$$\min E(u, v) = E(u_1, v_1)$$

---

[2]If the class of admissible functions is $\mathcal{A}_{k,\alpha}$ then $\partial\Omega$ must be $C^{k+3,\alpha}$. If we are considering the set $\mathcal{A}_{p,q}$ then we assume that $\partial\Omega$ is Lipschitz.

and if f is such that[3]

$$\begin{cases} \det \nabla f(x) = 1 & \text{in } \Omega \\ f(x) = x & \text{on } \partial\Omega, \end{cases}$$

then $(u_1 \circ f, v_1 \circ f)$ is admissible and, as $f(\Omega) = \Omega$ we obtain

$$\begin{aligned} E(u_1 \circ f, v_1 \circ f) &= \int_\Omega W(\nabla u_1(f(x))\nabla f(x)\,(\nabla f(x))^{-1}(\nabla v_1(f(x)))^{-1})\,dx \\ &= \int_\Omega W(\nabla u_1(f(x))(\nabla v_1(f(x)))^{-1})\,dx \\ &= \int_{f(\Omega)} W(\nabla u_1(y)(\nabla v_1(y))^{-1})\,dx \\ &= E(u_1, v_1). \end{aligned}$$

In Remark 3.9 we noted that if $W(X) = \|X\|^r$, $r \geq N$, and if there exists a minimizing sequence $\{(u_\varepsilon, v_\varepsilon)\}$ where $\{v_\varepsilon\}$ is bounded in $W^{1,q}$ then (P) admits a solution in $\mathscr{A}_{p,q}$. By the preceeding remark, it would suffice to show that given a sequence $\{v_\varepsilon\}$ in $W^{1,q}$ then there exists a sequence $f_\varepsilon \in W^{1,\infty}(\Omega, \Omega)$ such that

$$\begin{cases} \det \nabla f_\varepsilon(x) = 1 & \text{in } \Omega \\ f_\varepsilon(x) = x & \text{on } \partial\Omega, \end{cases}$$

and $\{v_\varepsilon \circ f_\varepsilon\}$ is bounded in $W^{1,q}$. However, such sequence $\{f_\varepsilon\}$ may fail to exist since (P) has no solution if $u_0(x) = Ax + b$, $b \in \mathbb{R}^N$, $A \in M^{N\times N}$, $\det A = 0$ and $A \neq 0$ (see Theorem 3.6).

As we mentioned before, the minimization of (P) corresponds, formally, to a minimization of a functional where the domain is varying. Theorem 3.4 provides a sufficient condition under which there is existence of solution. Here, $v(\Omega)$ becomes the domain of the solution. It is natural to ask what type of domains may correspond to solutions of (P). The following proposition partially answers that question.

---

[3]Here $f \in C^{k,\alpha}$ in the case where the class of admissible functions is $\mathscr{A}_{k,\alpha}$ and f is Lipschitz if we are considering the class $\mathscr{A}_{p,q}$.

Proposition 5.1.

Let $k \geq 1$ be an integer, $0<cc < 1$, let Q ^ [R^2 be a bounded, open set with $C^{k+3,ct}$ boundary and let u_0 € Diff^C$\overline{Q}$) with det $Vu_0 > 0$ in $\overline{Q}$. Let W(X) = IIXII^2 and assume that 3uo(f2) is an analytic Jordan curve. If Y C IR^2 is such that meas Y = meas *Q* and if 9Y is an analytic Jordan curve then there exists a minimizer (u, v) of E(.,.) on £#k,a such that *v(Q)* is a translation of Y.

Proof. By the Riemann Mapping Theorem there exists a conformal equivalence f *e* Diff^~($\overline{Y}$, $\overline{u_o(£2)}$). Thus we have f = (fi, *fi)* where

$$\begin{cases} \dfrac{\partial f_1}{\partial y_1} = \dfrac{3f_2}{dy_2} \\[2ex] \dfrac{\partial f_1}{\partial y_2} = -\dfrac{3f_2}{\partial y_1}. \end{cases} \tag{5.1}$$

Set

vorsf^uoiQ^BCO, R).

As v_0 € Diff^{k,ot}($\overline{ii}$), we have

meas vo(Q) = meas Y = meas Q

and since dQ e C^{k+3}-^{a}, by Theorem 1 in DACOROGNA & MOSER [3] there exists vi € Diff$^{k,\alpha}$(S; $\overline{Y}$) such that

$$\begin{cases} \det Vv!(x)=l & \text{in} Q \\[1ex] v_2(x) = v_o(x) & \text{on } \partial\Omega. \end{cases}$$

Finally, set

v(x) := vi(x) + C, where the constant C is such that

$$\int v(x)dx = 0,$$

and define

u := fo(v - C) € $C^{k,\alpha}(\Omega)$.

Clearly

$u = u_o$ on $\partial\Omega$

and as v is invertible (see BALL [1], Theorems 1 **and** 2), we have

$$E(u, v) := \int \| Vu(x)(Vv(x))^{-1} \|^2 \, dx$$

$$= \int \| Vu(x)(Vv(x))^{-1} \|^2 \, dx$$

$$= \int \| \nabla u(v_1^{-1}(y)) \, \nabla v_1^{-1}(y) \|^2 \, dy$$

$$= \int \| Vf(y) \|^2 \, dy.$$

Therefore, by (5.1) we deduce that

$$E(u, v) = 2 \int \left[\left(\frac{\partial f_1}{\partial y_1}(y) \right)^2 + \left(\frac{\partial f_1}{\partial y_2}(y) \right)^2 \right] dy$$

$$= 2 \int \det Vf(y) \, dy$$

$$= 2 \int \det Vu(v_1^{-1}(y)) \det Vv_1^{-1}(y) \, dy$$

$$= 2 \int \det Vu(x) \, dx$$

$$= 2 \int_\Omega \det Vu_o(x) \, dx. \qquad (5.2)$$

From (2.4) and (5.2) we deduce that (u, v) is a solution of (P) and

$$v(Q) = v_o(Q) + C = Y + C.$$

Next, and pursuing the discussing of the nature of the set of solutions of (P), we give some uniqueness results.

Proposition 5.2.

Let Q be an open, bounded, Lipschitz domain in $\mathbb{R}^N$, let $r \geq N$ and let $p \geq r \geq N$, $q \geq N$. $W(X) = \|X\|^r$ and if (u_1, v) and (u_2, v) are solutions of (P) in $Pt_{f,p,q}$ then $u_1 = u_2$ a. e. in Q.

Proof. Clearly, if 6 e (0, 1) then (0ui + 8u_2, v) is admissible and

JllV(eui+eu2)(x)(Vv(x))-^{1}ll^{r}dx < 6 JllVui(x)(Vv(x))-^{1}ll^{r}dx

d ***d***

+ (1 - 0) JllVu$_2$(x)(Vv(x))-i|l^{r}dx

d

unless Vui(x)= Vu$_2$(x) a. e. in Q, and so, as ui = u_2 on *dQ.* we conclude that ui = u_2 a. e. in Q.

Proposition 5.3.

Let Q be an open, bounded, Lipschitz domain in [R^N, let r ≳ N and let p ≳ r ≳ N, q > N. If W(X) = IIXII^r and if (u, vi) is a solution of (P) in £0pq such that vi is invertible and *vi(Q)* is a Lipschitz domain[4], then (u, v_2) is another solution of (P) if and only if there exist a constant rotation R and a constant C € [R^N such that v_2(x) = Rvi(x) + C a. e. in *Cl.*

Proof. Suppose that v_2(x) = Rvi(x) + C a. e. in Q. By Corollary 2.5

∇u(x) = λ(x) Q(x) Vvi(x) a.e. in Q

for some rotation Q(.) and some scalar *X(.).* Hence,

Vu(x) = *X(x)* Q(x)R^{T}Vv_2(x) a. e. in *Q*

and so

JllVu(x)(Vv_2(x))-i|l^r dx = N^r/2 JlA.(x)l' dx

= JllVu(x)(Vvi(x))-i|l'dx

and so, (u, v_2) is also a minimizer. Conversely, if (u, vi) and (u, v_2) are solutions of (P) then by Corollary 2.5 we must have det Vu > 0, Vu(x) = Xi(x)Qi(x)Vvi(x) and Vu(x)=X_2(x)Q2(x)Vv_2(x) a. e. in £2, where *X*\, X_2 € [R and Qi, Q2 are rotations. Thus *X*\$_9$ *X2* > 0,

---

[4]Here we will use the fact that if v e W^1*, q > N, v is invertible, v(ft) is a strongly Lipschitz domain and if detVv = 1 a. e. then

(i) v^1 € W^1^^{1}), *Vv^l(y)* = (Vv(x))"^{1} a. e., where y = v(x);**

(ii) Wev € W^1-^{1} and V(w$_o$v)(x) = Vw(v(x)) Vv(x) a. e. in Q, whenever w € W^1*, p ≥ q/(q-l).

$$\mathbf{Vvi(x)} = \wedge^{1}(x)X_2(x)Qy(x)Q2(x)Vv_2(x) \quad \textbf{a.} \text{ e. in } \boldsymbol{Q}$$

and as det Vvi(x) = 1 we have

$$X\backslash(x) = \wedge 200 \quad \text{a. e. in } Q.$$

We conclude that

$$Vv_2(x)(V_{V1}(x))\text{-i} = Ro(x) \tag{5.3}$$

for some rotation R(.). Setting

$$0)_2(y) \ :=v_2(v;^{1}(y)) \quad \text{and} \quad \tilde{R}_0(y) := RoCv\wedge Cy)) \ ,$$

(5.3) reduces to

$$Vo>2(y) = \tilde{R}o(y) \quad \text{a. e. } y \in vi(Q)$$

and we conclude that (see FONSECA [9], Proposition A.I)

$\tilde{R}_o(.)$, and therefore Ro, must be constantly equal to a fixed rotation R

which, together with (5.3) implies that

$$v_2(x) = Rvi(x) + C \text{ a. e. in } Q.$$

Acknowledgments. This work started while the authors were visiting Heriot-Watt University in June 1990. The research of the second author was partially supported by the National Science Foundation under Grant No. DMS - 8803315, and it was partly done during the author's visit to the University of Bath (U. K.) in May-July 1990.

REFERENCES

[1] BALL, J. M. "Global invertibility of Sobolev functions and the interpenetration of matter", Proc. Roy. Soc. Edinburgh **88A** (1981), 315-328.

[2] DACOROGNA, B. ***Direct Methods in the Calculus of Variations,*** **Springer, 1989.**

[3] DACOROGNA, B. & MOSER, J. "On a partial differential equation involving the jacobian determinant", Ann. Inst. H. Poincare", Anal. Nonline"aire 7 (1990), 1-26.

[4] DAVINI, C. "A proposal for a continuum theory of defective crystals", Arch. Rat. Mech. Anal. 96 (1986), 295-317.

[5] DAVINI, C. & PARRY, G. "On defect-preserving deformations in crystals", Int. J. of Plasticity 5 (1989), 337-369.

[6] DAVINI, C. & PARRY, G. "A complete list of invariants for defective crystals", to appear.

[7] ERICKSEN, J. L. "Twinning of crystals I", in *Metastability and Incompletely Posed* ***Problems, IMA Vol. in Appl. Math* 3 (1987), 77-96.**

[8] FONSECA, I. "Variational methods for elastic crystals", Arch. Rat. Mech. Anal. 97 (1987), 189-220.

[9] FONSECA, I. "Phase transitions of elastic solid materials", Arch. Rat. Mech. Anal. **107** (1989), 195-223.

[10] FONSECA, I. & PARRY, G. "Equilibrium configurations of defective crystals", to appear.

[11] KINDERLEHRER, D. "Twinning of crystals H", in *Metastability and Incompletely Posed* ***Problems, IMA Vol. in Appl. Math* 3 (1987), 185-211.**

[12] TARTAR, L. "Compensated compactness and applications to partial differential equations," Nonlinear analysis and mechanics: Heriot-Watt Symposium, vol. IV, Res. Notes in Math. 39, Pitman, 1979, 136-212.

UNCLASSIFIED
SECURITY CLASSIFICATION OP THIS PAGE

MASTER COPY - FOR REPRODUCTION PURPOSES

REPORT DOCUMENTATION PAGE

I*. KtrVKI itLUKIIY CLASSIFICATION Unclassified	1b. RESTRICTIVE MARKINGS
2a. SECURITY CLASSIFICATION AUTHORITY	3 . DISTRIBUTION /AVAILABILITY OF REPORT Approved for public release; distribution unlimited.
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NAMS-8	S. MONITORING ORGANIZATION REPORT NUMBER(S)
6a. NAME OF PERFORMING ORGANIZATION Carnegie Mellon University tb. OFFICE SYMBOL *(If spplksble)*	7a. NAME OF MONITORING ORGANIZATION U, S. Army Research Office
6c ADDRESS *(Qty, Stste, snd ZIP Code)* Department of Mathematics Pittsburgh, PA 15213	7b. ADDRESS *[City, Stste. snd ZPCode)* P.0.Box 12211 Research Triangle Park, NC 27709-2211
6a. NAME OF FUNDING/SPONSORING ORGANIZATION U. S. Army Research Office Sb. OFFICE SYMBOL *Qfsppiksbie)*	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER
8c ADDRESS *(City, Stste, snd ZIP Code)* P.0.Box 12211 Research Triangle Park, NC 27709-2211	10. SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. / PROJECT NO. / TASK NO. / WORK UNIT ACCESSION NO.

11. TITLE *(Include Security Osszficstion)*
A Minimization Problem Involving Variation of the Domain

12. PERSONAL AUTHOR(S) Bernard Dacorogna and irene Fonseca

13*. TYPE OF REPORT	13b. TIME COVERED	14. DATE OF REPORT *(Yesr, Month, Osy)*	15. PAGE COUNT
Technical	FROM_____TO_____	November 1990	28

16. SUPPLEMENTARY NOTATION
The view, opinions and/or findings contained in this report are those of, the author(s). and should not. be construed as, an official Department of the Army position,

17. COSATI CODES			18. SUBJECT TERMS (Continue on *reverse if necesssry snd identify by block number)*
FIELD	GROUP	SUB-GROUP	minimization problems, weak lower semicontinuity, div-curl lemma

19. ABSTRACT (Continue on *reverse if necesssry snd identify by block number)*

Existence, regularity properties and uniqueness of solutions for a minimization problem involving variation of the domain are obtained. This model arises in the study of defective solid crystals and it is shown that direct methods of the Calculus of Variations cannot be applied.

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT D UNCLASSIFIED/UNLIMITED D SAME AS *PT. Q OTIC USERS	21. ABSTRACT SECURITY CLASSIFICATION Unclassified
22a. NAME OF RESPONSIBLE INDIVIDUAL	22b. TELEPHONE *Ondude Ares Code)* / 22c. OFFICE SYMBOL

DD FORM 1473, M MAR

13 APR edition may bt used until exhausted.
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

JUL 2 2 2034

Carnegie Mellon Umvereity Libraries

3 &H&2 DIMES 21bO