

6-21-2015

# Minimum-cost matching in a random bipartite graph with random costs

Alan Frieze

*Carnegie Mellon University*, [af1p@andrew.cmu.edu](mailto:af1p@andrew.cmu.edu)

Tony Johansson

*Carnegie Mellon University*

Follow this and additional works at: <http://repository.cmu.edu/math>

 Part of the [Mathematics Commons](#)

---

This Working Paper is brought to you for free and open access by the Mellon College of Science at Research Showcase @ CMU. It has been accepted for inclusion in Department of Mathematical Sciences by an authorized administrator of Research Showcase @ CMU. For more information, please contact [research-showcase@andrew.cmu.edu](mailto:research-showcase@andrew.cmu.edu).

# Minimum-cost matching in a random bipartite graph with random costs

Alan Frieze\* Tony Johansson†  
Department of Mathematical Sciences  
Carnegie Mellon University  
Pittsburgh PA 15213  
U.S.A.

June 21, 2015

## Abstract

Let  $G = G_{n,n,p}$  be the random bipartite graph on  $n+n$  vertices, where each  $e \in [n]^2$  appears as an edge independently with probability  $p$ . Suppose that each edge  $e$  is given an independent uniform exponential rate one cost. Let  $C(G)$  denote the expected length of the minimum cost perfect matching. We show that w.h.p. if  $d = np \gg (\log n)^2$  then  $\mathbf{E}[C(G)] = (1 + o(1))\frac{\pi^2}{6p}$ . This generalises the well-known result for the case  $G = K_{n,n}$ .

## 1 Introduction

There are many results concerning the optimal value of combinatorial optimization problems with random costs. Sometimes the costs are associated with  $n$  points generated uniformly at random in the unit square  $[0, 1]^2$ . In which case the most celebrated result is due to Beardwood, Halton and Hattersley [3] who showed that the minimum length of a tour through the points a.s. grew as  $\beta n^{1/2}$  for some still unknown  $\beta$ . For more on this and related topics see Steele [18].

The optimisation problem in [3] is defined by the distances between the points. So, it is defined by a random matrix where the entries are highly correlated. There have been many examples considered where the matrix of costs contains independent entries. Aside from the Travelling Salesperson Problem, the most studied problems in combinatorial optimization are perhaps, the shortest path problem; the minimum spanning tree problem and the matching problem. As a first example, consider the shortest path problem in the complete graph  $K_n$  where the edge lengths are independent exponential random variables with rate 1. We denote the exponential random variable with rate  $\lambda$  by  $E(\lambda)$ . Thus  $\mathbf{Pr}(E(\lambda) \geq x) = e^{-\lambda x}$  for  $x \in \mathbf{R}$ . Janson [9] proved (among other things) that if  $X_{i,j}$  denotes the shortest distance between vertices  $i, j$  in this model then  $\mathbf{E}[X_{1,2}] = \frac{H_n}{n}$  where  $H_n = \sum_{i=1}^n \frac{1}{i}$ .

---

\*Research supported in part by NSF Grant DMS1362785. Email: alan@random.math.cmu.edu

†Research supported in part by NSF Grant DMS1362785. Email: tjohanss@andrew.cmu.edu

As far as the spanning tree problem is concerned, the first relevant result is due to Frieze [6]. He showed that if the edges of the complete graph are given independent uniform  $[0, 1]$  edge weights, then the (random) minimum length of a spanning tree  $L_n$  satisfies  $\mathbf{E}[L_n] \rightarrow \zeta(3) = \sum_{i=1}^{\infty} \frac{1}{i^3}$  as  $n \rightarrow \infty$ . Further results on this question can be found in Janson [8], Beveridge, Frieze and McDiarmid [4], Frieze, Ruszinko and Thoma [7] and Cooper, Frieze, Ince, Janson and Spencer [5].

In the case of matchings, the nicest results concern the the minimum cost of a matching in a randomly edge-weighted copy of the complete bipartite graph  $K_{n,n}$ . If  $C_n$  denotes the (random) minimum cost of a perfect matching when edges are given independent exponential  $E(1)$  random variables then the story begins with Walkup [19] who proved that  $\mathbf{E}[C_n] \leq 3$ . Later Karp [10] proved that  $\mathbf{E}[C_n] \leq 2$ . Aldous [1], [2] proved that  $\lim_{n \rightarrow \infty} \mathbf{E}[C_n] = \zeta(2) = \sum_{k=1}^{\infty} \frac{1}{k^2}$ . Parisi [13] conjectured that in fact  $\mathbf{E}[C_n] = \sum_{k=1}^n \frac{1}{k^2}$ . This was proved independently by Linusson and Wästlund [11] and by Nair, Prabhakar and Sharma [12]. A short elegant proof was given by Wästlund [16], [17].

In the paper [4] on the minimum spanning tree problem, the complete graph was replaced by a  $d$ -regular graph  $G$ . Under some mild expansion assumptions, it was shown that if  $d \rightarrow \infty$  then  $\zeta(3)$  can be replaced asymptotically by  $\frac{n}{d}\zeta(3)$ .

Consider a  $d$ -regular bipartite graph  $G$  on  $2N$  vertices. Here  $d = d(N) \rightarrow \infty$  as  $N \rightarrow \infty$ . Each edge  $e$  is assigned a cost  $W(e)$ , each independently chosen according to the exponential distribution  $E(1)$ . Denote the total cost of the minimum-cost perfect matching by  $C(G)$ .

We conjecture the following (under some possibly mild restrictions):

**Conjecture 1.** *Suppose  $d = d(N) \rightarrow \infty$  as  $N \rightarrow \infty$ . For any  $d$ -regular bipartite  $G$ ,*

$$\mathbf{E}[C(G)] = (1 + o(1)) \frac{N \pi^2}{d \cdot 6}.$$

*Here the  $o(1)$  term goes to zero as  $N \rightarrow \infty$ .*

In this paper we prove the conjecture for random bipartite graphs. Let  $G = G_{n,n,p}$  be the random bipartite graph on  $n + n$  vertices, where each  $e \in [n]^2$  appears as an edge independently with probability  $p$ . Suppose that each edge  $e$  is given an independent uniform exponential rate one cost.

**Theorem 1.** *If  $d = np = \omega(\log n)^2$  where  $\omega \rightarrow \infty$  then w.h.p.  $\mathbf{E}[C(G)] \approx \frac{\pi^2}{6p}$ .*

Here  $A_n \approx B_n$  iff  $A_n = (1 + o(1))B_n$  as  $n \rightarrow \infty$  and the event  $\mathcal{E}_n$  occurs with high probability (w.h.p.) if  $\mathbf{Pr}(\mathcal{E}_n) = 1 - o(1)$  as  $n \rightarrow \infty$ .

Applying results of Talagrand [14] we can prove the following concentration result.

**Theorem 2.** *Let  $\varepsilon > 0$  be fixed, then*

$$\mathbf{Pr} \left( \left| C(G) - \frac{\pi^2}{6p} \right| \geq \frac{\varepsilon}{p} \right) \leq n^{-K},$$

*for any constant  $K > 0$ .*

## 2 Proof of Theorem 1

We find that the proofs in [16], [17] can be adapted to our current situation. Suppose that the vertices of  $G$  consist of  $A = \{a_i, i \in [n]\}$  and  $B = \{b_j, j \in [n]\}$ . Let  $C(n, r)$  denote the expected cost of the minimum cost matching

$$M_r = \{(a_i, \phi_r(a_i)) : i = 1, 2, \dots, r\} \text{ of } A_r = \{a_1, a_2, \dots, a_r\} \text{ into } B.$$

We will prove that w.h.p.

$$C(n, r) - C(n, r-1) \approx \frac{1}{p} \sum_{i=0}^{r-1} \frac{1}{r(n-i)}. \quad (1)$$

for  $r = 1, 2, \dots, n-m$  where

$$m = \left( \frac{n}{\omega^{1/2} \log n} \right).$$

Using this we argue that

$$\mathbf{E}[C(G)] = C(n, n) = (C(n, n) - C(n, n-m+1)) + \frac{1+o(1)}{p} \sum_{r=1}^{n-m} \sum_{i=0}^{r-1} \frac{1}{r(n-i)}. \quad (2)$$

We will then show that

$$\sum_{r=1}^{n-m} \sum_{i=0}^{r-1} \frac{1}{r(n-i)} \approx \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}. \quad (3)$$

$$C(n, n) - C(n, n-m+1) = o(p^{-1}). \quad (4)$$

Theorem 1 follows from these two statements.

Let  $B_r = \{\phi_r(a_i) : i = 1, 2, \dots, r\}$ .

**Lemma 1.**  $B_r$  is a random  $r$ -subset of  $B$ .

*Proof.* Let  $L$  denote the  $n \times n$  matrix of edge costs, where  $L(i, j) = W(a_i, b_j)$  and  $L(i, j) = \infty$  if edge  $(a_i, b_j)$  does not exist in  $G$ . For a permutation  $\pi$  of  $B$  let  $L_\pi$  be defined by  $L_\pi(i, j) = L(i, \pi(j))$ . Let  $X, Y$  be two distinct  $r$ -subsets of  $B$  and let  $\pi$  be any permutation of  $B$  that takes  $X$  into  $Y$ . Then we have

$$\mathbf{Pr}(B_r(L) = X) = \mathbf{Pr}(B_r(L_\pi) = \pi(X)) = \mathbf{Pr}(B_r(L_\pi) = Y) = \mathbf{Pr}(B_r(L) = Y),$$

where the last equality follows from the fact that  $L$  and  $L_\pi$  have the same distribution.  $\square$

We use the above lemma to bound degrees. For  $v \in A$  let  $d_r(v) = |\{w \in B \setminus B_r : (v, w) \in E(G)\}|$ . Then we have the following lemma:

**Lemma 2.**

$$|d_r(v) - (n-r)p| \leq \omega^{-1/5}(n-r)p \text{ w.h.p. for } v \in A, 0 \leq r \leq n-m.$$

*Proof.* This follows from Lemma 1 i.e.  $B \setminus B_r$  is a random set and the Chernoff bounds viz.

$$\Pr(|d_r(v) - (n-r)p| \geq \omega^{-1/5}(n-r)p) \leq 2e^{-\omega^{-2/5}(n-r)p/3} \leq 2n^{-\omega^{1/10}}.$$

□

We can now use the ideas of [16], [17]. We add a special vertex  $b_{n+1}$  to  $B$ , with edges to all  $n$  vertices of  $A$ . Each edge adjacent to  $b_{n+1}$  is assigned an  $E(\lambda)$  cost independently,  $\lambda > 0$ . We now consider  $M_r$  to be a minimum cost matching of  $A_r$  into  $B^* = B \cup \{b_{n+1}\}$ . We denote this matching by  $M_r^*$  and we let  $B_r^*$  denote the corresponding set of vertices of  $B^*$  that are covered by  $M_r^*$ .

**Lemma 3.** *Suppose  $r < n - m$ . Then*

$$\Pr(b_{n+1} \in B_r^* \mid b_{n+1} \notin B_{r-1}^*) = \frac{\lambda}{p(n-r+1)(1+\varepsilon_r) + \lambda} \quad (5)$$

where  $|\varepsilon_r| \leq \omega^{-1/5}$ .

*Proof.* Assume that  $b_{n+1} \notin B_{r-1}^*$ .  $M_r^*$  is obtained from  $M_{r-1}^*$  by finding an augmenting path  $P = (a_r, \dots, a_\sigma, b_\tau)$  from  $a_r$  to  $B^* \setminus B_{r-1}^*$  of minimum additional cost. Let  $\alpha = W(\sigma, \tau)$ . We condition on (i)  $\sigma$ , (ii) the lengths of all edges other than  $(a_\sigma, b_j), b_j \in B^* \setminus B_{r-1}^*$  and (iii)  $\min\{A(\sigma, j) : b_j \in B^* \setminus B_{r-1}^*\} = \alpha$ . With this conditioning  $M_{r-1} = M_{r-1}^*$  will be fixed and so will  $P' = (a_r, \dots, a_\sigma)$ . We can now use the following fact: Let  $X_1, X_2, \dots, X_M$  be independent exponential random variables of rates  $\alpha_1, \alpha_2, \dots, \alpha_M$ . Then the probability that  $X_i$  is the smallest of them is  $\alpha_i/(\alpha_1 + \alpha_2 + \dots + \alpha_M)$ . Furthermore, the probability stays the same if we condition on the value of  $\min\{X_1, X_2, \dots, X_M\}$ . Thus

$$\Pr(b_{n+1} \in B_r^* \mid b_{n+1} \notin B_{r-1}^*) = \frac{\lambda}{d_{r-1}(a_\sigma) + \lambda}.$$

□

**Corollary 1.**

$$\Pr(b_{n+1} \in B_r^*) = \frac{1}{p} \left( \frac{1}{n} + \frac{1}{n-1} + \dots + \frac{1}{n-r+1} \right) (1 + \varepsilon_k)\lambda + O(\lambda^2) \quad (6)$$

as  $\lambda \rightarrow 0$ , where  $|\varepsilon_r| \leq \omega^{-1/5}$ .

*Proof.* Let  $\nu(j) = p^{-1}(n-j)(1+\varepsilon_j)$ ,  $|\varepsilon_j| \leq \omega^{-1/5}$ . Then the probability is given by

$$\begin{aligned} & 1 - \frac{\nu(0)}{\nu(0) + \lambda} \cdot \frac{\nu(1)}{\nu(1) + \lambda} \cdots \frac{\nu(r-1)}{\nu(r-1) + \lambda} \\ &= 1 - \left(1 + \frac{\lambda}{\nu(0)}\right)^{-1} \cdots \left(1 + \frac{\lambda}{\nu(r-1)}\right)^{-1} \\ &= \left(\frac{1}{\nu(0)} + \frac{1}{\nu(1)} + \dots + \frac{1}{\nu(r-1)}\right) \lambda + O(\lambda^2) \\ &= \frac{1}{p} \left(\frac{1}{n(1+\varepsilon_0)} + \frac{1}{(n-1)(1+\varepsilon_1)} + \dots + \frac{1}{(n-r+1)(1+\varepsilon_{r-1})}\right) \lambda + O(\lambda^2) \end{aligned}$$

and each error factor satisfies  $|1 - 1/(1+\varepsilon_j)| \leq \omega^{-1/5}$ .

□

**Lemma 4.** *If  $r \leq n - m$  then*

$$\mathbf{E}[C(n, r) - C(n, r - 1)] = \frac{1}{rp} \sum_{i=0}^{r-1} \frac{1 + \varepsilon_k}{n - i} \quad (7)$$

where  $|\varepsilon_k| \leq \omega^{-1/5}$ .

*Proof.* Let  $X$  be the cost of  $M_r$  and let  $Y$  be the cost of  $M_{r-1}$ . Let  $w$  denote the cost of the edge  $(a_r, b_{n+1})$ , and let  $I$  denote the indicator variable for the event that the cost of the cheapest  $A_r$ -assignment that contains this edge is smaller than the cost of the cheapest  $A_r$ -assignment that does not use  $b_{n+1}$ . In other words,  $I$  is the indicator variable for the event  $\{Y + w < X\}$ .

It follows from Corollary 1 and symmetry (to obtain the factor  $1/r$ ) that the probability that  $(a_r, b_{n+1}) \in M_r^*$  is given by

$$\frac{1}{rp} \left( \frac{1}{n} + \frac{1}{n-1} + \cdots + \frac{1}{n-r+1} \right) (1 + \varepsilon_k) \lambda + O(\lambda^2) \quad (8)$$

as  $\lambda \rightarrow 0$ , since each edge adjacent to  $b_{n+1}$  is equally likely to participate in  $M_{r-1}$ . If  $(a_r, b_{n+1}) \in M_r^*$  then  $w < X - Y$ . Conversely, if  $w < X - Y$  and no other edge from  $b_{n+1}$  has cost smaller than  $X - Y$ , then  $(a_r, b_{n+1}) \in M_r^*$ , and when  $\lambda \rightarrow 0$ , the probability that there are two distinct edges from  $b_{n+1}$  of cost smaller than  $X - Y$  is of order  $O(\lambda^2)$ .

On the other hand,  $w$  is  $E(\lambda)$  distributed, so

$$\mathbf{E}[I] = \Pr\{w < X - Y\} = \mathbf{E}\left[1 - e^{-\lambda(X-Y)}\right] = 1 - \mathbf{E}\left[e^{-\lambda(X-Y)}\right]. \quad (9)$$

Hence  $\mathbf{E}[I]$ , regarded as a function of  $\lambda$ , is essentially the Laplace transform of  $X - Y$ . In particular  $\mathbf{E}[X - Y]$  is the derivative of  $\mathbf{E}[I]$  evaluated at  $\lambda = 0$ , so

$$\mathbf{E}[X - Y] = \frac{d}{d\lambda} \mathbf{E}[I] \Big|_{\lambda=0} = \frac{1}{rp} \left( \frac{1}{n} + \frac{1}{n-1} + \cdots + \frac{1}{n-r+1} \right) (1 + \varepsilon_k) \quad (10)$$

where  $|\varepsilon_k| \leq \omega^{-1/5}$ . Now clearly, as  $\lambda \rightarrow 0$ ,  $\mathbf{E}[X] \rightarrow C(n, r)$  and  $\mathbf{E}[Y] = C(n, r - 1)$  and the lemma follows.  $\square$

This confirms (2) and we turn to (3). We use the following expression from Young [20].

$$\sum_{i=1}^n \frac{1}{i} = \log n + \gamma + \frac{1}{2n} + O(n^{-2}), \quad \text{where } \gamma \text{ is Euler's constant.} \quad (11)$$

Let  $m_1 = \omega^{1/4}m$ . Observe first that

$$\begin{aligned} \sum_{i=0}^{m_1} \frac{1}{n-i} \sum_{r=i+1}^{n-m} \frac{1}{r} &\leq o(1) + \sum_{i=n^{3/4}}^{m_1} \frac{1}{n-i} \sum_{r=i+1}^{n-m} \frac{1}{r} \leq \\ &o(1) + \frac{2}{n} \sum_{i=n^{3/4}}^{m_1} \left( \log \left( \frac{n}{i} \right) + \frac{1}{2(n-m)} + O(n^{-3/2}) \right) \leq o(1) + \frac{2}{n} \log \left( \frac{n^{m_1}}{m_1!} \right) \\ &\leq o(1) + \frac{2m_1}{n} \log \left( \frac{ne}{m_1} \right) = o(1). \end{aligned} \quad (12)$$

Then,

$$\begin{aligned}
\sum_{r=1}^{n-m} \sum_{i=0}^{r-1} \frac{1}{r(n-i)} &= \sum_{i=0}^{n-m-1} \frac{1}{n-i} \sum_{r=i+1}^{n-m} \frac{1}{r}, \\
&= \sum_{i=m_1}^{n-m-1} \frac{1}{n-i} \sum_{r=i+1}^{n-m} \frac{1}{r} + o(1), \\
&= \sum_{i=m_1}^{n-m-1} \frac{1}{n-i} \left( \log \left( \frac{n-m}{i} \right) + \frac{1}{2(n-m)} - \frac{1}{2i} + O(i^{-2}) \right) + o(1), \\
&= \sum_{i=m_1}^{n-m-1} \frac{1}{n-i} \log \left( \frac{n-m}{i} \right) + o(1), \\
&= \sum_{j=m+1}^{n-m_1} \frac{1}{j} \log \left( \frac{n-m}{n-j} \right) + o(1), \\
&= \int_{x=m+1}^{n-m_1} \frac{1}{x} \log \left( \frac{n-m}{n-x} \right) dx + o(1).
\end{aligned} \tag{13}$$

We can replace the sum in (1) by an integral because the terms are all  $o(1)$  and the sequence of summands is unimodal.

Continuing, we have

$$\begin{aligned}
&\int_{x=m+1}^{n-m_1} \frac{1}{x} \log \left( \frac{n-m}{n-x} \right) dx \\
&= - \int_{x=m+1}^{n-m_1} \frac{1}{x} \log \left( 1 - \frac{x-m}{n-m} \right) dx \\
&= \sum_{k=1}^{\infty} \int_{x=m+1}^{n-m_1} \frac{1}{x} \frac{(x-m)^k}{k(n-m)^k} dx \\
&= \int_{y=1}^{n-m-m_1} \frac{1}{y+m} \frac{y^k}{k(n-m)^k} dy.
\end{aligned} \tag{14}$$

Observe next that

$$\int_{y=1}^{n-m-m_1} \frac{1}{y+m} \frac{y^k}{k(n-m)^k} dy \leq \int_{y=0}^{n-m-m_1} \frac{y^{k-1}}{k(n-m)^k} dy \leq \frac{1}{k^2}.$$

So,

$$0 \leq \sum_{k=\log n}^{\infty} \int_{x=m+1}^{n-m_1} \frac{1}{x} \frac{(x-m)^k}{k(n-m)^k} dx \leq \sum_{k=\log n}^{\infty} \frac{1}{k^2} = o(1). \tag{15}$$

If  $1 \leq k \leq \log n$  then we write

$$\int_{y=1}^{n-m-m_1} \frac{1}{y+m} \frac{y^k}{k(n-m)^k} dy = \int_{y=1}^{n-m-m_1} \frac{(y+m)^{k-1}}{k(n-m)^k} dy + \int_{y=1}^{n-m-m_1} \frac{y^k - (y+m)^k}{(y+m)k(n-m)^k} dy.$$

Now

$$\int_{y=1}^{n-m-m_1} \frac{(y+m)^{k-1}}{k(n-m)^k} dy = \frac{1}{k^2} \frac{(n-m_1)^k - (m+1)^k}{(n-m)^k} = \frac{1}{k^2} + O \left( \frac{1}{k\omega^{1/4} \log n} \right). \tag{16}$$

If  $k = 1$  then

$$\int_{y=1}^{n-m-m_1} \frac{(y+m)^k - y^k}{(y+m)k(n-m)^k} dy = \frac{m \log(n-m_1)}{n-m} = o(1).$$

And if  $2 \leq k \leq \log n$  then

$$\begin{aligned} \int_{y=1}^{n-m-m_1} \frac{(y+m)^k - y^k}{(y+m)k(n-m)^k} dy &= \sum_{l=1}^k \int_{y=1}^{n-m-m_1} \binom{k}{l} \frac{y^{k-l} m^l}{(y+m)k(n-m)^k} dy \\ &\leq \sum_{l=1}^k \int_{y=0}^{n-m-m_1} \binom{k}{l} \frac{y^{k-l-1} m^l}{k(n-m)^k} dy \\ &= \sum_{l=1}^k \binom{k}{l} \frac{m^l (n-m-m_1)^{k-l}}{k(k-l)(n-m)^k} \\ &= O\left(\frac{km}{k(k-1)n}\right) = O\left(\frac{1}{k\omega^{1/2} \log n}\right) \end{aligned}$$

It follows that

$$0 \leq \sum_{k=1}^{\log n} \int_{y=1}^{n-m-m_1} \frac{(y+m)^k - y^k}{(y+m)k(n-m)^k} dy = o(1) + O\left(\sum_{k=2}^{\log n} \frac{1}{k\omega^{1/2} \log n}\right) = o(1). \quad (17)$$

equation (3) now follows from (14), (15), (16) and (17).

Turning to (4) we prove the following lemma:

**Lemma 5.** *If  $r \geq n - m$  then  $0 \leq C(n, r+1) - C(n, r) = O\left(\frac{\log n}{np}\right)$ .*

This will prove that

$$0 \leq C(n, n) - C(n - m + 1) = O\left(\frac{m \log n}{np}\right) = O\left(\frac{n}{\omega^{1/2} np}\right) = o\left(\frac{1}{p}\right)$$

and complete the proof of (4) and hence Theorem 1.

## 2.1 Proof of Lemma 5

Let  $w(e)$  denote the weight of edge  $e$  in  $G$ . Let  $V_r = A_{r+1} \cup B$  and let  $G_r$  be the subgraph of  $G$  induced by  $V_r$ . For a vertex  $v \in V_r$  order the neighbors  $u_1, u_2, \dots$ , of  $v$  in  $G_r$  so that  $w(v, u_i) \leq w(v, u_{i+1})$ . Define the  $k$ -neighborhood  $N_k(v) = \{u_1, u_2, \dots, u_k\}$ .

Let the  $k$ -neighborhood of a set be the union of the  $k$ -neighborhoods of its vertices. In particular, for  $S \subseteq A_{r+1}$ ,  $T \subseteq B$ ,

$$N_k(S) = \{b \in B : \exists a \in S : y \in N_k(a)\}, \quad (18)$$

$$N_k(T) = \{a \in A_{r+1} : \exists b \in T : a \in N_k(b)\}. \quad (19)$$

Given a function  $\phi$  defining a matching  $M$  of  $A_r$  into  $B$ , we define the following digraph: let  $\vec{\Gamma}_r = (V_r, \vec{X})$  where  $\vec{X}$  is an orientation of

$$X = \{\{a, b\} \in G : a \in A_{r+1}, b \in N_{40}(a)\} \cup \{\{a, b\} \in G : b \in B, a \in N_{40}(b)\} \cup \{(\phi(a_i), a_i) : i = 1, 2, \dots, r\}.$$



An edge  $e \in M$  is oriented from  $B$  to  $A$  and has weight  $w_r(e) = -w(e)$ . The remaining edges are oriented from  $A$  to  $B$  and have weight equal to their weight in  $G$ .

The arcs of directed paths in  $\vec{\Gamma}_r$  are alternately forwards  $A \rightarrow B$  and backwards  $B \rightarrow A$  and so they correspond to alternating paths with respect to the matching  $M$ . It helps to know (Lemma 6, next) that given  $a \in A_{r+1}, b \in B$  we can find an alternating path from  $a$  to  $b$  with  $O(\log n)$  edges. The  $ab$ -diameter will be the maximum over  $a \in A_{r+1}, b \in B$  of the length of a shortest path from  $a$  to  $b$ .

**Lemma 6.** *W.h.p., for every  $\phi$ , the (unweighted)  $ab$ -diameter of  $\vec{\Gamma}_r$  is at most  $k_0 = \lceil 3 \log_4 n \rceil$ .*

*Proof.* For  $S \subseteq A_{r+1}, T \subseteq B$ , let

$$\begin{aligned} N(S) &= \{b \in B : \exists a \in S \text{ such that } (a, b) \in \vec{X}\}, \\ N(T) &= \{a \in A_{r+1} : \exists b \in T \text{ such that } (a, b) \in \vec{X}\}. \end{aligned}$$

We first prove an expansion property: that **whp**, for all  $S \subseteq A_{r+1}$  with  $|S| \leq \lceil n/5 \rceil$ ,  $|N(S)| \geq 4|S|$ . (Note that  $N(S), N(T)$  involve edges oriented from  $A$  to  $B$  and so do not depend on  $\phi$ .)

$$\begin{aligned} \Pr(\exists S : |S| \leq \lceil n/5 \rceil, |N(S)| < 4|S|) &\leq o(1) + \sum_{s=1}^{\lceil n/5 \rceil} \binom{r+1}{s} \binom{n}{4s} \left( \frac{\binom{4s}{40}}{\binom{n}{40}} \right)^s \\ &\leq \sum_{s=1}^{\lceil n/5 \rceil} \left( \frac{ne}{s} \right)^s \left( \frac{ne}{4s} \right)^{4s} \left( \frac{4s}{n} \right)^{40s} \\ &= \sum_{s=1}^{\lceil n/5 \rceil} \left( \frac{e^5 4^{36} s^{35}}{n^{35}} \right)^s \\ &= o(1). \end{aligned} \tag{20}$$

**Explanation:** *The  $o(1)$  term accounts for the probability that each vertex has at least 40 neighbors in  $\vec{\Gamma}_r$ . Condition on this. Over all possible ways of choosing  $s$  vertices and  $4s$  “targets”, we take the probability that for each of the  $s$  vertices, all 40 out-edges fall among the  $4s$  out of the  $n$  possibilities.*

Similarly, w.h.p., for all  $T \subseteq B$  with  $|T| \leq \lceil n/5 \rceil$ ,  $|N(T)| \geq 4|T|$ . Thus by the union bound, w.h.p. both these events hold. In the remainder of this proof we assume that we are in this “good” case, in which all small sets  $S$  and  $T$  have large vertex expansion.

Now, choose an arbitrary  $a \in A_{r+1}$ , and define  $S_0, S_1, S_2, \dots$  as the endpoints of all alternating paths starting from  $a$  and of lengths  $0, 2, 4, \dots$ . That is,

$$S_0 = \{a\} \text{ and } S_i = \phi^{-1}(N(S_{i-1})).$$

Since we are in the good case,  $|S_i| \geq 4|S_{i-1}|$  provided  $|S_{i-1}| \leq n/5$ , and so there exists a smallest index  $i_S$  such that  $|S_{i_S-1}| > n/5$ , and  $i_S - 1 \leq \log_4(n/5) \leq \log_4 n - 1$ . Arbitrarily discard vertices from  $S_{i_S-1}$  to create a smaller set  $S'_{i_S-1}$  with  $|S'_{i_S-1}| = \lceil n/5 \rceil$ , so that  $S'_{i_S} = N(S'_{i_S-1})$  has cardinality  $|S'_{i_S}| \geq 4|S'_{i_S-1}| \geq 4n/5$ .

Similarly, for an arbitrary  $b \in B$ , define  $T_0, T_1, \dots$ , by

$$T_0 = \{b\} \text{ and } T_i = \phi(N(T_{i-1})).$$

Again, we will find an index  $i_T \leq \log_4 n$  whose modified set has cardinality  $|T'_{i_T}| \geq 4n/5$ .

With both  $|S'_{i_S}|$  and  $|T'_{i_T}|$  larger than  $n/2$ , there must be some  $a' \in S'_{i_S}$  for which  $b' = \phi(a') \in T'_{i_T}$ . This establishes the existence of an alternating walk and hence (removing any cycles) an alternating path of length at most  $2(i_S + i_T) \leq 2 \log_4 n$  from  $a$  to  $b$  in  $\vec{\Gamma}_r$ .  $\square$

We will need the following lemma,

**Lemma 7.** *Suppose that  $k_1 + k_2 + \dots + k_M \leq a \log N$ , and  $X_1, X_2, \dots, X_M$  are independent random variables with  $Y_i$  distributed as the  $k_i$ th minimum of  $N$  independent exponential rate one random variables. If  $\mu > 1$  then*

$$\Pr \left( X_1 + \dots + X_M \geq \frac{\mu a \log N}{N - a \log N} \right) \leq N^{a(1+\log \mu - \mu)}.$$

*Proof.* Let  $Y_{(k)}$  denote the  $k$ th smallest of  $Y_1, Y_2, \dots, Y_N$ , where we assume that  $k = O(\log N)$ . Then the density function  $f_k(x)$  of  $Y_{(k)}$  is

$$f_k(x) = \binom{N}{k} k (1 - e^{-x})^{k-1} e^{-x(N-k+1)} dx$$

and hence the  $i$ th moment of  $Y_{(k)}$  is given by

$$\begin{aligned} \mathbf{E} \left[ Y_{(k)}^i \right] &= \int_0^\infty \binom{N}{k} k x^i (1 - e^{-x})^{k-1} e^{-x(N-k+1)} dx \\ &\leq \int_0^\infty \binom{N}{k} k x^{i+k-1} e^{-x(N-k+1)} dx \\ &= \binom{N}{k} k \frac{(i+k-1)!}{(N-k+1)^{i+k}} \\ &\leq \left( 1 + O\left(\frac{k^2}{N}\right) \right) \frac{k(k+1) \cdots (i+k-1)}{(N-k+1)^i}. \end{aligned}$$

Thus, if  $0 \leq t < N - k + 1$ ,

$$\mathbf{E} \left[ e^{tY_{(k)}} \right] \leq \left( 1 + O\left(\frac{k^2}{N}\right) \right) \sum_{i=0}^\infty \left( \frac{-t}{N-k+1} \right)^i \binom{-k}{i} = \left( 1 + O\left(\frac{k^2}{N}\right) \right) \left( 1 - \frac{t}{N-k+1} \right)^{-k}.$$

If  $Z = X_1 + X_2 + \dots + X_M$  then if  $0 \leq t < N - a \log N$ ,

$$\mathbf{E} \left[ e^{tZ} \right] = \prod_{i=1}^M \mathbf{E} \left[ e^{tX_i} \right] \leq \left( 1 - \frac{t}{N - a \log N} \right)^{-a \log N}.$$

It follows that

$$\Pr \left( Z \geq \frac{\mu a \log N}{N - a \log N} \right) \leq \left( 1 - \frac{t}{N - a \log N} \right)^{-a \log N} \exp \left\{ -\frac{t \mu a \log N}{N - a \log N} \right\}.$$

We put  $t = (N - a \log N)(1 - 1/\mu)$  to minimise the above expression, giving

$$\Pr \left( Z \geq \frac{\mu a \log N}{N - a \log N} \right) \leq (\mu e^{1-\mu})^{a \log N}.$$

$\square$

**Lemma 8.** *W.h.p., for all  $\phi$ , the weighted ab-diameter of  $\bar{\Gamma}_r$  is at most  $c_1 \frac{\log n}{np}$  for some absolute constant  $c_1 > 0$ .*

*Proof.* Let

$$Z_1 = \max \left\{ \sum_{i=0}^k w(a_i, b_i) - \sum_{i=0}^{k-1} w(b_i, a_{i+1}) \right\}, \quad (21)$$

where the maximum is over sequences  $a_0, b_0, a_1, \dots, a_k, b_k$  where  $(a_i, b_i)$  is one of the 40 shortest arcs leaving  $a_i$  for  $i = 0, 1, \dots, k \leq k_0 = \lceil 3 \log_4 n \rceil$ , and  $(b_i, a_{i+1})$  is a backwards matching edge.

We compute an upper bound on the probability that  $Z_1$  is large. For any  $\zeta > 0$  we have

$$\Pr \left( Z_1 \geq \zeta \frac{\ln n}{np} \right) \leq o(1) + \sum_{k=0}^{k_0} ((r+1)n)^{k+1} \left( \frac{1+o(1)}{np} \right)^{k+1} \times \int_{y=0}^{\infty} \left[ \frac{1}{(k-1)!} \left( \frac{y \ln n}{np} \right)^{k-1} \sum_{\rho_0 + \rho_1 + \dots + \rho_k \leq 40(k+1)} q(\rho_0, \rho_1, \dots, \rho_k; \zeta + y) \right] dy$$

where

$$q(\rho_0, \rho_1, \dots, \rho_k; \eta) = \Pr \left( X_0 + X_1 + \dots + X_k \geq \eta \frac{\log n}{np} \right),$$

$X_0, X_1, \dots, X_k$  are independent and  $X_j$  is distributed as the  $\rho_j$ th minimum of  $r$  independent exponential random variables. (When  $k = 0$  there is no term  $\frac{1}{(k-1)!} \left( \frac{y \log n}{n} \right)^{k-1}$ ).

**Explanation:** *The  $o(1)$  term is for the probability that there is a vertex in  $V_r$  that has fewer than  $(1 - o(1))np$  neighbors in  $V_r$ . We have at most  $((r+1)n)^{k+1}$  choices for the sequence  $a_0, b_0, a_1, \dots, a_k, b_k$ . The term  $\frac{1}{(k-1)!} \left( \frac{y \ln n}{np} \right)^{k-1} dy$  bounds the probability that the sum of  $k$  independent exponentials,  $w(b_0, a_1) + \dots + w(b_{k-1}, a_k)$ , is in  $\frac{\ln n}{np}[y, y + dy]$ . (The density function for the sum of  $k$  independent exponentials is  $\frac{x^{k-1} e^{-x}}{(k-1)!}$ .) We integrate over  $y$ .*

*$\frac{1+o(1)}{np}$  is the probability that  $(a_i, b_i)$  is the  $\rho_i$ th shortest edge leaving  $a_i$ , and these events are independent for  $0 \leq i \leq k$ . The final summation bounds the probability that the associated edge lengths sum to at least  $\frac{(\zeta+y) \ln n}{np}$ .*

It follows from Lemma 7 that if  $\zeta$  is sufficiently large then, for all  $y \geq 0$ ,

$$q(\rho_1, \dots, \rho_k; \zeta + y) \leq (np)^{-(\zeta+y) \log n / (2 \log np)} = n^{-(\zeta+y)/2}.$$

Since the number of choices for  $\rho_0, \rho_1, \dots, \rho_k$  is at most  $\binom{41k+40}{k+1}$  (the number of positive integral solutions to  $a_0 + a_1 + \dots + a_{k+1} \leq 40(k+1)$ ) we have

$$\begin{aligned} \Pr \left( Z_1 \geq \zeta \frac{\ln n}{np} \right) &\leq 2n^{2-\zeta/2} \sum_{k=0}^{k_0} \frac{(\ln n)^{k-1}}{(k-1)!} \binom{41k+40}{k+1} \int_{y=0}^{\infty} y^{k-1} n^{-y/2} dy \\ &\leq 2n^{2-\zeta/2} \sum_{k=0}^{k_0} \frac{(\ln n)^{k-1}}{(k-1)!} \left( \frac{84e}{\ln n} \right)^k \Gamma(k) \\ &\leq 2n^{2-\zeta/2} (k_0 + 1) (84e)^{k_0+2} \\ &= o(n^{-4}). \end{aligned}$$

□

Lemma 8 shows that with probability  $1 - o(n^{-2})$  in going from  $M_r$  to  $M_{r+1}$  we can find an augmenting path of weight at most  $\frac{c_1 \log n}{np}$ . This completes the proof of Lemma 5 and Theorem 1. (Note that to go from w.h.p. to expectation we use the fact that w.h.p.  $w(e) = O(\log n)$  for all  $e \in A \times B$ .)  $\square$

Notice that in the proof of Lemmas 6 and 8 we can certainly make the failure probability less than  $n^{-\text{anyconstant}}$ .

### 3 Proof of Theorem 2

The proof of Lemma 8 allows us to claim that with probability  $1 - O(n^{-\text{anyconstant}})$  the maximum length of an edge in the minimum cost perfect matching of  $G$  is at most  $\mu = c_2 \frac{\log n}{np}$  for some constant  $c_2 > 0$ . We can now proceed as in Talagrand's proof of concentration for the assignment problem. We let  $\hat{w}(e) = \min\{w(e), \mu\}$  and let  $\hat{C}(G)$  be the assignment cost using  $\hat{w}$  in place of  $w$ . We observe that

$$\Pr(\hat{C}(G) \neq C(G)) = O(n^{-\text{anyconstant}}) \quad (22)$$

and so it is enough to prove concentration of  $\hat{C}(G)$ .

For this we use the following result of Talagrand [14]: consider a family  $\mathcal{F}$  of  $N$ -tuples  $\alpha = (\alpha_i)_{i \leq N}$  of non-negative real numbers. Let

$$Z = \min_{\alpha \in \mathcal{F}} \sum_{i \leq N} \alpha_i X_i$$

where  $X_1, X_2, \dots, X_N$  are an independent sequence of random variables taking values in  $[0, 1]$ .

Let  $\sigma = \max_{\alpha \in \mathcal{F}} \|\alpha\|_2$ . Then if  $M$  is the median of  $Z$  and  $u > 0$ , we have

$$\Pr(|Z - M| \geq u) \leq 4 \exp \left\{ -\frac{u^2}{4\sigma^2} \right\}. \quad (23)$$

We apply (23) with  $N = n^2$  and  $X_e = \hat{w}(e)/\mu$ . For  $\mathcal{F}$  we take the  $n!$   $\{0, 1\}$  vectors corresponding to perfect matchings and scale them by  $\mu$ . In this way,  $\sum_e \alpha_e X_e$  will be the weight of a perfect matching. In this case we have  $\sigma^2 \leq n\mu^2$ . Applying (23) we obtain

$$\Pr \left( |\hat{C}(G) - \widehat{M}| \geq \frac{\varepsilon}{p} \right) \leq 4 \exp \left\{ -\frac{\varepsilon^2}{4p^2} \cdot \frac{1}{n\mu^2} \right\} = \exp \left\{ -\frac{\varepsilon^2 n}{(c_2 \log n)^2} \right\}, \quad (24)$$

where  $\widehat{M}$  is the median of  $\hat{C}(G)$ . Theorem 2 follows easily from (22) and (24).

### 4 Final remarks

We have generalised the result of [2] to random bipartite graphs. It would seem that in the absence of proving Conjecture 1 we should be able to replace  $\omega(\log n)^2$  in Theorem 1 by  $\omega \log n$ . It would be of interest to prove the analogous result for  $G_{n,p}$ . Here we would expect to find that the expected cost of a minimum matching was asymptotically  $\frac{\pi^2}{12p}$ , given that Wästlund has proved this for  $p = 1$  in [15].

## References

- [1] D. Aldous, Asymptotics in the random assignment problem, *Probability Theory and Related Fields* 93 (1992) 507-534.
- [2] D. Aldous, The  $\zeta(2)$  limit in the random assignment problem, *Random Structures and Algorithms* 4 (2001) 381-418.
- [3] J. Beardwood, J. H. Halton and J. M. Hammersley, The shortest path through many points, *Mathematical Proceedings of the Cambridge Philosophical Society* 55 (1959) 299-327.
- [4] A. Beveridge, A.M. Frieze and C.McDiarmid, Random minimum length spanning trees in regular graphs, *Combinatorica* 18 (1998) 311-333.
- [5] C. Cooper, A.M. Frieze, N. Ince, S. Janson and J. Spencer, On the length of a random minimum spanning tree, see arxiv.org.
- [6] A.M. Frieze, On the value of a random minimum spanning tree problem, *Discrete Applied Mathematics* 10 (1985) 47-56.
- [7] A.M. Frieze, M. Ruszinko and L. Thoma, A note on random minimum length spanning trees, *Electronic Journal of Combinatorics* 7 (2000) R41.
- [8] S. Janson, The minimal spanning tree in a complete graph and a functional limit theorem for trees in a random graph, *Random Structures Algorithms* 7 (1995) 337-355.
- [9] S. Janson, One, two and three times  $\log n/n$  for paths in a complete graph with random weights, *Combinatorics, Probability and Computing* 8 (1999) 347-361.
- [10] R.M. Karp, An upper bound on the expected cost of an optimal assignment, *Discrete Algorithms and Complexity: Proceedings of the Japan-US Joint Seminar (D. Johnson et al., eds.)*, Academic Press, New York, 1987, 1-4.
- [11] S. Linusson and J. Wästlund, A proof of Parisi's conjecture on the random assignment problem, *Probability Theory and Related Fields* 128 (2004) 419-440.
- [12] C. Nair, B. Prabhakar and M. Sharma, Proofs of the Parisi and Coppersmith-Sorkin random assignment conjectures, *Random Structures and Algorithms* 27 (2005) 413-444.
- [13] G. Parisi, A conjecture on Random Bipartite Matching, Physics e-Print archive (1998).
- [14] M. Talagrand, Concentration of measure and isoperimetric inequalities in product spaces, *Publications Mathématiques de L'IHÉS* (1995).
- [15] J. Wästlund, Random matching problems on the complete graph, *Electronic Communications in Probability* (2008) 258-265.
- [16] J. Wästlund, A simple proof of the Parisi and Coppersmith-Sorkin formulas for the random assignment problem, *Linköping Studies in Mathematics* 6 (2005).
- [17] J. Wästlund, An easy proof of the  $\zeta(2)$  limit in the random assignment problem, *Elec. Comm. in Probab.* 14 (2009) 261 – 269

- [18] J. Michael Steele, Probability Theory and Combinatorial Optimization, SIAM CBMS series, 1996.
- [19] D.W. Walkup, On the expected value of a random assignment problem, *SIAM Journal on Computing* 8 (1979) 440-442.
- [20] R.M. Young, Euler's constant, *Mathematical Gazette* 75 (1991) 187-190.