

7-2015

Cops and robbers on planar directed graphs

Po-Shen Loh

Carnegie Mellon University, ploh@cmu.edu

Siyoung Oh

Daum Kakao Corp.

Follow this and additional works at: <http://repository.cmu.edu/math>

 Part of the [Mathematics Commons](#)

This Working Paper is brought to you for free and open access by the Mellon College of Science at Research Showcase @ CMU. It has been accepted for inclusion in Department of Mathematical Sciences by an authorized administrator of Research Showcase @ CMU. For more information, please contact research-showcase@andrew.cmu.edu.

Cops and robbers on planar directed graphs

Po-Shen Loh *

Siyoung Oh †

Abstract

Aigner and Fromme initiated the systematic study of the *cop number* of a graph by proving the elegant and sharp result that in every connected planar graph, three cops are sufficient to win a natural pursuit game against a single robber. This game, introduced by Nowakowski and Winkler, is commonly known as *Cops and Robbers* in the combinatorial literature. We extend this study to directed planar graphs, and establish separation from the undirected setting. We exhibit a geometric construction which shows that a more sophisticated robber strategy can indefinitely evade three cops on a particular strongly connected planar directed graph.

1 Introduction

The general study of pursuit games on graphs drew a substantial amount of research attention over the last decade. Its appeal stemmed from the combination of its apparent proximity to natural applications, some combinatorially elegant results and conjectures, and the challenge of developing tools to analyze game-theoretic dynamics on graphs. Indeed, dynamic processes are typically already significantly more difficult to analyze than properties of static graphs, and game theoretic interactions between opposing parties drive the complexity to another level.

This paper considers the most extensively studied game in this area, commonly known as *Cops and Robbers*, introduced by Nowakowski and Winkler [16], and independently by Quillot [18]. In its classical setting, a graph is fixed, and fully known to two players, *the cops* and *the robber*. The cops move first, placing k cops on the vertices of the given graph, at any locations of choice (multiple cops are allowed to reside on the same vertex). The robber then chooses a single vertex at which to start. Players alternate turns, starting with the cops, and on each turn, they choose a subset of their agents to move across one edge each. Note that the robber has only one agent, and so decides whether or not to move to an adjacent vertex. If the robber ends up on the same vertex as a cop, then the cops win.

The main question is to determine, for each graph, the minimum value of k (known as the *cop number* of the graph) for which there is a strategy for the cops that guarantees a win within finite time. This game-theoretic graph invariant was introduced by Aigner and Fromme [1] shortly after the game's appearance in the combinatorial literature, and in that same paper, the authors proved the elegant and sharp result that every planar graph has cop number at most three.

*Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, PA 15213. E-mail: ploh@cmu.edu. Research supported in part by an NSA Young Investigators Grant, NSF grants DMS-1201380 and DMS-1455125, and by a USA-Israel BSF Grant.

†Daum Kakao Corp., South Korea. Email: unbing@gmail.com. Research conducted for Masters Thesis at Carnegie Mellon University.

This basic setting is a natural prototype for a general class of pursuit games on graphs, and it has been the subject of numerous papers, including multiple surveys [3, 4, 8, 12] and ultimately a book by Bonato and Nowakowski [7]. Many variants have been studied, including random graphs [6, 15, 17], Cayley graphs [10], geometric graphs [5], directed graphs [11], and fast robbers [2], to name just a few.

The central open conjecture in this area, due to Meyniel (communicated by Frankl [9]), is that every n -vertex graph has cop number at most $O(\sqrt{n})$, which would be asymptotically tight. The current best bounds of $O(n/e^{\Theta(\sqrt{\log n})})$ were proven by Lu and Peng [14], with alternate proofs independently discovered by Frieze, Krivelevich, and Loh [11], and Scott and Sudakov [19].

The same paper of Frieze, Loh, and Krivelevich also formally started the systematic study of the game in directed graphs (where the cops and robber can only move along the direction of each edge), mainly in the context of Meyniel’s conjecture. Specifically, the focus was on n -vertex strongly connected digraphs, because the problem for a general digraph easily reduces to the problems on its strongly connected components. As usual, digraphs turn out to be more complicated than undirected graphs, and they obtained a weaker upper bound of $O(n \cdot \frac{(\log \log n)^2}{\log n})$, which is still the current best bound for the cop number in directed graphs. The lower bound from undirected graphs carries over to the directed case (simply replace each edge with a pair of antiparallel directed edges), but there was no improvement.

This paper combines the directed graph inquiry with the original focus of Aigner and Fromme on planar graphs. Specifically, we ask to determine, for each n , the maximum cop number of any n -vertex planar digraph. Unfortunately, the approach of Aigner and Fromme for their upper bound (of three cops) completely breaks down, as it relied on repeated clever applications of the following simple and elegant observation.

Lemma 1.1. *In an undirected graph, if P is a geodesic (a shortest path between a pair of vertices), then a single cop can guard all of the vertices of P : after a bounded number of turns, if the robber ever moves onto a vertex of P , it will be captured by that cop.*

Geodesics are particularly useful in planar graphs because they can provide powerful separation properties in the plane, thereby efficiently trapping the robber in successively smaller regions. The proof of Lemma 1.1 employs the cop strategy of always moving towards the vertex of P which is nearest to the robber. The geodesic’s minimality guarantees that the cop will eventually be able to reach every vertex of P at least as quickly as the robber. However, this strategy is clearly impossible in directed graphs, and indeed, there is no upper bound written in the literature. For completeness, we observe that the Lipton-Tarjan Planar Separator Theorem gives a nontrivial upper bound, but it is still far from constant. The proof will appear in Section 4.

Proposition 1.1. *Every n -vertex strongly connected planar digraph has cop number at most $O(\sqrt{n})$.*

The main open question for planar digraphs is whether the cop number is always bounded by a constant in each strongly connected component. It is then interesting to examine lower bound constructions. Again, basic methods do not work, as many previous results (including the tightness of Meyniel’s conjecture) relied on another simple observation.

Lemma 1.2. *Every undirected graph with minimum degree δ and no 3- or 4-cycles has cop number at least δ .*

This follows from the elementary robber strategy of remaining stationary until a cop moves to an adjacent vertex, and then moving to another adjacent vertex which has no cops adjacent to it. This lemma obviously extends to directed graphs with a similar strategy: every digraph with minimum out-degree δ^+ and an appropriate girth condition (e.g., undirected girth at least five) has cop number at least $1 + \delta^+$. It suffices to focus on constructions in which each unordered pair of vertices induces at most one directed edge, because if antiparallel edge pairs exist, the entire digraph can be replaced by one with no antiparallel pairs by subdividing every edge with a new unique vertex, and replacing each antiparallel edge pair by a pair of independent 2-edge directed paths. Since every n -vertex planar graph has fewer than $3n$ edges, any such construction will always have $\delta^+ \leq 2$, and so the standard robber strategy cannot even be used to improve the lower bound by any amount at all.

The main contribution of this paper is a geometrically-inspired construction which introduces and employs a more sophisticated strategy for the robber, and breaks through the lower bound for undirected graphs.

Theorem 1.1. *There is a strongly connected planar digraph which requires more than three cops to capture the robber.*

The rest of this paper is organized as follows. The construction is described in the next section, and analyzed in the section thereafter. We close with a short proof of Proposition 1.1 in the final section.

2 Construction

We will create a geometric construction which is clearly embeddable on the surface of a sphere, at which point a standard stereographic projection produces a planar oriented graph. Although it is impossible to construct a planar oriented graph with all out-degrees at least 3, it is natural to start with an object which is as close as possible. Indeed, consider an icosahedron (Figure 1), which as an undirected graph is a triangulation with all degrees equal to 5. We use this as the starting point for a series of steps which ultimately produce our construction.




Figure 1: Icosahedron




Figure 2: Truncated icosahedron

Truncate each of its vertices to obtain the truncated icosahedron in Figure 2. Observe that each vertex is replaced with a pentagon, and each of the triangular faces is replaced by a hexagon.

Next, truncate again, this time along all edges of the original icosahedron (which are now precisely the edges between pentagons). This operation produces the Archimedean solid in Figure 3, which is known as the *great rhombicosidodecahedron*, or *truncated icosidodecahedron*.¹

Observe that the original icosahedron vertices have been replaced by decagons, the original icosahedron edges have been replaced by quadrilaterals, and the original icosahedron faces have been replaced by hexagons. Also observe that each quadrilateral links two decagons, and naturally identifies a pair of parallel edges between decagons, as highlighted in Figure 4.




Figure 3: Great rhombicosidodecahedron




Figure 4: Highlighting parallel edges

We use this fundamental structure as the backbone for our construction. Introduce a new vertex at the center of each decagon, called a *center*, and join it to each vertex of its decagon with a *spoke*, as in Figure 5. We will use the term *unit* to refer to the whole structure of a single decagon, including its center and spokes. Two units are *neighbors* if they are joined by one of the highlighted edges in Figure 4.

The last step is to give directions and lengths to all edges. Observe that between each pair of neighboring units, we can always find a hexagon as highlighted in Figure 6. In each such hexagon, orient all edges counter-clockwise. This orientation is consistent because the sphere is an orientable surface. Ultimately, we obtain the structure of Figure 7.

All green edges between units are now consistently oriented. Subdivide each of them with 999 new vertices so that it takes 1,000 turns to move from one end of a green edge to the other end. Likewise, subdivide each spoke between a center and its decagon with 9 new vertices so that each spoke now has length 10. The only edges remaining to be oriented are the decagon edges. Replace each of them with a directed chain-like structure as in Figure 8. It now takes 16 turns to move from one original decagon vertex to another, and at most one additional turn to reverse direction when traversing the chain.

3 Analysis

In this section, we prove that our construction requires more than three cops to capture the robber. We achieve this by analyzing the robber's travel from unit to unit. Note that each unit has five neighboring units, and in order to travel from a unit U_1 to a neighboring unit U_2 , there is exactly

¹All polyhedron images were generated by <http://gratrix.net/polyhedra/webgl/poly.xhtml>.




Figure 5: Point of view from any unit (without directions on edges). Note that edges connecting units are green.




Figure 6: Point of view from any unit, highlighting one hexagon.




Figure 7: Point of view from any unit, with directions determined.




Figure 8: Replacement for one original decagon edge.

one *exit* vertex on the perimeter of U_1 from which the robber can directly travel along a directed path of length 1000 to reach U_2 . Each unit therefore has five exits. It is convenient to make the following observation.

Lemma 3.1. *Suppose that the robber is at the center of a unit U , and it is the robber's turn. Let c be the number of cops in unit U which are not on spokes that are oriented towards the center. Then, for any set S of more than c exits of U , there is at least one vertex in S that the robber can reach in 10 turns, without being captured by any of the cops currently in U .*

Proof. The robber will move directly toward one of the vertices of S . It is clear that with this strategy, the robber cannot be stopped by any cop who is currently on a spoke oriented towards the center. Furthermore, since each decagon edge was replaced by a chain which takes 16 turns to traverse from end to end, it is clear that every other cop in U can only be within 10 moves of at most one vertex of S . Since $|S| > c$, there will be a choice for the robber which avoids all of the cops. \square

We will ultimately break into cases based upon how many cops are in the robber's current unit. It turns out that the most delicate case is when there is exactly one cop in the robber's unit, and the following observation will cleanly handle that situation, in conjunction with the previous result.

Lemma 3.2. *Suppose that the robber is on the perimeter of a unit, and that unit has exactly one cop, located one vertex away from the center along a spoke which is oriented away from the center. Then, the robber can reach the center within 27 moves, without that cop catching it.*

Proof. The robber will follow a shortest directed path from its current location to the center. Since the spokes alternate in orientation toward and away from the center, and it only takes at most one turn to change direction along a chain on the perimeter, it is clear that the robber will reach the center without any interference from the cop, who starts 9 moves away from the perimeter along an outward spoke. The maximum number of moves required for the robber is 27, because it may take the robber 1 move to change direction along a perimeter chain, 16 moves to traverse the perimeter to the nearest inward spoke, and 10 more moves to traverse the inward spoke to the center. \square

We are now ready to prove that three cops are insufficient to capture the robber on our construction.

Proof of Theorem 1.1. At the start of the game, three cops select their positions. Since the icosahedron has 12 vertices, the robber is free to select a position which is the center of a unit that starts with no cops. The analysis now proceeds by considering how the robber moves from unit to neighboring unit.

The robber remains at a center until a cop arrives at an adjacent vertex (along an inbound spoke in its unit). It is then the robber's turn. It suffices to show that the robber can always move from this state to a state in which it is again at a center (possibly of a different unit), with a cop adjacent along an inbound spoke, and it is the robber's turn. This will prove that the robber can escape capture indefinitely.

So, let us focus on the situation in which the robber is at a center of some unit U , with an immediately adjacent inbound cop. If there are any cops on length-1000 unit-to-unit directed paths towards U , without loss of generality, assume that they are already at the corresponding entry vertices along the perimeter of U . (This only makes it more difficult for the robber to escape to a neighboring unit center, because cops that are trapped in transit along a unit-to-unit path have nowhere to go but forward to U .) Note that there are five exit vertices from U , each to a distinct neighboring unit. The robber's strategy will be to use one of them to exit to a neighboring unit which cannot be reached more quickly by any cop.

We split into three cases, depending on how many cops are in unit U . If all three cops are in U , then Lemma 3.1 implies that of the five exits from U , there is at least one option which can be reached with no interference from the cops. The robber moves directly toward one of these, takes the length-1000 directed path to the neighboring unit, and then moves directly toward that unit's center, completing this case.

Next, consider the case when there are exactly two cops in U . The cop outside U is in some unit U' , but even if one considers U' together with its five neighboring units, these six total units overlap with at most three neighboring units of U . (We have used the fact that in an icosahedron, among the five neighbors of a fixed vertex u , the overlap size with a different vertex u' and its neighborhood is largest when u' is a neighbor of u , at which point it has size three.) The robber will seek an exit which does not go to U' or a neighboring unit of U' . Since there were five exits, there are still at least two left. Lemma 3.1 then implies that there is at least one option which can be reached with no interference from the cops in U , and the robber safely proceeds through that exit to the center of that neighboring unit.

The final case has exactly one cop in U . The previous argument no longer works, because each of the two cops outside U can in theory block up to three neighboring units of U (as in the previous case's analysis), and could together block all five neighboring units of U . The robber counters with a different strategy. It is this twist in this case which improves the lower bound from three to four, and here we leverage the length-1000 paths between neighboring units. The key insight is that as long as no other cops start moving toward U , the robber can evade the single cop in U indefinitely. However, the moment a cop starts down a long one-way street towards U , it stops guarding three neighboring units, and in fact guards zero neighboring units. We formalize this as follows.

The robber begins by moving directly along an outbound spoke, and reaches a vertex on the perimeter of unit U in 10 moves. It then stays stationary until the cop in U moves onto an adjacent vertex. It then moves along the perimeter of U , in a direction away from the cop, moving only when the cop moves onto an adjacent vertex along an edge directed towards it. (So, it is possible that the robber spends a substantial amount of time not moving at all, if the cop is moving through vertices which are not adjacent to the robber.)

The robber continues this simple evasion strategy until one of the other two cops takes a step into a length-1000 unit-to-unit directed path leading to U . At this point, the robber switches strategy again, to exit U within 200 turns. To achieve this, observe that the remaining cop who is neither in U nor trapped in the length-1000 path to U can reach at most three neighboring units

of U within 1500 turns. Let S be the set of exits of U which do not lead to those units.

The robber selects a direction away from the cop in U , and consistently moves along the perimeter in that direction until he reaches one of the (at least two) exits in S . If during this process, the cop never passes through the center, then the robber will definitely reach its exit without being captured by the cop in U . Since each decagon edge was replaced by a 16-turn chain, this will take at most 160 turns.

Otherwise, if the cop attempts to route through the center, the robber suddenly changes strategy again at the moment the cop moves onto a vertex adjacent to the center along an outbound spoke (which must happen on any route through the center). At that moment, the robber employs the strategy in Lemma 3.2, and reaches the center within 27 more moves without interference from that cop. Then, the robber changes strategy again to that in Lemma 3.1, and since $|S| > 1$, it will definitely be able to reach an exit in S in 10 more turns, without any interference from the cop in U . Therefore, the robber will be able to reach an exit in S within a total of 200 turns, during which the cop en route from a neighboring unit (along a length-1000 directed path) is still far off. The robber then traverses the length-1000 directed path out of this exit, and reaches the center of the corresponding neighboring unit within 1500 turns, without any interference from any cops, as claimed. \square

4 Upper bound

We close with a very short treatment of the upper bound, which applies the Planar Separator Theorem of Lipton and Tarjan [13].

Theorem 4.1. [*Lipton and Tarjan CITE*] *There is an absolute constant c for which the following holds. Every n -vertex planar graph can be partitioned into three sets A , B , and C such that $|A| \leq 2n/3$, $|B| \leq 2n/3$, $|C| \leq c\sqrt{n}$, and there is no edge between A and B .*

Proof of Theorem 1.1. Initially, put $k\sqrt{n}$ cops at an arbitrary vertex, and call them *free* cops. The constant k will be determined at the end of the proof. By continuously separating G with these cops, we will show that the cops win the game. Let $G_1 = G$, and for $i \geq 1$, construct G_{i+1} with the following algorithm.

Using Lemma 4.1, let $G_i = A_i \cup B_i \cup C_i$ such that $|A_i| \leq 2|G_i|/3$, $|B_i| \leq 2|G_i|/3$, $|C_i| \leq c\sqrt{|G_i|}$, and there is no edge in G_i between A_i and B_i . Send free cops to each vertex of C_i without regard to the robber's actions. Note that this is always possible because G is strongly connected. This step costs at most $c\sqrt{|G_i|}$ free cops and these cops will not move afterward, thus becoming *static* cops. If the robber was not caught during this process, the robber is now permanently trapped in either A_i or B_i . Set G_{i+1} to be the graph induced by the corresponding set. Thus, $|G_1| = n$, $|G_2| \leq 2n/3$, \dots , $|G_i| \leq (\frac{2}{3})^{i-1}n$. The i -th step costs $c\sqrt{|G_i|}$ free cops. Thus, the number of cops needed to win is at most

$$c \sum_{i=0}^{\infty} \sqrt{\left(\frac{2}{3}\right)^i n} = c\sqrt{n} \sum_{i=0}^{\infty} \left(\sqrt{\frac{2}{3}}\right)^i,$$

which is at most $k\sqrt{n}$ for some k , because the sum is a convergent geometric series. \square

References

- [1] M. Aigner and M. Fromme, A game of cops and robbers, *Discrete Appl. Math.* **8** (1984), 1–12.

- [2] N. Alon and A. Mehrabian, Chasing a fast robber on planar graphs and random graphs, *J. Graph Theory* **78** (2015), 81–96.
- [3] B. Alspach, Searching and sweeping graphs, *Le Matematiche* **59** (2004), 5–37.
- [4] W. Baird and A. Bonato, Meyniel’s conjecture on the cop number: a survey, *J. Comb.* **3** (2012), 225–238.
- [5] A. Beveridge, A. Dudek, A. Frieze, and T. Müller, Cops and robbers on geometric graphs. *Combin. Probab. Comput.* 21 (2012), no. 6, 816834.
- [6] B. Bollobás, G. Kun, and I. Leader, Cops and robbers in a random graph, *Journal of Combinatorial Theory, Series B* **103** (2013), 226–236.
- [7] A. Bonato and R. Nowakowski, **The Game of Cops and Robbers on Graphs**, American Mathematical Society Student Mathematical Library vol. 61 (2011).
- [8] F. Fomin and D. Thilikos, An annotated bibliography in guaranteed graph searching, *Theoret. Comput. Sci.* **399** (2008), 236–245.
- [9] P. Frankl, Cops and robbers in graphs of large girth and Cayley graphs, *Discrete Appl. Math.* **17** (1987), 301–305.
- [10] P. Frankl, On a pursuit game on Cayley graphs, *Combinatorica* **7** (1987), 67–70.
- [11] A. Frieze, M. Krivelevich, and P. Loh, Variations on Cops and Robbers, *J. Graph Theory* **69** (2012), 383–402.
- [12] G. Hahn, Cops, robbers, and graphs, *Tatra Mt. Math. Publ.* **36** (2007), 163–176.
- [13] R. Lipton and R. Tarjan, *A separator theorem for planar graphs*, *SIAM J. Appl. Math.* 36 (1979), 177–189.
- [14] L. Lu and X. Peng, On Meyniel’s conjecture of the cop number. *J. Graph Theory* **71** (2012), 192–205.
- [15] T. Łuczak and P. Prałat, Chasing robbers on random graphs: zigzag theorem, *Random Structures and Algorithms* **37** (2010), 516–524.
- [16] R. Nowakowski and P. Winkler, Vertex to vertex pursuit in a graph, *Discrete Math.* **43** (1983), 235–239.
- [17] P. Prałat and N. Wormald, Meyniel’s conjecture holds for random graphs, *Random Structures and Algorithms*, to appear.
- [18] A. Quilliot, A short note about pursuit games played on a graph with a given genus, *J. Combin. Theory Ser. B* **38** (1985), 89–92.
- [19] A. Scott and B. Sudakov, A bound for the cops and robbers problem, *SIAM J. Discrete Math.* **25** (2011), 1438–1442.